

CATALOGUE
OF THE
MESOZOIC PLANTS
IN THE
DEPARTMENT OF GEOLOGY.
PART V.

CATALOGUE
OF THE
MESOZOIC PLANTS
IN THE
BRITISH MUSEUM
(NATURAL HISTORY).
THE CRETACEOUS FLORA.

PART I.—BIBLIOGRAPHY, ALGÆ & FUNGI.

BY
MARIE C. STOPES, D.Sc. (Lond.), Ph.D. (Munich), F.L.S.,
FELLOW OF UNIVERSITY COLLEGE; LECTURER IN PALEOBOTANY
AT UNIVERSITY COLLEGE, LONDON UNIVERSITY.


LONDON:
PRINTED BY ORDER OF THE TRUSTEES OF THE BRITISH MUSEUM.

SOLD BY
LONGMANS, GREEN & Co., 39 PATERNOSTER ROW, E.C.;
B. QUARITCH, 11 GRAFTON STREET, NEW BOND STREET, W.;
DULAU & Co., Ltd., 37 SOHO SQUARE, W.;
AND AT THE
BRITISH MUSEUM (NATURAL HISTORY), CROMWELL ROAD, S.W.

1913

[All rights reserved.]

ALERE FLAMMAM.



PRINTED BY TAYLOR AND FRANCIS,
RED LION COURT, FLEET STREET.

PREFACE.

THE present work is a continuation of the series of Catalogues of the Mesozoic Plants in the British Museum. The fossils discussed are even more fragmentary than most of those by which the Jurassic and Wealden Floras are represented; but the Cretaceous Flora is of so much interest as affording possible links with the succeeding Tertiary Flora, that even imperfect evidence is worthy of the closest attention. The literature of the subject is already voluminous, but so many names have been given to these fossils without adequate definition, that it seems unnecessary at present to do more than enumerate them. The greater portion of Part I. of the Catalogue is therefore devoted to an exhaustive bibliography, containing many references to published work which will not occur again in the Descriptive Catalogue.

A. SMITH WOODWARD.

DEPARTMENT OF GEOLOGY,
November, 1913.

AUTHOR'S PREFACE.

WHEN, nearly five years ago, I began a general consideration of the Cretaceous flora, at the request of the British Museum, I discovered that the task could make no progress until a list of the material had been compiled. The literature dealing with Cretaceous plants was widely scattered and had never been brought together and sifted, and there was no standard work to which reference could be made to place the innumerable names which have been published. Consequently, I first endeavoured to compile a list, as nearly complete as possible, of all the Cretaceous species originally described. This list proved immensely greater than was anticipated, and bulks largely in the present volume. As my own work would have been impossible without it, I trust it may prove useful to others; and therefore propose to mention a few points in explanation of its apparent lack of uniformity and its arbitrariness.

In the first place, while including the American Lower Cretaceous species, the list does not take account of most of the European Wealden species. This is not my own choice, but depends on the fact that in this series of Museum Catalogues, the Wealden has been dealt with by Prof. Seward, and the work allotted to me is to complete the Cretaceous, but to leave out of consideration the parts covered by Prof. Seward's work.

Secondly, anyone using the list will notice that the same strata are mentioned under different names, and that the most recent stratigraphical nomenclature is seldom used. This is

because the list does not profess to be critical, but takes the name and the horizon given by the original describer of each species. Therefore, the names of a number of the horizons will be recognisably out of date; but I thought it useful to have the original descriptions catalogued for reference. Further, a number of plants from the Laramie have been included in the list, because in many cases they were described as Cretaceous, though they may be well known to-day to belong to the Tertiary. I included most of those species which had been much quoted as Cretaceous, because it appeared to me that for a reader not intimately acquainted with the Cretaceous, it would be useful to have a list in which references to the original descriptions could readily be found.

Lastly, a point which may make the list appear very incomplete, must be mentioned. Many of the species originally described under one genus have been removed from genus to genus at the hands of various writers. If all the species had been entered under all the generic names ever given to them, the list would have been even bulkier than it is at present. Consequently, while I endeavoured to enter in my list any re-naming that was enlightening, or accompanied by any original work on the specimen, I deliberately left out many of the cases (which are so numerous relating to badly preserved leaf-impressions) in which a species had been transferred to various genera on the grounds of nomenclatorial rules or the personal preferences of an author who neither re-figured nor re-described the specimen, nor in any way added to our knowledge of it. Nevertheless, a good many cross-references to the different genera under which a species appears, will be found. As an example of the type of name which I have tended (perhaps arbitrarily) to omit, *Dryopteris virginica* may be mentioned. Under *Aspidium* Fontaine described a considerable number of species, *A. virginicum* among them. In Knowlton's list of Cretaceous plants, 1898, he transfers most of these to the genus *Dryopteris*, his entry regarding them being in this form: "*Dryopteris virginica* (Font.) n. comb.," followed by reference to Fontaine's

original description. Now, in my list, this species appears under *Aspidium*, with a remark that it was re-named by Knowlton *Dryopteris*; but it is not entered again by me under *Dryopteris*. On the other hand, various species similarly treated by Knowlton and others do find their place in my list under two or even more genera: in most cases this depends on the fact that where I have found a name used in the general literature, or where for some reason it appeared useful to have the different references to the plant, I have added it to my list.

Even after the list was restricted in this fashion, it proved so bulky that only the THALLOPHYTES could be dealt with critically in the present volume. Other portions of the Cretaceous Flora will be considered as is convenient in future volumes.

It is certain that there are many errors in this work, though I have laboured to avoid them. Those who have done any task of the kind will realise how difficult it is to deal with such enormous numbers of references, particularly when parts of the text have to be printed off before the later sections are revised.

In the course of the preparatory work I have travelled much, and though much of the help here mentioned will not bear fruit till the future volumes are completed, I should like to say here how greatly I am indebted to friends and officials for help in the work and for facilities for examining specimens in numerous museums on the Continent and in America. Above all I must thank Mr. W. E. Balston and the Government Grant Committee of the Royal Society of London, for grants partially to defray the cost of these journeys.

Among so many who have rendered assistance in various forms, it seems invidious to single out persons for special thanks, but mention must be made of the kindness of the authorities and staff of the National Museum in Washington, where I had the advantage not only of examining many type-specimens, but also of comparing their bibliographic slips with mine, whereby I gained many additions to my lists. Prof. Nathorst has been most helpful both when on his visits to London and in Stockholm, where he placed the remarkable

State Collections at my disposal. In America, Drs. Knowlton, Hollick, and Berry, and Mr. White were most kind and helpful; as were Dr. Britton of New York, whose great Institute gave me hospitality and the use of the plant collections for some time, and Prof. Jeffrey of Cambridge, who showed me several of his valuable specimens. In Canada, the Director of the Geological Survey was most kind in Ottawa, and also loaned some specimens for further examination in London. In Europe the following must be specially mentioned:—Prof. Rothpletz, Dr. Gothan, Dr. Halle, Dr. Ravn, Prof. Seward, Prof. Oliver, and Dr. Arber, for all have from time to time exerted themselves on my behalf, loaned specimens, or given me help. In the British Museum I also have to thank Dr. Smith Woodward, Dr. Rendle, and Dr. Bather, besides Mr. Gepp and Mr. Newton: while Mr. Sherborn's unique knowledge of old and troublesome books and incomplete bibliographical references has several times proved invaluable.

MARIE C. STOPES.

August, 1913.

CONTENTS.

	Page
INTRODUCTION	xiii
LITERATURE OF CRETACEOUS PLANTS	1
LIST OF SPECIES OF CRETACEOUS PLANTS	49
DESCRIPTIVE CATALOGUE OF CRETACEOUS PLANTS	233
Group ThallopHYta	233
Class Algæ	233
Order Diatomaceæ	235
Order Siphonaceæ	236
Suborder Codiaceæ	236
Genus Bouëina	236
<i>Bouëina Hochstetteri</i> , Toulou	237
Suborder Dasycladaceæ	239
Genus Neomeris	239
<i>Neomeris cretacea</i> , Steinmann	240
Genus Munieria	240
<i>Munieria baconica</i> , v. Hantken	240
Genus Diplopore	243
<i>Diplopore Mühlbergii</i> , T. Lorenz	243
Genus Triploporella	244
<i>Triploporella Fraasi</i> , Steinmann	245
Tribe Acetabulariæ?	247
Order Phæophyceæ?	247
Genus Chondrites	247
<i>Chondrites Targionii</i> (Brongn.), Sternberg	249

	Page
<i>Chondrites intricatus</i> (Brongn.), Sternberg	252
" <i>patulus</i> , Fischer-Ooster	253
Genus <i>Algites</i>	254
<i>Algites furcatus</i> (Brongn.)	255
Order Rhodophyceæ—Corallinaceæ	256
Genus <i>Lithothamnium</i>	256
<i>Lithothamnium mamillosum</i> , GümbeL	257
" <i>perulatum</i> , GümbeL	258
" <i>procoenum</i> , GümbeL	259
" <i>palmatum</i> (Goldf.), GümbeL	259
" <i>Goldfussi</i> , GümbeL	260
" <i>cenomanicum</i> , Rothpletz	260
" <i>turonicum</i> , Rothpletz	262
" <i>amphiroceformis</i> , Rothpletz	262
" <i>gosaviense</i> , Rothpletz	264
Algæ?	265
Class Fungi	267
Subclass Ascomycetes	268
Order Pyrenomycetes	268
Genus <i>Pleosporites</i>	268
<i>Pleosporites Shirainus</i> , Suzuki	268
Genus <i>Petrosphaeria</i>	270
<i>Petrosphaeria japonica</i> , Stopes & Fujii	270
Genus <i>Sphærites</i>	272
<i>Sphærites cretaceus</i> (Heer), Meschinelli	272
" <i>problematicus</i> (Knowlton)	273
Order Discomycetes	273
Genus <i>Phacidites</i>	273
<i>Phacidites circumscriptus</i> (Bayer)	274
Genus <i>Rhytismites</i>	275
<i>Rhytismites hederæ</i> (Heer), Meschinelli	275

CONTENTS.

xi

	Page
Genus <i>Hysterites</i>	276
<i>Hysterites protogæus</i> (Heer), Meschinelli	276
Order Hyphomycetes	276
Genus <i>Cercosporites</i>	276
<i>Cercosporites coriococcus</i> (Bayer)	276
Order Basidiomycetes	273
Genus <i>Trametites</i>	278
<i>Trametites Pini</i> (Conwentz), Meschinelli	278
Genus <i>Trichosporites</i>	279
<i>Trichosporites Conwentzi</i> , Felix.	279
Doubtful Fungi	280

LIST OF FIGURES IN THE TEXT.

	Page
Fig. 1. <i>Boučina Hochstetteri</i> , Toulà; transverse sections . . .	238
2. <i>Halimeda</i> ; cross-section	239
3. <i>Munieria baconica</i> , v. Hantken	240
4. <i>Munieria baconica</i> , v. Hantken; sections & restoration . . .	241
5. <i>Triploporella Fraasi</i> , Steinmann	244
6. <i>Triploporella Fraasi</i> , Steinmann; portion of radial section	245
7. <i>Triploporella Fraasi</i> , Steinmann; reconstruction . . .	246
8. <i>Chondrites Targionii</i> (Brongn.), Sternberg	251
9. <i>Chondrites intricatus</i> (Brongn.), Sternberg	253
10. <i>Chondrites patulus</i> , Fischer-Ooster	254
11. <i>Lithothamnium mamillosum</i> , Gümbel	257
12. <i>Lithothamnium mamillosum</i> , Gümbel; microscopic section	258
13. <i>Lithothamnium perulatum</i> , Gümbel	258
14. <i>Lithothamnium procoenum</i> , Gümbel	259
15. <i>Lithothamnium palmatum</i> (Goldf.), Gümbel	260
16. <i>Lithothamnium Goldfussi</i> , Gümbel	261
17. <i>Lithothamnium cenomanicum</i> , Rothpletz	261
18. <i>Lithothamnium turonicum</i> , Rothpletz	263
19. <i>Lithothamnium amphiroæformis</i> , Rothpletz	263
20. <i>Lithothamnium gosaviense</i> , Rothpletz; section	264
21. <i>Pleosporites Shiraiinus</i> , Suzuki; section of perithecium . . .	268
22. <i>Petrospheria japonica</i> , Stopes & Fujii	271
23. <i>Phacidites circumscriptus</i> (Bayer)	274
24. <i>Cercosporites coriocoecus</i> (Bayer)	277
25. <i>Trichosporites Conwentzi</i> , Felix	280

INTRODUCTION.

THE Cretaceous Flora differs fundamentally from all the older floras in the presence (and in many places the preponderance) of Dicotyledonous and Monocotyledonous plants with the older and "simpler" families. The Angiosperms must undoubtedly trace their ancestors further back than the lowest Cretaceous, but for practical purposes of geology they may almost be described as appearing in the Cretaceous period. Numerous species of Angiosperms have been described and figured from all parts of the world, and these with the other families, which are represented to a greater or less extent, make the Cretaceous flora exceedingly rich as well as interesting. The deposits in some parts of Europe, in Japan, and the Arctic regions, but more particularly in the North American continent, cover large geographical areas and have yielded a great number both of specimens and of species.

To encompass in any one work a description which is more than a mere compilation, of the whole of this vast flora, is impossible in the present state of the science. For in the work on the Cretaceous epoch more perhaps than in that on any other geological period, the data on which determinations have been made are often very unsatisfactory, and the species described have in many cases been named with no regard to the necessity of previous careful comparisons; while generalisations regarding distribution and climate have often been risked on little foundation. There exists no general account of the flora, the bibliography is widely scattered and has never been brought together, and the numerous specific names scattered through the literature have never been listed. The present volume by bringing these things together may prove a useful foundation for future work. I do not think the time has arrived for generalisations or broad conclusions of the kind sometimes attempted by palaeobotanists.

One of the difficulties of work on the Cretaceous flora is the correlation of the many scattered deposits in which the plants have been found. So many local names have been given to beds in different places, that it is difficult to realise sometimes to which period the described plants belong. Added to this is the fact that, particularly in America, the relative positions of the beds are still largely undecided, and individual writers use terms in very different senses. I have not attempted exact correlations, but in the accompanying table* (opposite) have placed in series roughly in their relative positions, the principal beds from which important Cretaceous plants are described. It does not represent the relative thicknesses of the deposits, or, indeed, the final and exact relation of the series, but has merely a temporary value for the convenience of workers not specially acquainted with the beds.

It may also prove of use to give a short summary of the more important work that has been done on the deposits in various parts of the world, thus indicating the distribution of Cretaceous plants in the various countries. Full references to the papers mentioned will be found in the bibliography which follows.

AUSTRIA.

Unger (in 1867) described both Senonian and Cenomanian plants from three principal localities, viz. Ischl, Sanct Wolfgang, and Neue Welt. In 1871 A. Schenk published a monograph on the Urgonian deposits of Wernsdorf and Teschen, with notes on some specimens from neighbouring localities. Schenk also (1876) described eight species of Senonian plants from the northern Tyrol, at Brandenburg.

Of BOHEMIA, much is known owing to the work of many notable palaeobotanists. Corda (1846) in Reuss, described a number of Cenomanian plants from Trzibitz, Perutz and other localities. The Perucer Beds have been the subject of numerous

* NOTE.—This table is for the use of those to whom the local names of horizons are unfamiliar. It does not possess any finality, as the correlations vary with individual judgment. It is based partly on published correlation-tables, partly on the advice of several European and American geologists, and may be regarded as representing approximately the relations of the principal horizons in the more important localities where Cretaceous plants are found.

RAP

[To face page xiv.]

ENT.

America Atlantic Coast.	Mount Livingston.	Pacific Coast.	Black Hills.	Canada.
	Lance.			
nasquan.	Lemnep.			
nocas.	Bear Paw. Judith River.	Wanting.		Vancouver Island.
nmouth.	Claggett. Eagle. Niobrara.	Chico.		
tawan.	Colorado.			Peace River.
?				
gotly				Mill Creek.
fwood.				
itan?				
conformity			Fuson.	
apsco.		Unconformity.		
ndel.			Lakota.	Queen Charlotte Island.
	Kootanie.	Horsetown } Shasta. Knoxville }		Kootanie.
uxent.				

NORTH AMERICAN CONTINENT.																
General European.	England.	Germany.	France.	Bohemia.	Greenland.	America. Atlantic Coast.	Maryland.	America. Eastern Gulf Region.	Western Gulf Region.	Western Interior.	Colorado.	Wyoming.	Mount Livingston.	Pacific Coast.	Black Hills.	Canada.
UPPER CRETACEOUS.	Danian.		Danien			Manasquan.		Wanting.	Wanting.	Denver & Livingston.		Lance.	Lance.			
	Senonian.	Upper Chalk (with flints).	Senon. "Quader" of Saxony.	Senonian.	Senon. { Chlomeker. Priesener. Teplitzer. Iser.	Rancous.	Rancous.	Ripley.	Montana. Navarro.	Laramie.	Laramie (conform. with Cret.).	Leemp.	Bear Paw, Judith River.	Wanting.		Vancouver Island.
	Turonian.	Middle Chalk (without flints).	Turon. Planer Quader.	Turonien.	Turon. { Malnitzer. Weissenberger.	Patoot.	Monmouth.	Monmouth.	Colorado = (a) Austin, (b) Eagleford.	Montana = (a) Foxhills, (b) Fort Pierre & Belly River.	Foxhills, Pierre.	Pierre.	Chico.			
	Cenomanian.	Lower Chalk.	Cenoman. "Pläner" in Westphalia, "Quader" in Saxony.	Cenomanien.	Cenoman. { Korytzauer. Perucer.	Atane.	Magothy.	Magothy.	Wanting.	Dakota.	Dakota.	Dakota.	Dakota.			Peace River.
	Albian.	Gault. Upper Greensand.	Gault.	Albien (Gault).			Raritan?	Raritan?	Unconformity?	Woodbine.	Woodbine.					Mill Creek.
LOWER CRETACEOUS.	Unconformity in places.						Patapsco.	Patapsco.	Unconformity?	Tuscaloosa.						
	Aptian.	Lower Greensand.		Aptien.			Arundel.	Arundel.	Washita.	Unconformity.		Bear River.	Fuson.	Unconformity.		
	Urgonian.								Friedricksburg.							
	Neocomian.								Trinity.							
	Wealden.	Wealden.														
			Hils.			Kome.	Arundel.	Arundel.			Comanche.	Lakota.	Kootanie.	Horsetown	Lakota.	Queen Charlotte Island.
			Neocom.	Neocomien.			Patuxent.	Patuxent.		Kootanie & Morrison.				Shasta. Knoxville		Kootanie.
			Wealden.													

² Raritan used to be placed among the lower series, but Berry (1910) regards it as Upper Cretaceous.

later papers. Krejci (1853) wrote in Bohemian, as did Renger in 1866. Renger (1866 A) had a paper on the Cretaceous tree-ferns of Bohemia. Important work was done by Feismantel (1870, 1872 & 1874) and by Stur (1873), but the deposits are principally known through the publications of Velenovsky. On the Dicotyledons we have Velenovsky's papers published in 1882 A and B, and 1883. A special monograph on the Gymnosperms appeared in 1885. Further papers on the rest of the flora followed in 1886 and 1887; in 1887 A a paper appeared on the Cenomanian plants, and papers in 1888 A, 1888 B (on ferns) and 1889 complete a series of works covering the floras of numerous localities and dealing with most of the families of plants. The most noted localities are perhaps Lipence, Raudnitz, Chlomek, Vyserovic, and Kaunic. Bayer carried on the work and published in 1893, 1896 and 1899; while Fritsch in 1877, 1883, 1889, 1893 and 1897, issued a series of papers dealing with the geology of the deposits, in the later parts of which Bayer deals with the plants. In 1901 Fritsch and Bayer published an important monograph on the Cretaceous plants of the Perucer Beds of the country, which summarises and brings together the previous work on the deposits of this age in Bohemia. In 1901 Marik also published a paper, with plates, in Bohemian.

Unger, in 1865, described a few plants from HUNGARY, and Staub (1888) recorded some Cenomanian plants from Nadrog. Tuzson (1908) has added an illustrated paper on the fossil flora of this region.

For MORAVIA, Heer (1869 A) wrote a large memoir on the Cenomanian deposits at Moletein, near the Bohemian frontier. Krasser, in 1889 and 1896, and Krasser with Kubart (1906) added to the Cretaceous flora of Moletein. The specimens are leaf-impressions, many of them dicotyledonous.

A number of Cretaceous plants from the Island of LESINA were described and illustrated by Kerner in 1896.

BELGIUM.

Coemans in 1867 and Briart and Cornet (1867) described the Belgian Cretaceous plants probably of the age of the Gault, Coemans dealing with a comparatively small number of plant-impressions.

ENGLAND.

Except for the important specimens of *Bennettites* from the Lower Greensand, few Cretaceous plants have been described from England. Some fragments are mentioned by Mantell (1822 & 1844), and supposed fruits from the Chalk were described in 1846, but their nature remains doubtful. Lindley and Hutton (1833-1837) figured several species, principally Gymnospermic cones, of Gault and Greensand age, and some of these have their internal anatomy petrified, though it has not yet been described. Maidstone is an important locality, the Greensand there having yielded a number of plants (see Bensted, 1862, and Mackie, 1862 b). Carruthers (1865) made known *Caulopteris* from the Upper Greensand, and also coniferous fruits from various Cretaceous deposits, in 1866 A, 1866 B, 1868 & 1869. The genus *Cycadeoidea* from the Lower Greensand was described by him in 1867, and he published a paper on Cycadean stems in 1870. Further coniferous fruits were described in 1871 and 1875. In Dixon's 'Geology of Sussex' Carruthers gave a list of the Cretaceous plants (1878), and with Gardner and others (1886) drew up the British Association Committee's report on British Secondary plants.

Williamson (1887) described a further specimen of Gymnospermic cone originally figured by Lindley and Hutton; and in 1898 Barber gave a detailed account of a *Cupressinoxylon* with the internal anatomy preserved from the Lower Greensand. The flora is actually much richer than these published papers would lead one to expect, as will be shown in vol. 2 of this Catalogue.

FRANCE.

The Cretaceous Flora in France is also rather scattered and fragmentary. Brongniart in 1849 described a few plants, and Cornuel (1866) wrote on coniferous fruits of Neocomian age, and published further studies on the subject in 1882. Cricé, in 1877, 1884, 1890 & 1892, published papers dealing with the Cretaceous Flora, the most important being that of 1892.

Saporta in 1880 published on the Lower Cretaceous of Havre, and Bertrand in 1883 described a new genus of Taxineau fossil

from the Upper Cretaceous. In 1890 Marion had a short paper on the Turonian; Saporta described species of *Nelumbium* from the Uppermost Cretaceous (Danian); and Vasseur recorded a Turonian Flora near Martigues. Fliche (1892, 1893) described two Albian plants, and in 1894 had a paper on the fructifications of Cenomanian palms. In 1896 he published a large illustrated paper on the Albian and Cenomanian deposits, and in 1900 one on the Lower Cretaceous.

Lignier (1907) described several specimens of wood from Normandy, with the internal structure petrified.

GERMANY.

PRUSSIA.—The deposits of Aix-la-Chapelle (or Aachen) are classical ground for the palæobotanist, and were first worked by Goeppert (1842 B). The locality, however, is generally associated with the name of Debey, and then Debey & Ettingshausen. Debey (1848 A, 1848 B, 1849, 1850) published several short papers on the Aix Cretaceous plants, and then with Ettingshausen (1859 A & B) prepared two large, well-illustrated monographs on the "Thallophyten" and the "Acrobryen" of the Cretaceous of Aix and Maestricht. Debey made large collections and was apparently preparing voluminous further publications, but in 1877 only a very short note appeared about a paper he gave on the Coniferæ of the Cretaceous of Aix. His collections with the specimens bearing unpublished names, are evidence of the richness of the deposits. In 1890 Lange also contributed to our knowledge of these beds.

The Senonian sandstones of Quedlinburg and neighbourhood have also yielded an important Flora, first noticed by Geinitz (1850 D), but more exhaustively described by Stiehler (1854, 1857, and especially 1858). Heer (1871 A) made a small contribution, and Richter has continued the work during more recent years. In 1899 B he published a paper on the Conifers of Quedlinburg, and in 1901 a monograph on the Neocomian plants of the same district. In 1905 he described numerous Senonian plants, and published a valuable revision of the genus *Credneria*. In 1906 he prepared a similar revision of *Hausmannia*, with notes on the Lower Cretaceous Flora; and in 1909 there followed an equally exhaustive monograph on *Nathorstiana*.

WESTPHALIA.—This district, rich in fossil plants, has Cretaceous Floras of several successive horizons.

Roemer (1841) referred to the Cretaceous plants, but the earliest work of any note is W. von der Marek's monograph in the 'Palæontographica,' vol. xi. (1864), in which he described a number of species from the Plattenkalk. Saporta (1867) made Hal'dem a classical locality; and Hosius (1870) dealt with the more geological aspect of the beds of Cretaceous age in several localities. Hosius (1870 A) also described and figured a number of Dicotyledonous leaf-impressions from these deposits. Hosius and von der Marek (1880 & 1885) finally gave full and excellent accounts of the Upper and Lower Cretaceous plants of Westphalia, with good descriptions and figures, noticing a number of localities.

SILESIA.—The Quadersandstein (Upper Cretaceous) of Silesia was described by Goeppert in 1842 A and 1847. He illustrated and named plants from Kieslingswalde, Bunzlau, and elsewhere. Roemer (1889) also published an account of Senonian leaf-impressions from Bunzlau.

SAXONY.—The well-known fossil plants from the Cenomanian of Niederschoena have been described by several authors. They are often mentioned by Sternberg, 'Flora der Vorwelt' (1820-1838), some of them were noticed by Cotta (1836), and numerous plant-impressions were well described and illustrated by Geinitz (1842). The first really important work on this Flora, however, was that by Ettingshausen (1867 A), who dealt with dicotyledonous leaf-impressions; and his researches were followed by those of Engelhardt (1892 A).

The neighbourhoods of Dippoldiswalde and Dresden are also known for Cretaceous plant-bearing deposits, and here Glocker (1841) discovered his so-called *Gyrophyllites*. Otto in his *Addimenta* (1852 & 1854) enumerated and gave an account of the Quader plants of this region. Geinitz (1875 A) also published a paper on the plants of the Lower Quader, with a second part on those of the Middle and Upper Quader of Saxony.

HOLLAND.

The Cretaceous plants of Holland are all impressions, and are similar to those of Aix. Debey (1851) studied the specimens from Maestricht; and Miquel (1853) published a monograph on

the deposits, with a number of plates and text-figures of the plants. In 1861 Bosquet enumerated a hundred species from the Limburg Chalk of Maestricht and elsewhere.

ITALY.

Fossil Algæ and Cycads are the principal plants described from the Cretaceous of Italy. In 1891 Bozzi wrote a paper on the Cretaceous Flora which was illustrated with two plates. The most important work is that of Capellini and Solms-Laubach (1892) dealing with the Bennettitæ, some of which are of Cretaceous age.

PORTUGAL.

The Cretaceous plants of Portugal are of special interest because among them are Dicotyledons, supposed to be the earliest to reach Europe. Neocomian or Wealden plants from Almargem and elsewhere were described by Heer (1881). Cenomanian Dicotyledons were described in 1888 by Saporta, who subsequently (1891) gave a brief account of the "most ancient European Dicotyledons" of Cretaceous age, and in 1894 published his classic Memoir. De Lima (1901) also published a short paper on Senonian plants from this country.

RUSSIA.

In Eichwald's work on the fossil Flora of Russia (1853, 1860, 1861, 1862, 1865) he dealt with a number of Cretaceous plants, mostly of Neocomian age. Mercklin (1855) described petrified woods with their structure preserved, and Krendowsky (1880) described in Russian and figured several new species of Cretaceous plants.

SWEDEN.

Surprisingly few fossil plants of Cretaceous age are known from this country, though the other Mesozoic deposits are very rich in such remains. Nilsson (1824) described and figured some plants from the Greensand of Scania, but did not name them fully and they have been re-named from time to time. Nathorst mentions and re-figures some of these in his works on

Swedish geology. Structurally-preserved woods, including fungal hyphae, were described and figured by Conwentz in 1892 and further investigated by Felix (1894).

SWITZERLAND.

In Heer's 'Die Urwelt der Schweiz,' which first appeared in 1865, he dealt with the fossil Floras of all geological periods in Switzerland, and included an account of the Lower Cretaceous Flora of several localities. In his 'Flora fossilis Helvetiae' (1877) he described the Neocomian plants of Berne and elsewhere, and the Upper Cretaceous plants of Freiburg and Berne. He also referred to the Flysch deposits from which Fischer-Ooster (1858) had already described the peculiar "fucoïds." Though chiefly Tertiary, the lower horizons of these beds are probably of Cretaceous age.

MADAGASCAR.

Petrified wood from Madagascar has proved interesting and important (see Fliche, 1900 A, 1905).

EGYPT.

The petrified forests of Egypt were the subject of much comment even by the early writers. Among later authors, Unger (1859) and Carruthers (1870 A) may be mentioned, but the most important work was done in connection with Rohlfs' expedition (in Zittel, Palaeontographica, 1883), the woods being determined by Schenk. The plants are supposed to be Upper Cretaceous, and geological evidence tends to prove this for some of the horizons; but there is still doubt as to whether much of that described as Cretaceous wood is not really Tertiary. The same plant-bearing deposits extend into Nubia, where much of the wood is probably of Cretaceous age.

AUSTRALASIA.

The plant-bearing beds of this continent are not so well known as could be wished. Unger (1866) first mentioned Cretaceous plants from New Zealand, and Ettingshausen (1887 A) subsequently described numerous species from Upper Cretaceous

formations. A translation of the latter work appeared in New Zealand in 1891. In 1893 the same author published a preliminary essay on the Cretaceous Flora of Australia, followed by an illustrated paper in 1895; and a general account of the Cretaceous Flora of the Southern hemisphere appeared in 1896. Finally in 1896 Johnston described and figured several plants, probably of Upper Cretaceous age, from Tasmania.

JAPAN.

Lower Cretaceous plant-impressions from Japan were first described by Yokoyama (1889) in his paper on the Jurassic Flora, and additional species were subsequently described and figured by Nathorst (1890) and Yokoyama (1894). He pointed out the very close affinity of many of them to the Potomac plants of North America. The Japanese Cretaceous formations, however, are notable as being the only deposits of Mesozoic age which have hitherto yielded petrified masses of plants in many respects similar to the well-known Carboniferous "coal-balls." In a paper largely dealing with Tertiary specimens Reiss (1907) described a few fragments of Gymnospermic wood from these nodules; and the remains of an extensive flora, including the leading plant phyla, have been discovered in them by Stopes and Fujii (1909, 1910), Stopes (1909, 1910), Kershaw (1910), and Suzuki (1910).

INDIA.

Feistmantel (1877) described a tree-fern from the Cretaceous of India, but all other records of Cretaceous fossil plants from this region are doubtful.

CHINA.

Yokoyama (1906) records from China a few impressions of plants which are probably of Lower Cretaceous age.

NORTH AMERICA.

CANADA.—Much work on the extensive deposits of Cretaceous plants was done by Dawson (1873-1894), who dealt principally with the numerous impressions of dicotyledonous leaves and

fragments of gymnosperms and ferns, though he found and described several woods with petrified structure. Penhallow continued the investigation of the impressions, and also treated of Gymnospermic woods, but his most interesting addition to the Flora was that of *Osmundites* with its internal anatomy (1902 A).

UNITED STATES.—The enormous extent of the Cretaceous deposits in this country, and the richness of the beds in plant-impressions, have led many workers to investigate them. The list of American publications is indeed lengthy, and instead of summarising them here, it may be better to refer to Ward's exhaustive paper (Ward, 1889, pp. 834-926) for an account of the earlier literature. Knowlton's Catalogue (1898) of the American Tertiary and Cretaceous plants brings the lists more up to date. Since this time the most active workers have been Berry (1901-1910), Knowlton (1889-1910), Hollick (1892-1910), Hollick & Jeffrey (1909), and finally Wieland, whose publications are almost confined to the Cycadophyta, which are principally from older deposits but afford results of great importance in a consideration of the Cretaceous Flora. The majority of the American specimens are in the form of leaf-impressions, and they occur eastwards in the clays and shales, and westwards in the great Coal-bearing series. Several species of petrified (silicified) wood have been described by Knowlton; but the most important plants with their internal anatomy are those with which Hollick & Jeffrey and their pupils have made the locality of Staten Island famous. These plants, preserved in the fine clay (Amboy Clay), are not exactly petrifications in the ordinary sense, but are fragments in which the internal parts are fairly well preserved, much as they would be in peat. Dr. Berry's important monograph on the Lower Cretaceous plants of America has now appeared. Unfortunately the present work only includes literature and species appearing before the end of 1910, and as so much of the present volume was in type before Dr. Berry's work was published, it is impossible to include his results.

MEXICO.—Petrifications of wood have been described by Felix & Nathorst from the State of Oaxaca; and Steinmann (1899) recorded and described interesting Siphonaceæ with their structure well preserved.

SOUTH AMERICA.

Kurtz (1902) directed attention to the existence of a Flora of the age of the "Dakota Group" in Argentina. The Neocomian flora has also received attention from Neumann (1907).

ARCTIC REGIONS.

The Cretaceous Floras of the Arctic are so well known through Heer's classical work (see 'Flora Fossilis Arctica,' various parts in bibliography) that it will be necessary merely to mention his papers. The Kome, Atane, and Patoot Beds of Greenland are the best known, and are most important from a geological point of view on account of the proof they afford of the great changes of climate which must have occurred in that region. They have yielded a large number of plant-impressions, principally Gymnosperms and Angiosperms, with a few characteristic Ferns. Unfortunately, however, petrified specimens are not available, although something can be done with many of the cuticles, and the material on the whole is well preserved. Nathorst has continued and amplified Heer's work, his large collections of impressions testifying to the richness of the deposits; and his description of a Cenomanian plant from Greenland (1907) forms a valuable contribution to the literature of the Cycadophyta.



LITERATURE

OF

CRETACEOUS PLANTS

UP TO DECEMBER, 1910.

-
- AGARDH, C. A.—1824. *Systema Algarum*. Pp. xxxviii & 312. Lund.
- ARBENZ, P.—1908. Ueber Diploporon aus dem Schrattenkalk des Santisgebietes: Vierteljahrsschrift Naturf. Ges. Zürich, vol. 53, pp. 387-392.
- D'ARCHIAC, VICOMTE.—1837. Mémoire sur la Formation Crétacée du Sud-ouest de la France: Mém. Soc. géol. France, vol. 11, pp. 157-192.
- BAILEY, I. W.—1910. Anatomical characters in the evolution of *Pinus*: Amer. Nat., vol. 44, pp. 284-293, pl. i.
- BARBER, C. A.—1898. *Cupressinoxylon vectense*; a fossil Conifer from the Lower Greensand of Shanklin, in the Isle of Wight: Annals Bot., vol. 12, pp. 329-361, pls. xxiii-xxiv.
- BARROIS, C.—1876. Recherches sur le Terrain Crétacé supérieur de l'Angleterre et de l'Irlande. Mém. Soc. géol. Nord, vol. 1, pp. 1-232 & maps, pls. i-iii.
- BARTSCH, P.—1896. Notes on the Cretaceous Flora of Western Iowa: Bull. Iowa State Univ. Lab. Nat. Hist., vol. 3, no. 4, pp. 178-182.
- BAYER, E.—1893. Die Flora der Priesener Schichten, résumé of O Rostlinstvu vrstev Březenských: Vestník Kral. Cezké Společnosti Nauk, pp. 1-50, résumé pp. 34-50, text illust.
- 1893 A. See FRITSCH.—1893.
- 1896. Die Flora der Chlomeker Schichten, résumé of O Rostlinstvu vrstev Chlomeckých: Vestník Kral. Cezké Společnosti Nauk, pp. 1-36, résumé pp. 29-36, text illust.
- 1899. Einige neue Pflanzen der Perucr Kreideschichten in Böhmen: Sitzb. k. böhm. Ges. Wiss., pp. 1-51, pls. i, ii.
- 1897. See FRITSCH.—1897.
- 1901. See FRITSCH & BAYER.—1901.

- BENNETT, E.—1831. A Catalogue of the Organic Remains of the County of Wilts. 9 pp., xviii pls. Warminster.
- BENSTED, W. H.—1862. The Geology of Maidstone: Geologist, vol. 5, see pp. 336, 337, pl. xix.
- BERRY, E. W.—1901. Notes on *Liriodendron* Leaves: Torrey, vol. 1, pp. 105-107, pls. i, ii.
- 1901 A. The Origin of Stipules in *Liriodendron*: Bull. Torrey Bot. Club, vol. 28, pp. 493-498, pls. xli, xlii.
- 1902 A. *Liriodendron Celakovskii*, Velen.: Bull. Torrey Bot. Club, vol. 29, pp. 478-480.
- 1902 B. Notes on *Sassafras*: Bot. Gazette, vol. 34, pp. 426-450, pl. xviii.
- 1902 C. Additional Notes on *Liriodendron* Leaves: Torrey, vol. 2, no. 3, pp. 33-37, pls. i, ii.
- 1902 D. Notes on the Phylogeny of *Liriodendron*: Bot. Gazette, vol. 34, pp. 44-63, text-fig.
- 1903 A. *Avatia* in American Paleobotany: Bot. Gazette, vol. 36, pp. 421-428.
- 1903 B. *Liriodendron* Notes: Torrey, vol. 3, no. 9, pp. 129-132, figs. 1-3.
- 1903 C. New Species of Plants from the Matawan Formation: Amer. Naturalist, vol. 37, pp. 677-684.
- 1903 D. The American Species referred to *Thinnfeldia*: Bull. Torrey Bot. Club, vol. 30, pp. 438-445.
- 1904 A. Additions to the Flora of the Matawan Formation: Bull. Torrey Bot. Club, vol. 31, pp. 67-82, pls. i-v.
- 1904 B. The Cretaceous Exposure near Cliffwood, N.J.: Amer. Geologist, vol. 34, pp. 253-260, pl. xv.
- 1904 C. Otto Kuntze on *Sequoia*: Torrey, vol. 4, pp. 153, 154.
- 1905 A. Additions to the Fossil Flora from Cliffwood, New Jersey: Bull. Torrey Bot. Club, vol. 32, pp. 43-48, pls. i, ii.
- 1905 B. A *Ficus* confused with *Proteoides*: Bull. Torrey Bot. Club, vol. 32, no. 6, pp. 327-330, pl. xxi.
- 1905 C. A Palm from the Mid-Cretaceous: Torrey, vol. 5, pp. 30-33, figs. 1, 2.
- 1905 D. Fossil Grasses and Sedges: Amer. Naturalist, vol. 29, pp. 345-348.
- 1905 E. The Flora of the Matawan Formation (Crosswick Clays): Bull. New York Bot. Gard., vol. 3 (1903-05), pp. 45-103, pls. xliii-lvii.
- 1906. A Note on the Mid-Cretaceous Geography: Science, vol. 23, pp. 509-510.
- 1906 A. Living and Fossil Species of *Comptonia*: Amer. Naturalist, vol. 40, pp. 485-520, pls. i-iv.
- 1906 B. Contributions to the Mesozoic Flora of the Atlantic Coastal Plain.—I: Bull. Torrey Bot. Club, vol. 33, pp. 163-182, pls. vii-ix.
- 1906 D. A Brief Sketch of Fossil Plants: Ann. Rep. Geol. Surv. New Jersey for 1905, pp. 97-133.

- BERRY, E. W.—1906 E. The Flora of the Cliffwood Clays: Ann. Rep. Geol. Surv. New Jersey for 1905, pp. 135-172, pls. xix-xxvi.
- 1907. Contributions to the Mesozoic Flora of the Atlantic Coastal Plain.—II. North Carolina: Bull. Torrey Bot. Club, vol. 34, pp. 185-206, pls. xi-xvi.
- 1907 A. Palaeobotanical Notes: Cretaceous Floras in North and South Carolina. New Species of Plants from the Magothy Formation: Johns Hopkins Univ. Circ., 1906-7, no. 7, pp. 667-679 = 79-91, text-figs. 1-6.
- 1908 A. Some Araucarian Remains from the Atlantic Coastal Plain: Bull. Torrey Bot. Club, vol. 35, pp. 249-260, pls. xi-xvi.
- 1908 B. A Mid-Cretaceous Species of *Torreya*: Amer. Journ. Sci., ser. 4, vol. 25, pp. 382-386, text-figs.
- 1908 C. A New Cretaceous *Bauhinia*: Torreya, vol. 8, pp. 218-219, text-figs.
- 1909. Contributions to the Mesozoic Flora of the Atlantic Coastal Plain.—III. New Jersey: Bull. Torrey Bot. Club, vol. 36, pp. 245-264, pls. xviii, xviii a.
- 1910 A. Contributions to the Mesozoic Flora of the Atlantic Coastal Plain.—V. North Carolina: Bull. Torrey Bot. Club, vol. 37, pp. 181-200, pls. xix-xxiv.
- 1910 B. Contributions to the Mesozoic Flora of the Atlantic Coastal Plain.—VI. Georgia: Bull. Torrey Bot. Club, vol. 37, no. 10, pp. 503-511, text-figs. 1, 2.
- *— 1910 C. A Revision of the Fossil Plants of the Genus *Nageiopsis* of Fontaine: Proc. U.S. Nat. Mus., vol. 38, pp. 185-195.
- 1910 D. A Revision of the Fossil Plants of the Genera *Acrostichopteris*, *Teniopteris*, *Nilssonia*, and *Sapindopsis* from the Potomac Group: Proc. U.S. Nat. Mus., vol. 38, pp. 625-644.
- 1910 E. A New Species of *Deualquesa* from the American Cretaceous: Torreya, vol. 10, pp. 34-38, text-fig.
- 1910 F. The epidermal characters of *Frenelopsis ramosissima*: Bot. Gazette, vol. 50, no. 4, pp. 305-309, text-figs. 1, 2.
- 1910 G. A Cretaceous *Lycopodium*: Amer. Journ. Sci., ser. 4, vol. 30, pp. 275-6, text-fig. 1.
- 1910 H. Contributions to the Mesozoic Flora of the Atlantic Coastal Plain.—IV. Maryland: Bull. Torrey Bot. Club, vol. 37, pp. 19-29, pl. viii.
- 1910 J. The evidence of the Flora regarding the Age of the Baritan Formation: Journ. Geol., vol. 18, pp. 252-258.
- 1910 K. A New Cretaceous *Bauhinia* from Alabama: Amer. Journ. Sci., ser. 4, vol. 29, pp. 256-258.
- 1910 L. Geologic Relations of the Cretaceous Floras of Virginia and North Carolina: Bull. Geol. Soc. Amer., vol. 20, pp. 655-659.

- BERTRAND, C. E.—1883. Sur le genre *Vesquia*, Taxinée fossile du Terrain Aachénien de Tournai : Comptes Rendus Acad. Sci. Paris, vol. 97, pp. 1382-1384.
- 1883 a. Note sur le Genre *Vesquia*, Taxinée fossile du Terrain Aachénien de Tournai : Bull. Soc. bot. France, vol. 30, pp. 293-299.
- BEUST, F.—1884. Untersuchung ueber fossile Hölzer aus Gröuland : Neue Denkschr. schweiz. natur. Ges., pp. 1-43, pls. i-vi.
- BESSEY, C. E.—1897. The Phylogeny and Taxonomy of Angiosperms : Bot. Gazette, vol. 24, pp. 145-178.
- BIBBINS, A.—1895. Notes on the Palaeobotany of the Potomac Formation : Johns Hopkins Univ. Circ., vol. 15, no. 121, pp. 17-20, 1 pl. unnumb.
- 1898. A fossil Cypress Swamp in Maryland : The Plant World, vol. 1, pp. 164-166, 1 pl.
- 1905. See WARD, L. F.—1905.
- BLANCKENHORN, M.—1890. Beiträge zur Geologie Syriens : Die Entwicklung des Kreidesystems in Mittel- und Nord-Syrien mit besonderer Berücksichtigung der paläontologischen Verhältnisse nebst einem Anhang über den jurassischen Glandarienkalk. 135 pp., xi pls. Cassel [see p. 61].
- BÖHM, J.—1885. Der Grünsand von Aachen und seine Molluskenfauna : Inaug.-Diss. Bonn. Universität, 155 pp., ii pls. Bonn.
- 1903. Ueber cretaceische und eocene Versteinerungen aus Fergana : Durchasien, vol. 3, lief. 1, Palaeontol. pp. 91-112, 1 pl. Berlin.
- BOMMER, C.—1903. Les Causes d'Erreur dans l'étude des Empreintes végétales : Nouv. Mém. Soc. belge Géol., fasc. i, pp. 1-32, pls. i-x.
- BONARELLI, G.—1899. I fossili senoniani dell' Apennino centrale che si conservano a Perugia nella Collezione Bellucci : Atti R. Accad. Sci. Torino, vol. 24, pp. 1020-1027, pl. i.
- BORNEMANN, J. G.—1856. Ueber organische Reste der Lettenkohlengruppe Thüringens. Ein Beitrag zur Fauna und Flora dieser Formation besonders ueber fossile Cycadeen, nebst vergleichenden Untersuchungen ueber die Blattstruktur der jetztleblichen Cycadeengattungen. 85 pp., xii pls. Leipzig.
- BORNHARDT, W.—1900. Zur Oberflächengestaltung und Geologie Deutsch-Ostafrikas : Deutsch-Ost-Afrika, vol. 7, pp. 1-595, & maps. Berlin.
- BOSQUET, J.—1861. Coup d'œil sur la répartition géologique et géographique des espèces d'animaux et de végétaux citées dans le tableau des fossiles Crétacés du Limbourg inséré dans la dernière livraison de l'ouvrage du Dr. W. C. H. Staring sur le sol de la Néerlande : Verslag. K. Akad. Wet. Natuurk., vol. 11, pp. 108-120, table.
- BOULENGER, G. A., and LYDEKKER, R.—1889. A Wooden Dinosaur : Geol. Mag., dec. 3, vol. 6, pp. 191-192.
- BOZZI, L.—1888. Sulle filliti cretacee di Vernasso nel Friuli : Atti Soc. Ital. Sci. Nat., vol. 31, pp. 399-405, pl. vi.

- Bozzi L.—1891. La Flora Cretacea di Vernasso nel Friuli: Bol. Soc. Geol. Ital., vol. 10, pp. 371-382, pls. xv, xvi.
- BRAUN, A.—1854. Einige Beiträge zur Flora der Tertiär-Zeit: Neues Jahrb. f. Min., pp. 138-147, pl. iii.
- BRAUNS, D.—1876. Die senonen Mergel des Salzbergs bei Quedlinburg und ihre organischen Einschlüsse: Zeitschr. gesamt. Naturwiss., vol. 46, pp. 325-420, pls. vii-x.
- BRIART, A., and CORNET, F. L.—1867. Description minéralogique et stratigraphique de l'Étage Inférieur du Terrain Crétacé du Hainault (System Aachénien de Dumont). Suivie de la Description des Végétaux fossiles de cet Étage par M. E. Coemans: Mém. Acad. roy. Belgique, vols. 33 & 36. See COEMANS.—1867.
- BRISTOW, H. W.—1889. Geology of the Isle of Wight, ed. 2, revised by C. Reid & A. Stralman: Mem. Geol. Surv. England & Wales. Pp. 349, pls. i-v.
- BRITTON, N. L.—1889. Remarks on recent discoveries in local Cretaceous and Quaternary Geology: Trans. New York Acad. Sci., vol. 8, pp. 177-181.
- BRONGNIART, A.—1824. Observations sur les Fucoides, et sur quelques autres Plantes marines fossiles: Mém. Soc. d'Hist. nat. Paris, vol. i, pt. 2, pp. 301-321, pls. xix-xxi.
- 1828. Histoire des Végétaux Fossiles, ou Recherches botaniques et géologiques sur les Végétaux renfermés dans les divers couches du globe, vol. 7, 488 pp., clxvi pls. Paris. (See Note on p. 48.)
- 1828 A. Prodrôme d'une Histoire des Végétaux Fossiles. 223 pp. Paris.
- 1849 A. Tableau des genres de Végétaux fossiles considérés sous le point de vue de leur classification botanique et leur distribution géologique: repaged Extract from Dict. univ. d'Hist. Nat. Pp. 127.
- 1849 B. Exposition chronologique des périodes de végétation et des Flores diverses qui se sont succédé à la surface de la terre: Ann. Sci. Nat., Bot., ser. 3, vol. 11, pp. 285-338.
- BRONX, H. G.—1837. Lethæa Geognostica, oder Abbildungen und Beschreibungen der . . . Versteinerungen, 1346 pp., xlvii pls. Stuttgart.
- 1849. Index Palæontologicus. Vol. 2, Enumerator Palæontologicus, 980 pp.
- and ROEMER, F.—1852. Lethæa Geognostica, vol. 2, pp. 1-56. Stuttgart.
- BROWN, R.—(1851) 1855. *Cycadites Saxbyanus*: Proc. Linn. Soc., vol. 2, p. 130.
- 1877. Geological Notes on the Noursoak Peninsula, Disco Island, and the country in the vicinity of Disco Bay, North Greenland: Trans. Geol. Soc. Glasgow, vol. 5, pp. 55-112 & map.
- CALVIN, S.—1894. On the Geological Position of *Bennettites dacotensis*, MacBride, with remarks on the Stratigraphy of the region in which the species was discovered: Amer. Geologist, vol. 13, pp. 79-84.

- CAPELLINI, G.—1890. *Ichthyosaurus campylodon* e Tronchi di Cicadee nelle argille scagliose dell' Emilia : Mem. R. Accad. Sci. Bologna, ser. 4, vol. 10, pp. 431-450, pls. i, ii.
- and HEER, O.—1867. Les Phyllites crétacées du Nebraska.—Sur les gisements des Phyllites du Nebraska.—Note géologique par G. Capellini.—Sur les plantes fossiles du Nebraska par O. Heer : Nouv. Mém. Soc. helv. Sci. Nat., vol. 22, pp. 1-22, pls. i-iv.
- and SOLMS-LAUBACH, E.—1892. I Tronchi di Benettitee dei Musci Italiani Notizie storiche, geologiche, botaniche : Mem. R. Accad. Sci. Bologna, ser. 5, vol. 2, pp. 67-120, pls. i-v.
- CARRUTHERS, W.—1865. On *Caulopteris punctata*, Goepp., a Tree-fern from the Upper Greensand of Shaftesbury in Dorsetshire : Geol. Mag., vol. 2, pp. 484-487, pl. xiii.
- 1866 A. On Araucarian Cones from the Secondary Rocks of Britain : Geol. Mag., vol. 3, pp. 249-252, pl. xi.
- 1866 B. On some Fossil Coniferous Fruits : Geol. Mag., vol. 3, pp. 534-546, pls. xx, xxi.
- 1867. On *Cycadoidea Yatesii*, a Fossil Cycadean Stem from the Potton Sands, Bedfordshire : Geol. Mag., vol. 4, pp. 199-201, pl. ix.
- 1868. British Fossil Pandanæ : Geol. Mag., vol. 5, pp. 153-156, pl. ix.
- 1869. On some Undescribed Coniferous Fruits from the Secondary Rocks of Britain : Geol. Mag., vol. 6, pp. 1-7, pls. i, ii.
- 1870. On Fossil Cycadean Stems from the Secondary Rocks of Britain : Trans. Linn. Soc., vol. 26, pp. 675-708, pls. liv-lxiii.
- 1870 A. On the Petrified Forest near Cairo : Geol. Mag., vol. 7, pp. 306-310, pl. xiv.
- 1871. On two Undescribed Coniferous Fruits from the Secondary Rocks of Britain : Geol. Mag., vol. 8, pp. 540-544, pl. xv.
- 1875. Note on the Flora of the Gault, with Description of a New Pine Cone : Proc. Geol. Assoc., vol. 4, pp. 278-281, text-fig.
- 1878. The Plant Remains of the Upper and Lower Cretaceous (Neocomian) Formations in England, in Dixon's Geol. Sussex, ed. 2, ref. pp. 277-282. See Dixon, F.—1878.
- CARUEL, T.—1870. Osservazioni sul genere di Cicadacee fossili *Rauumeria*, e descrizione di una specie nuova : Boll. R. Comitato Geol. Italia, vol. 1, pp. 181-186, pl. i.
- CASPARY, R.—1888. Einige fossile Hölzer Preussens : Schrift. pays.-ökonom. Ges. Königsberg, Abhandl., vol. 28, pp. 29-45.
- 1889. Einige fossile Hölzer Preussens, bearbeitet v. Triebel : Abhandl. geol. Specialkarte Preussens, vol. 9, pt. 2, pp. viii, 1-86, pls. i-xv.
- CAYEUX, L.—1892. Sur la présence de nombreuses Diatomées dans les gaizes crétacées du Bassin de Paris : Comptes Rendus Acad. Sci. Paris, vol. 114, pp. 375-377.
- 1897. Contribution à l'étude micrographique des Terrains sédimentaires.—II. Craie du Bassin de Paris : Mém. Soc. géol. Nord, vol. 4, pp. 206-563, pls. vii-x.

- CHOFFAT, P.—1885. Sur le Système crétacique du Portugal.—1^{re} étude : Impr. Acad. roy. Sci. Sect. Trav. géol. Portugal, pp. 1-68, pls. i-iii.
- 1889. Note sur le Crétacique des Environs de Torres-Vedras, de Peniche et de Cerenl [Fossil plants identified by Saporta]: Commun. Commissao Trabalhos Geol. Portugal, vol. 2, pp. 171-215.
- CHRYSLER, M. A.—1906. See JEFFREY, E. C., and CHRYSLER.—1906.
- CLARK, W. B.—1894. Origin and Classification of the Greensands of New Jersey: Journ. Geology, vol. 2, pp. 161-177.
- 1895. Cretaceous Deposits of the Northern Half of the Atlantic Coastal Plain: Bull. Geol. Soc. America, vol. 6, pp. 479-482.
- 1904. The Matawan Formation of Maryland, Delaware, and New Jersey, and its relation to overlying and underlying Formations: Amer. Journ. Sci., ser. 4, vol. 18, pp. 435-440.
- 1908. Results of a recent investigation of the Coastal Plain Formations in the area between Massachusetts and North Carolina: Bull. Geol. Soc. America, vol. 20, pp. 646-654, pl. iii.
- COCKERELL, T. D. A.—1909. Amber in the Laramie Cretaceous: Torreya, vol. 9, pp. 140-142, text-fig.
- COEMANS, E.—1867. Description de la Flore fossile du premier Étage du Terrain Crétacé du Hainaut: Mém. Acad. roy. Belgique, vol. 36, pp. 1-20, pls. iii-v. [plates wrongly numbered as being from vol. 33.] [Reprinted with Briart & Cornet, 1867.]
- CONRAD, T. A.—1869. Notes on American Fossiliferous Strata: Amer. Journ. Sci., ser. 2, vol. 47, pp. 358-364.
- CONWENTZ, H.—1892. Untersuchungen ueber fossile Hölzer Schwedens: K. Svensk. Vet.-Akad. Handl., vol. 24, no. 13, pp. 1-99, pls. i-xi.
- CORDA, A. J.—1838. Skizzen zur vergleichenden Phytotomie vor- und jetztweltlicher Pflanzen-Stämme, in STERNBERG.—1838. Pp. i-lxxi.
- 1845. Beiträge zur Flora der Vorwelt, 128 pp., lx pls. Prague.
- 1846. Pflanzen Versteinerungen der böhmischen Kreideformation, in A. E. Reuss, Die Versteinerungen der böhmischen Kreideformation, pt. ii, pp. 81-96, pls. xvi-li. Stuttgart.
- CORNET, F. L.—1867. See BRIART, A., & CORNET, F. L.—1867.
- CORNET, J.—1908. Sur quelques Bois fossiles du Crétacique marin du Hainaut: Ann. Soc. géol. Belgique, vol. 35, pp. 322-324.
- CORNUEL, J.—1866. Description des cônes de pins trouvés dans les couches fluviolacustres de l'étage néocomien du bassin parisien, précédée de diverses appréciations d'après leur état, et d'observations sur l'origine des eaux de la lagune dans laquelle ces cônes ont été fossilisés: Bull. Soc. géol. France, ser. 2, vol. 23, pp. 658-673, pl. xii.
- 1882. Note sur les cônes de *Pinus elongata* découverts à Saint-Dizier (Haute-Marne), et sur des cônes de Cèdre du sable vert de la Houpelte (Meuse): Bull. Soc. géol. France, ser. 3, vol. 10, pp. 259-263, pl. vii.

- COTTA, B.—1836. Ueber die Niederschöna-Schichten: Neues Jahrb. f. Min., pp. 584-587.
- COUFFON, O.—1909. Les Grès à *Sabalites Andegavensis* en Anjou: Bull. Soc. Sci. d'Angers, vol. 38, pp. 9-22, pls. i-vii.
- 1910. Les Grès à *Sabalites* en Anjou (Supplément): *loc. cit.*, vol. 39, pp. 21-31.
- COUYAT, J., & FRITEL, P. H.—1910. Sur la présence d'empreintes végétales dans le grès nubien des environs d'Assouan: Comptes Rendus Acad. Sci. Paris, vol. 151, pp. 961-964.
- CRAGIN, F. W.—1889. Contributions to the Palaeontology of the Plains. —No. 1: Bull. Washburn College Laboratory Nat. Hist., vol. 2, no. 10, pp. 65-68.
- CREDNER, H.—1902. Elemente der Geologie, 802 pp. Leipzig.
- CRÉÉ, L.—1877. Considérations sur la Végétation de l'Ouest et du Nord-Ouest de la France, aux époques géologiques: Bull. Soc. Linn. Normandie, ser. 3, vol. 1, pp. 225-232.
- 1879? Les Anciens Climats et les Flores Fossiles de l'ouest de la France, 74 pp. and frontispiece. Rennes.
- 1884. Contributions à la Flore crétacée de l'ouest de la France: Comptes Rendus Acad. Sci. Paris, vol. 99, pp. 511-513.
- 1889. Exposition Paléophytique.—Paléontologie des Colonies Françaises, Expos. Univ. Paris, pp. 1-32.
- 1890. Recherches sur les Végétaux fossiles de l'Île d'Aix (Charente-Inférieure): Ann. Soc. Sci. nat. Charente-Inférieure, no. 26, pp. 231-237, pls. i, ii.
- 1892. Recherches sur les Palmiers silicifiés des terrains crétacés de l'Anjou: Bull. Soc. d'Etudes Sci. d'Angers, vol. 21, pp. 97-103, 2 plates, figs. 1-4.
- CROSS, W.—1896. See EMMONS, CROSS, and ELDRIDGE.—1896.
- CUMMINS, W. F.—1892. Report on the Geography, Topography, and Geology of the Llano Estacado or Staked Plains, with notes on the Geology of the Country west of the Plains, in 3rd Ann. Rep. Geol. Surv. Texas, pp. 129-223, text-figs.
- DARTON, N. H.—1893. The Magothy Formation of Northeastern Maryland: Amer. Journ. Sci., ser. 3, vol. 45, pp. 407-419.
- DAVIDSON, T.—1869. Notes on Continental Geology and Palaeontology (Cretaceous): Geol. Mag., vol. 6, pp. 251-263.
- DAWSON, J. W.—1873. Note on the Fossil Plants from British Columbia, collected by Mr. James Richardson in 1872: Geol. Surv. Canada, Rep. Progress, 1872-3, pp. 66-71, unnumb. pl.
- 1874. Note on Fossil Woods from British Columbia, collected by Mr. Richardson: Amer. Journ. Sci., ser. 3, vol. 7, pp. 47-51.
- 1883. On the Cretaceous and Tertiary Floras of British Columbia and the North-West Territory: Proc. & Trans. Roy. Soc. Canada, vol. 1, pp. 15-34, pls. i-viii.
- 1884. The Cretaceous Flora of North America: Nature, vol. 30, pp. 631-632.

- DAWSON, J. W.—1885. A modern type of plant in the Cretaceous: *Science*, vol. 5, p. 514, text-fig.
- 1886. On the Mesozoic Floras of the Rocky Mountain Region of Canada: *Proc. & Trans. Roy. Soc. Canada*, vol. 3, sect. iv, pp. 1-22, pls. i-iv.
- 1887. On the Fossil Plants of the Laramie Formation of Canada: *Proc. & Trans. Roy. Soc. Canada*, vol. 4, sect. iv, pp. 19-32, pls. i, ii.
- 1888. Note on Fossil Woods and other Plant Remains, from the Cretaceous and Laramie Formations of the Western Territories of Canada: *Proc. & Trans. Roy. Soc. Canada*, vol. 5, sect. iv, pp. 31-37.
- 1890. On Fossil Plants collected by Mr. R. A. McConnell, on Mackenzie River, and by Mr. T. O. Weston, on Bow River: *Proc. & Trans. Roy. Soc. Canada*, vol. 7, pp. 69-74, pls. x, xi.
- 1893. On the Correlation of early Cretaceous Floras in Canada and the United States, and on some new plants of this period: *Proc. & Trans. Roy. Soc. Canada*, vol. 10, pp. 79-93.
- 1894. On New Species of Cretaceous Plants from Vancouver Island: *Proc. & Trans. Roy. Soc. Canada*, vol. 11, sect. iv, pp. 53-72, pls. v-xiv.
- 1905. *The Geological History of Plants*, 290 pp. London. [1st ed. 1888.]
- and DAWSON, G. M.—1889. On Cretaceous Plants from Port McNeill, Vancouver Island: *Trans. Roy. Soc. Canada*, vol. 6, sect. iv, pp. 71, 72.
- DEBEY, M. H.—1848 A. Uebersicht der urweltlichen Pflanzen des Kreidegebirges überhaupt und der Aachener Kreideschichten insbesondere: *Verhandl. naturhist. Vereins preuss. Rheinland*, 5th year, pp. 113-125.
- 1848 B. Ueber eine neue Gattung urweltlicher Coniferen aus dem Eisensand der Aachener Kreide: *Verhandl. naturhist. Vereins preuss. Rheinland*, 5th year, pp. 126-142.
- 1849. Entwurf zu einer geognostisch-geogenetischen Darstellung der Gegend von Aachen: *Amtlich. Bericht Ges. Deutsch. Naturforsch. Ärzte Versamml. in Aachen 1847*, pp. 269-328.
- 1850. Uebersicht der urweltlichen Pflanzen des Kreide-Gebirges überhaupt und der Aachener Kreide-Schichten insbesondere: *Neues Jahrb. f. Min.*, pp. 115-117. [Referat.]
- 1850 A. Ueber eine neue Gattung urweltlicher Coniferen aus dem Eisensande der Aachener Kreide: *Neues Jahrb. f. Min.*, pp. 117-118.
- 1851. Beitrag zur fossilen Flora der holländischen Kreide: *Verhandl. naturhist. Vereins preuss. Rheinland*, 8th year, pp. 568-569.
- 1857. See ETTINGSHAUSEN and DEBEY.—1857.
- 1858. Ueber die fossile Flora der Kreideformation der Umgebungen von Aachen und Maestricht: *Amtlich. Bericht 32. Versamml. Deutsch. Naturforsch. Ärzte in 1856*, pp. 142-144.

- DEBEY, M. H.—1858 A. See ETTINGSHAUSEN and DEBEY.—1858.
 — 1865. [On Cretaceous plants of Aachen:] Correspondenzblatt no. 2, Verhandl. naturhist. Vereins preuss. Rheinland, 22nd year, pp. 56-58.
 — 1877. [On Fossil Conifers of the Aachen Chalk:] Correspondenzblatt no. 1, Verhandl. naturhist. Vereins preuss. Rheinland, 34th year, p. 110.
 — 1881. Sur les Feuilles Querciformes des sables d'Aix-la-Chapelle: Comptes Rendus Congrès Bot. Horticult., pt. 2, pp. 83-97, pl.
 — and ETTINGSHAUSEN, C. von.—1859 A. Die urweltlichen Thallophten des Kreidegebirges von Aachen und Maestricht: Denkschr. k. Akad. Wiss. Wien, vol. 16, pp. 131-214, pls. i-iii.
 — and ETTINGSHAUSEN, C. von.—1859 B. Die urweltlichen Aerobryen des Kreidegebirges von Aachen und Maestricht: Denkschr. k. Akad. Wiss. Wien, vol. 17, pp. 183-248, pls. i-vii.
 DEECKE, W.—1883. Ueber einige neue Siphoneen: Neues Jahrb. f. M. n., vol. 1, pp. 1-14, pl. i.
 — 1901. Ueber *Hexagonaria* v. Hag. und *Goniolina* Rœm.: Centralbl. f. Min., no. 15, pp. 469-473, text-figs. 1, 2.
 DEWALQUE, G.—1880. Fragments Paléontologiques. Sur une Algue nouvelle de la Craie: Ann. Soc. géol. Belgique, vol. 8, pp. 43-44, pl. i.
 DILLER, J. S.—1908. Strata containing the Jurassic Flora of Oregon: Bull. Geol. Soc. Amer., vol. 19, pp. 367-402 [see Cret. plants by Knowlton, pp. 384-387].
 DILLER, J. S., and STANTON, T. W.—1894. The Shasta-Chico Series: Bull. Geol. Soc. Amer., vol. 5, pp. 435-464.
 DIXON, F.—1850. The Geology and Fossils of the Tertiary and Cretaceous Formations of Sussex, 422 pp., xl pls., London. Also ed. 2, revised and augmented by T. Rupert Jones, Brighton, 1878.
 DOLLO, L.—1888. *Achenosaurus multidentis*: Bull. Soc. belge Géol., vol. 2, p. 300.
 DORMITZER.—1853. See KREPI, M.—1853.
 DUNKER, W.—1846 Monographie der Norddeutschen Wealdenbildung. Ein Beitrag zur Geognosie und Naturgeschichte der Vorwelt, 63 pp., xxi pls. Brunswick.
 — 1856. Ueber mehre Pflanzenreste aus dem Quadersandstein von Blankenburg: Palaeontographica, vol. 4, pp. 179-183, pls. xxxii-xxxv.
 EHRENBURG, C.—1841. Ueber noch jetzt zahlreich lebende Thierarten der Kreidebildung und den Organismus der Polythalamien: Abhandl. k. preuss. Akad. Wiss., 1839, pp. 81-174, pls. i-iv.
 EICHWALD, E. von.—1853. Einige palaeontologische Bemerkungen über den Eisensand von Kursk: Bull. Soc. Imp. Nat. Moscou, vol. 26, pt. i, pp. 209-231.
 — 1860. Lethaea Rossica, Plants in vol. 1, pp. 268, pls. xxi. Stuttgart.
 — 1861. Der Grünsand in der Umgegend von Moskwa: Bull. Soc. Imp. Nat. Moscou, vol. 34, pp. 278-313.

- ERCHWALD, E. VON.—1862. Die Vorweltliche Fauna und Flora des Grünsandes der Gegend von Moskwa: Bull. Soc. Imp. Nat. Moscou, vol. 35, no. 2, pp. 355–410.
- 1865–68. *Lethæa Rossica*, vol. 2, pp. 1304, pls. iii (Cret.). Stuttgart.
- ELDRIDGE, G. H.—1896. See EMMONS, CROSS, and ELDRIDGE—1896.
- EMMONS, S. F., CROSS, W., and ELDRIDGE, G. H.—1896. Geology of the Denver Basin in Colorado: Mon. U. S. Geol. Surv., no. 27, pp. 1–527, pls. i–xxx.
- ENDLICHER, S.—1840. *Genera Plantarum secundum ordines naturales disposita*, 1483 pp. Vienna. Also Supplement (114 pp.), 1842.
- 1847 A. *Synopsis Coniferarum fossilium*, 52 pp. Sangalli.
- 1847 B. *Synopsis Coniferarum*, 368 pp. Sangalli.
- ENGELHARDT, H.—1885. Die Crednerien im unteren Quader Sachsens: Festschrift naturwiss. Ges. Isis Dresden, pp. 55–62, pl. i.
- 1892. Ueber boehmische Kreidepflanzen aus dem Geolog. Institute der deutschen Universität Prag: Mitteil. Osterland. herausgegeb. Naturforsch. Ges. Osterlandes, vol. 5, pp. 86–118, pl. i.
- 1892 A. Ueber Kreidepflanzen von Niederschöna: Abhandl. naturwiss. Ges. Isis Dresden, 1891, pp. 79–105, pl. ii.
- 1897. See VANHOFFEN, G.—1897.
- ETHERIDGE, R.—1904. An Endophyte (*Stichus mermisoides*) occurring in the test of a Cretaceous Bivalve: Rec. Australian Mus., vol. 5, no. 4, pp. 255–257, pls. xxx, xxxi.
- ETTINGSHAUSEN, C. VON.—1851. Die Proteaceen der Vorwelt: Sitzb. k. Akad. Wiss. Wien, vol. 7, pt. x, pp. 711–745, pls. xxx–xxxiv.
- 1852. Ueber fossile Pandaneen: Sitzb. k. Akad. Wiss. Wien, vol. 8, pp. 489–495, pls. xxiii–xxvi.
- 1852 A. Ueber fossile Proteaceen: Sitzb. k. Akad. Wiss. Wien, vol. 9, pp. 820–824, pls. lvii–lviii.
- 1852 B. Ueber *Palæobromelia*, ein neues fossiles Pflanzengeschlecht: Abhandl. k. k. geol. Reichsanst., vol. 1, pt. 3, no. 1, pp. 1–10, pls. i, ii.
- 1852 C. Beitrag zur näheren Kenntniss der Flora der Wealdenperiode: Abhandl. k. k. geol. Reichsanst., vol. 1, pt. 3, no. 2, pp. 1–32, pls. i–v.
- 1853 A. Die Tertiär-Flora der Oesterreichischen Monarchie, no. 1. —Fossile Flora von Wien: Abhandl. k. k. geol. Reichsanst., vol. 2, pt. 3, no. 1, pp. 1–36, pls. i–v.
- 1853 B. Die Tertiäre-Flora von Häring in Tirol: Abhandl. k. k. geol. Reichsanst., vol. 2, pt. 3, no. 2, pp. 1–118, pls. i–xxxi.
- 1859 A. See DEBEY and ETTINGSHAUSEN.—1859 A.
- 1859 B. See DEBEY, M. H., and ETTINGSHAUSEN, C. VON.—1859 B.
- 1863. Die fossilen Algen des Wiener und des Karpathen-Sandsteines: Sitzb. k. Akad. Wiss. Wien, vol. 48, pp. 444–467, pls. i, ii.
- 1867 A. Die Kreideflora von Niederschöna in Sachsen, ein Beitrag zur Kenntniss der ältesten Dicotyledonengewächse: Sitzb. k. Akad. Wiss. Wien, vol. 55, pt. 1, pp. 235–264, pls. i–iii.
- 1867 B. Die fossile Flora des Tertiärbeckens von Bilin, pt. i: Denkschr. k. Akad. Wiss. Wien, vol. 26, pp. 79–174, pls. i–xxx.

- ETTINGSHAUSEN, C. von.—1872. Die fossile Flora von Sazor in Krain: Denkschr. k. Akad. Wiss. Wien, vol. 32, pp. 159-202, pls. i-x.
- 1874. Die Florenelemente in der Kreideflora: Sitzb. k. Akad. Wiss. Wien, vol. 69, pt. 1, pp. 510-518.
- 1887. On the Fossil Flora of New Zealand: Geol. Mag., dec. 3, vol. 4, pp. 363-367.
- 1887 A. Beiträge zur Kenntniss der Fossilen Flora Neuseelands: Denkschr. k. Akad. Wiss. Wien, vol. 53, pp. 143-192, pls. i-ix.
- 1888. Contributions to the Tertiary Flora of Australia: Mem. Geol. Surv. New South Wales, Palaeont. no. 2, 189 pp., xv pls.
- 1891. Contributions to the Knowledge of the Fossil Flora of New Zealand: Trans. New Zealand Institute, vol. 23, pp. 237-310, pls. xxiv-xxxii (see p. 286 *et seq.*). [Descriptions of species translated from Denkschr. k. Akad. Wiss. Wien, vol. 53, 1887.]
- 1893. Ueber fossile Pflanzenreste aus der Kreideformation Australiens: Sitzb. k. Akad. Wiss. Wien, vol. 102, pp. 126-151.
- 1895. Beiträge z. Kennt. d. Kreideflora Australiens: Denkschr. k. Akad. Wiss. Wien, vol. 62, pp. 1-57, pls. i-iv.
- 1896. Ueber die Kreideflora der südlichen Hemisphäre: Mittheil. Naturwiss. Ver. Steiermark, vol. 32, pp. 155-164.
- 1896 A. Ueber die Nervation der Blätter bei der Gattung *Quercus* mit besonderer Berücksichtigung ihrer vorweltlichen Arten: Denkschr. k. Akad. Wiss. Wien, vol. 63, pp. 117-180, pls. i-xii.
- and DEBEY, M. H.—1857. Die urweltlichen Thallophyten des Kreidegebirges von Aachen und Maestricht: Sitzb. k. Akad. Wiss. Wien, vol. 25, pp. 507-512.
- 1858. Die vorweltlichen Acrobryen des Kreidegebirges von Aachen und Maestricht: Sitzb. k. Akad. Wiss. Wien, vol. 27, pp. 167-170.
- 1859 A. See DEBEY and ETTINGSHAUSEN.—1859 A.
- 1859 B. See DEBEY and ETTINGSHAUSEN.—1859 B.
- FEISTMANTEL, O.—1870. Ueber die Reste der Kreideformation bei Kuchelbad: Sitzb. k. böhm. Ges. Wiss., 1870, pp. 73-75.
- 1872. Ueber Baumfarnenreste der böhmischen Steinkohlen-, Perm- und Kreideformation: Abhandl. k. böhm. Ges. Wiss., vol. 5, pp. 1-30, pls. i, ii.
- 1874. Vorbericht über die Peraccer Kreideschichten in Böhmen und ihre fossilen Reste: Sitzb. k. böhm. Ges. Wiss., 1874, pp. 253-276.
- 1877. Notes on Fossil Floras of India.—XIV. On a tree fern stem from the Cretaceous Rocks near Trichinopoly in Southern India: Rec. Geol. Surv. India, vol. 10, pp. 133-140, plate.
- FELIX, J.—1894. Studien über fossile Pilze: Zeitschr. deutsch. geol. Ges., vol. 46, pp. 269-280, pl. xix.
- and NATHORST, A. G.—1893. Versteinerungen aus dem mexicanischen Staat Oaxaca: Beitr. Geol. Palaeont. Republ. Mexico, pt. ii, pp. 41-54, pl. iii. Leipzig.

- FISCHER, G., DE WALDHEIM.—1826. Notice sur les végétaux fossiles du Gouvernement de Moscou: Seance publ. Soc. Imp. Nat. Imp. Universit. Moscou, pp. 1-23, pl. i.
- FISCHER-OOSTER, C. von.—1858. Die fossilen Fucoiden der Schweizer-Alpen, nebst Brörterungen über deren geologisches Alter, 72 pp., xviii pls. Berne.
- FISHER, C. A.—1909. Geology of the Great Falls Coalfield, Montana: Bull. U.S. Geol. Surv., no. 356, pp. 1-85 [see pp. 33-36].
- FITTON, W. H.—1836. Observations on some of the Strata between the Chalk and the Oxford Oolite, in the South-East of England: Trans. Geol. Soc., ser. 2, vol. 4, pp. 103-388, pls. xi-xxiii.
- FLICHE, P.—1892. Sur une dicotylédone trouvée dans l'Albien supérieur, aux environs de Sainte-Menehould (Marne): Comptes Rendus Acad. Sci. Paris, vol. 114, pp. 1084-1085.
- 1893. Sur un nouveau genre de Conifère rencontré dans l'Albien de l'Argonne: Comptes Rendus Acad. Sci. Paris, vol. 116, pp. 1002-1004.
- 1894. Sur des fruits de Palmiers trouvés dans le Cénomanien aux environs de Sainte-Menehould: Comptes Rendus Acad. Sci. Paris, vol. 118, pp. 889-890.
- 1896. Études sur la Flore fossile de l'Argonne Albien-Cénomanien: Bull. Soc. Sci. Nancy, ser. 2, vol. 14, pp. 114-306, pls. i-xvii.
- 1896 a. Note sur les nodules et les bois minéralisés trouvés à Saint-Parres-les-Vaudes (Aube) dans les Grès verts infracrétacés: Mém. Soc. Acad. Aube, vol. 33, pp. 177-189, pl. iv.
- 1900. Contribution à la Flore fossile de la Haute-Marne (Infracrétacé): Bull. Soc. Sci. Nancy, ser. 2, vol. 16, pp. 11-31, pls. i, ii, & unnumb. pl.
- 1900 a. Note sur un bois fossile de Madagascar: Bull. Soc. géol. France, ser. 3, vol. 28, pp. 470-472, text-fig. 1.
- 1902. Note sur un *Zosterites* trouvé dans le Crétacé supérieur du Dévoluy: Bull. Soc. géol. France, ser. 4, vol. 2, pp. 112-126, pl. ii.
- 1905. Note sur des bois fossiles de Madagascar: Bull. Soc. géol. France, ser. 4, vol. 5, pp. 346-358, pl. x.
- FONTAINE, W. M.—1879. Notes on the Mesozoic Strata of Virginia: Amer. Journ. Sci., ser. 3, vol. 17, pp. 229-239.
- 1889. The Potomac or Younger Mesozoic Flora: Mon. U. S. Geol. Surv., no. 15, 377 pp., clxxx pls.
- 1893. Description of some Fossil Plants from the Great Falls Coalfield of Montana: Proc. U. S. Nat. Mus., vol. 15, pp. 487-495, pls. lxxxii-lxxxiv.
- 1894. Notes on some Fossil Plants from the Trinity Division of the Comanche series of Texas: Proc. U.S. Nat. Mus., vol. 16, pp. 261-282, pls. xxxvi-xliii.
- 1894 a. Flora of the Shasta-Chico series, in DILLER and STANTON,—1894.

- FONTAINE, W. M.—1895. See STANTON, T. W.—1895.
- 1896. The Potomac Formation in Virginia: *Bull. U. S. Geol. Surv.*, no. 145, pp. 1-147, & map.
- 1898. See WEED, W. H., and PIRSSON, L. V.—1898.
- 1899. Notes on Lower Cretaceous Plants from the Hay Creek Coalfield, Crook County, Wyoming: *U. S. Geol. Surv. 19th Ann. Rep.*, pt. ii, pp. 645-702, pls. clx-clxix.
- 1905 a. Notes on some Lower Cretaceous (Kootanie) Plants from Montana, in L. F. Ward's Status of Mesozoic Floras: *Mon. U. S. Geol. Surv.*, no. 48, pp. 281-315, pls. lxxi-lxxiii.
- 1905 b. Report on various collections of Fossil Plants from the Older Potomac of Virginia and Maryland, in L. F. Ward's Status of Mesozoic Floras: *Mon. U. S. Geol. Surv.*, no. 48, pp. 476-503, pls. cvii-cix.
- FORD, S. O.—1906. See SEWARD and FORD.—1906.
- FRAAS, O.—1867. *Aus dem Orient*, pt. 1, p. 40. Stuttgart. [Reprinted from *Württ. Jahresh.*, vol. 23.]
- 1878. *Aus dem Orient*, pt. 2, p. 43. Stuttgart. [Reprinted from *Württ. Jahresh.*, vol. 34.]
- FRITEL, P. H.—1903. *Histoire naturelle de la France*. Pt. 24, *Paléo botanique (Plantes Fossiles)*, 347 pp., xxxvi pls. Paris.
- 1908. *Étude sur les Nymphéacées fossiles*, pts. i-iii: *Le Naturaliste*, vol. 30, pp. 53-56; pp. 137-139; pp. 149-150.
- 1910. See COUVAT, J., and FRITEL, P. H.—1910.
- FRITSCH, A.—1877. Studien im Gebiete der böhmischen Kreideformation. Paläontologische Untersuchungen der einzelnen Schichten. II. Die Weissenberger und Malnitzer Schichten: *Archiv. Naturwiss. Landesdurchf. Böhmen*, vol. 4, pp. 1-152.
- 1883. Studien im Gebiete der böhmischen Kreideformation. Paläontologische Untersuchungen der einzelnen Schichten. III. Die Iserschichten: *loc. cit.*, vol. 5, pp. 1-137.
- 1889. Studien im Gebiete der böhmischen Kreideformation. Paläontologische Untersuchungen der einzelnen Schichten. IV. Die Teplitzer Schichten: *loc. cit.*, vol. 7, pp. 1-119.
- 1893. Studien im Gebiete der böhmischen Kreideformation. Paläontologische Untersuchungen der einzelnen Schichten. V. Priesener Schichten [Plants by Bayer, see p. 128]: *loc. cit.*, vol. 9, pp. 1-134.
- 1897. Studien im Gebiete der böhmischen Kreideformation. Paläontologische Untersuchungen der einzelnen Schichten. VI. Die Chlumek Schichten [Plants by Bayer, see pp. 72-83]: *loc. cit.*, vol. 10, pp. 1-83.
- and BAYER, E.—1901. Studien im Gebiete der böhmischen Kreideformation: *loc. cit.*, vol. 11, no. 2, pp. 1-180, text-figs.
- FRÜH, J.—1890. Zur Kenntniss der Gesteinbildenden Algen der schweizer Alpen, mit besonderer Berücksichtigung des Säntisgebietes: *Abhandl. schweiz. paläont. Ges.*, vol. 17, pp. 1-32, pl. i.

- FUCHS, T.—1895. Studien über Fucoiden und Hieroglyphen: Denkschr. k. Akad. Wiss. Wien, vol. 62, pp. 369-448, pls. i-ix.
- FUJII, K.—1909. See STOKES, M. C., and FUJII, K.—1909.
- 1910 A. See STOKES, M. C., and FUJII, K.—1910.
- 1910 B. Some Remarks on the Cretaceous Fossil Flora and the Causes of Extinction: Bot. Mag. Tokyo, vol. 23, no. 284, pp. 197-220.
- GARDNER, J. S.—1884. On the Relative Ages of the American and the English Cretaceous and Eocene Series: Geol. Mag., dec. 3, vol. 1, pp. 492-506.
- 1886 A. On Mesozoic Angiosperms: Geol. Mag., dec. 3, vol. 3, pp. 193-204, pl. v.
- 1886 B. Second Report of the Committee . . . for . . . reporting on the Fossil Plants of the Tertiary and Secondary Beds of the United Kingdom: Rep. British Assoc., Birmingham, pp. 241-250 and pl. [Also in Geol. Mag., 1886, p. 495.]
- GEINITZ, H.—1901. *Aphrocallistes* (*Hexagonaria*) als Senongeschiebe: Centralbl. f. Min., no. 19, pp. 584-585.
- GEINITZ, H. B.—1842. Charakteristik der Schichten und Petrefakten des sächsisch-böhmischen Kreidegebirges. Pt. 3. Die sächsisch-böhmische Schweiz, die Oberlausitz und das Innere von Böhmen, pp. i-xxii, 63-116, pls. xvii-xxiv. Leipzig and Dresden. [Ed. 2 in 1850.]
- 1850 A. Das Quadersandsteingebirge, oder Kreidegebirge in Deutschland, 292 pp., xii pls. Freiberg.
- 1850 B. Das Quadergebirge, oder die Kreideformation in Sachsen, mit besonderer Berücksichtigung der Glaukonitreichen Schichten: Fürstl. Jablon. Ges. Preisschrift, pp. 1-44 & 1 pl. Leipzig.
- 1850 D. Ueber die Zusammensetzung und Lagerung der Kreideformation in der Gegend zwischen Halberstadt, Blankenburg, und Quedlinburg: Neues Jahrb. f. Min., pp. 133-138.
- 1875 A. Das Elbthalgebirge in Sachsen. Erster Theil. Der untere Quader. IX. Pflanzenreste des unteren Quaders: Paläontogr., vol. 20, part 1, pp. 304-310, pls. lxi-lxvii.
- 1875 B. Das Elbthalgebirge in Sachsen. Zweiter Theil. Der mittlere und obere Quader: Paläontogr., vol. 20, part 2, pp. 231-233.
- 1879. Ueber zwei neue Kreide-Pflanzen: Neues Jahrb. f. Min., pp. 113-115, pl. iv.
- GERRY, E.—1910. The Distribution of the "Bars of Sanio" in the Coniferales: Annals Bot., vol. 24, pp. 119-124, pl. xiii.
- GIRTY, G. H.—1899. See HAGUE, A., et alii.—1899.
- GLOCKER, E. F.—1841. Ueber die kalkführende Sandsteinformation auf beiden Seiten der mittleren March, in der Gegend zwischen Kwassitz und Kremsier: Nova Acta Acad. Caes. Leop.-Carol., vol. 19, suppl. pp. 309-334, pl. iv.
- GOEPPERT, H. R.—1836. Die fossilen Farnkräuter: Nova Acta Acad. Caes. Leop.-Carol., vol. 17, suppl., pp. 1-486, pls. i-xliv.
- 1839. Bemerkungen über die des Geschiebe im nördlichen Deutschland vorkommenden versteinerten Hölzer: Neues Jahrb. f. Min., 1839, pp. 518-521, pl. 8 B.

- GOEPPERT, H. R. — 1842 A. Ueber die fossile Flora der Quadersandsteinformation in Schlesien, als erster Beitrag zur Flora der Tertiaergebilde: *Nova Acta Acad. Caes. Leop.-Carol.*, vol. 19, pp. 97–134, pls. xlv–liii.
- 1842 B. Fossile Pflanzenreste des Eisensandes von Aachen, als zweiter Beitrag zur Flora der Tertiaergebilde: *Nova Acta Acad. Caes. Leop.-Carol.*, vol. 19, pt. 2, pp. 137–160, pl. liv.
- 1844. Ueber die fossilen Cycadeen überhaupt, mit Rücksicht auf die in Schlesien vorkommenden Arten: *Uebers. schles. Ges. Vaterländ. Kultur* in 1843, pp. 114–144, pl. i.
- 1845. Uebersicht der fossilen Flora Schlesiens. See WIMMER, F. — 1845.
- 1847. Zur Flora des Quadersandsteins im Schlesien, als Nachtrag zu der früher erschienenen Abhandlung über denselben Gegenstand: *Nova Acta Acad. Caes. Leop.-Carol.*, vol. 22, pp. 355–365, pls. xxxv.–xxxviii.
- 1848. Zur Flora des Quadersandsteins im Schlesien, als Nachtrag zu der Abhandlung in *N. Acta Acad. C. Leop. Nat. Cur.* xix, 1841: *Neues Jahrb. f. Min.*, pp. 269–278.
- 1850 A. Monographie der fossilen Coniferen. Pp. 1–286, pls. i–lviii; Anhang, pp. 1–73. Preisschrift, Leiden.
- 1865. Ueber die fossile Kreideflora und ihre Leitpflanzen: *Zeitschr. deutsch. geol. Ges.*, vol. 17, pp. 638–648.
- 1865 A. Die fossile Flora der Permischen Formation: *Palaeontogr.*, vol. 12, pp. 1–316, pls. i–lxiv [see p. 259].
- 1865 B. Ueber das Vorkommen von Baumfarn in der fossilen Flora, insbesondere in der Kreideformation: *Neues Jahrb. f. Min.*, pp. 395–399.
- 1865 C. Beiträge zur Kreideflora und ihre gegenwärtige geognostische Bedeutung: *Nova Acta Acad. Caes. Leop.-Carol.*, vol. 32, part 1, Heft 5, pp. 11, 12.
- 1866. Beiträge zur Kenntniss fossiler Cycadeen: *Neues Jahrb. f. Min.*, pp. 129–135, pl. ii.
- 1881. *Arboretum fossile. Sammlung von Dünnschliffen fossiler Coniferen-Hölzer der paläozoischen Formation.* [Actual slides of fossil plants.] Descript., pp. 1–6. Breslau.
- GOLDFUSS, A. — 1826. *Petrefacta Germaniae*, part 1. Pp. 1–242, pls. i–lxxi. Düsseldorf.
- GOTHAN, W. — 1904. *Rhizodendron oppolense* Göppert (nach der anatomischen Struktur) *Alsophylla* sp. (nach der Oberflächenskulptur des Stammes): in Hef. 2, *Potonic, Abbild. u. Beschreib. foss. Pflanzenreste*, pp. 1–12, text-figs. Berlin.
- 1905. Zur Anatomie lebender und fossiler Gymnospermen-Hölzer: *Abhandl. k. preuss. geol. Landesanst.*, vol. 44, pp. 1–108.
- 1908. Die Frage der Klimadifferenzierung im Jura und in der Kreideformation im Lichte paläobotanischer Tatsachen: *Jahrb. k. preuss. geol. Landesanst.*, vol. 29, pp. 220–242, pls. xvi–xix.

- GOTHAN, W.—1908 A. Die fossilen Hölzer von der Seymour- und Snow-Inseln: Wiss. Ergebn. Schwed. Sudpol. Exped. 1901-03, vol. 3, pt. 8, pp. 33, pls. ii. Stockholm.
- 1910. *Weichselia reticulata*: in lief. 7, Potonié, Abbild. u. Beschreib. foss. Pflanzenreste, pp. 1-14, text-figs. Berlin.
- GOWAN, J.—1900. See SEWARD and GOWAN.—1900.
- GREGORY, J. W.—1897. Some problems of Arctic Geology, II. Former Arctic Climates: Nature, vol. 56, pp. 351-352.
- GRIEPENKERL, O.—1889. Die Versteinerungen der senonen Kreide von Königsutter im Herzogthum Braunschweig: Palaeont. Abhandl., vol. 14, pp. 1-116, pls. i-xii [see p. 14].
- GÜMBEL, C. W.—1861. Geognostische Beschreibung des bayerischen Alpengebirges und seines Vorlandes. Pp. 1-950, pls. i-xlii. Gotha.
- 1871. Die sogenannten Nulliporen (*Lithothamnium* und *Dactylopora*) und ihre Betheiligung an der Zusammensetzung der Kalkgesteine: Abhandl. k. Akad. Wiss. München, vol. 11, pt. 1, pp. 11-52, pls. i, ii; & pp. 232-290, pls. Di-Div.
- 1896. Vorläufige Mittheilung über Flyschalgen: Neues Jahrb. f. Min., vol. 1, pp. 227-232.
- GÜRICH, G.—1885. Ein neues fossiles Holz aus der Kreide Armeniens, nebst Bemerkungen über paläozoische Hölzer: Zeitschr. deutsch. geol. Ges., vol. 37, pp. 433-440.
- GWYNNE-VAUGHAN, D. T.—1908. See KIDSTON and GWYNNE-VAUGHAN.—1908.
- HAAST, J. von.—1887. Notes on the Age and Subdivisions of the Sedimentary Rocks in the Canterbury Mountains, based upon the Palaeontological Researches of Professor Dr. C. Baron von Ettingshausen in Gratz (Austria): Trans. & Proc. New Zealand Inst., ser. 2, vol. 19, pp. 449-451.
- HAGUE, A., IDINGS, J. P., WEED, W. H., WALCOTT, C. D., Girty, G. H., STANTON, T. W., KNOWLTON, F. H.—1899. Geology of the Yellowstone National Park, part ii: Mon. U.S. Geol. Surv., no. 32, pp. 1-791, pls. lxxvii-cxli.
- HAMPE, E.—1852. Vortrag über Petrefacten der Kreideformation (Q. S. St.) bei Blankenburg: Bericht naturwiss. Ver. Harzes, 1852, pp. 6, 7.
- 1855. [Fossil Plants from Quadersandstein]: Bericht naturwiss. Ver. Harzes, 1853-54, p. 12.
- HARTIG, T.—1848. Beiträge zur Geschichte der Pflanzen und zur Kenntniss der nord-deutschen Braunkohlen-Flora: Bot. Zeit., vol. 6, pp. 166-172.
- HATCHER, J. B.—1901. *Sabal rigida*; a new species of Palm from the Laramie: Ann. Carnegie Mus., vol. 1, pp. 263-264, text-fig.
- 1903. Osteology of *Haplocanthosaurus* with Description of a New Species, and Remarks on the probable Habits of the Sauropoda and the Age and Origin of the *Atlantosaurus* Beds: Mem. Carnegie Mus., vol. 2, no. 1, pp. 1-75, pls. i-vi.
- 1905. See STANTON and HATCHER.—1905.

- HAUPTFLEISCH, P.—1897. Die als fossile Algen beschriebenen Pflanzenreste oder Abdrücke : in Engler's Pflanzenfamilien, I, pt. 2, pp. 543-569.
- HAYDEN, F. V.—1859 A. See MEEK, F. B., and HAYDEN, F. V.—1859.
- 1859 B. See MEEK and HAYDEN.—1859 B.
- 1859 C. See MEEK and HAYDEN.—1859 C.
- 1869. Geological Report of the Exploration of the Yellowstone and Missouri Rivers under the direction of Captain W. F. Reynolds, Corps of Engineers, 1859-60. Washington.
- 1871. Preliminary Report of the United States Geological Survey of Wyoming. 2nd Ann. Rep. [=4th Ann. Rep.]. Pp. 1-511. Washington.
- 1872. Preliminary Report of the United States Geological Survey of Montana. 5th Ann. Rep. Pp. 1-538. Washington.
- 1873. Sixth Annual Report of the United States Geological Survey of the Territories. Pp. 1-844. Washington.
- HECTOR, J.—1879. On the Fossil Flora of New Zealand (Abstract): Trans. & Proc. New Zealand Inst., vol. 11, pp. 536-537.
- HEER, O.—1859. See MEEK and HAYDEN.—1859.
- 1859 A. See LESQUEREUX.—1859.
- 1861. Reply to Dr. Newberry on the Age of the Nebraska Leaves: Amer. Journ. Sci., ser. 2, vol. 31, pp. 435-440.
- 1863. On the Fossil Flora of Bovey Tracey: Phil. Trans. Roy. Soc. Lond., vol. 152, pp. 1039-1086, pls. lv-xxi.
- 1865. Die Umwelt der Schweiz. Zürich. [Die Zeit der Kreidebildung, pp. 167-220.]
- 1867. See CAPELLINI, J., and HEER, O.—1867.
- 1868. Flora Fossilis Arctica.—I. Die fossile Flora der Polarländer enthaltend die in Nordgrönland, auf der Melville-Insel, im Banksland, am Mackenzie, in Island und Spitzbergen entdeckten fossilen Pflanzen. Pp. 1-192, pls. i-i. Zürich.
- 1869 A. Beiträge zur Kreide-Flora.—I. Flora von Modetuin in Mähren: Neue Denkschr. allgem. schweizerisch. Ges. Naturwiss., vol. 23, pp. 1-24, pls. i-xi.
- 1870. Contributions to the Fossil Flora of North Greenland, being a Description of the Plants collected by Mr. Edward Whymper during the summer of 1867: Phil. Trans. Roy. Soc. Lond., vol. 159, pp. 445-488, pls. xxxix-lvi (Miocene plants). [Flora Fossilis Arctica, vol. 2.]
- 1870 A. Die Miocene Flora und Fauna Spitzbergens: K. Svensk. Vet.-Akad. Handl., vol. 8, pp. 1-98, pls. i-xvi.
- 1871. Förutskickade anmärkningar öfver Nordgrönlands Kritflora, grundade på den svenska expeditionens upptäckter, 1870: Öfvers. K. Svensk. Vet.-Akad., vol. 28, pp. 1175-1184.
- 1871 A. Beiträge zur Kreide-Flora.—II. Zur Kreide-Flora von Quedlinburg: Neue Denkschr. allgem. schweiz. Ges. gesamt. Naturwiss., vol. 24, pp. 1-15, pls. i-iii.
- 1871 B. Mittheilungen an Professor H. B. Geinitz: Neues Jahrb. f. Min., pp. 394-395.

- HEER, O.—1872. Vorläufige Bemerkungen über die Kreideflora Nordgrönlands, gegründet auf die Entdeckungen der schwedischen Expedition vom Jahre 1870: *Zeitschr. deutsch. geol. Ges.*, vol. 24, pp. 155–164. [Translation of Heer, 1871.]
- 1874 A. Die Kreide-Flora der Arctischen Zone, gegründet auf die von den Schwedischen Expeditionen von 1870 und 1872 in Grönland und Spitzbergen gesammelten Pflanzen: *K. Svensk. Vet.-Akad. Handl.*, vol. 12, pp. 1–138, pls. i–xxxviii. [*Flora Fossilis Arctica*, vol. 3.]
- 1874 B. Beiträge zur Steinkohlen-Flora der Arctischen Zone: *K. Svensk. Vet.-Akad. Handl.*, vol. 12, pp. 1–11, pls. i–vi [p. 8, pls. v, vi for Cretaceous plants]. [*Flora Fossilis Arctica*, vol. 3.]
- 1874 C. Ueber Ginkgo Thunberg: *Gartenflora*, vol. 23, pp. 260–262, pl. lxxvii.
- 1876 A. Beiträge zur fossilen Flora Spitzbergens, gegründet auf die Sammlungen der Schwedischen Expedition von Jahre 1872 auf 1873: *K. Svensk. Vet.-Akad. Handl.*, vol. 14, no. 5, pp. 1–141, pls. i–xxxii [pp. 48–50 for Cretaceous plants]. [*Flora Fossilis Arctica*, vol. 4, pt. 1.]
- 1876 B. Ueber fossile Früchte der Oase Charchah: *Neue Denkschr. allgem. schweiz. Ges. gesamt. Naturwiss.*, vol. 27, pp. 1–11, pl. i.
- 1876 C. Beiträge zur Jura-Flora Ostsibiriens und des Amurlandes: *Mém. Acad. Imp. Sci. St. Pétersbourg*, vol. 22, no. 12, pp. 1–122, pls. i–xxxi. [*Flora Fossilis Arctica*, vol. 4, pt. 2.]
- 1877. *Flora Fossilis Helvetiae*, Die Vorweltliche Flora der Schweiz. Dritte Lieferung. Die Pflanzen des Jura (Fortsetzung), der Kreide und des Eocen. Pp. 101–182, pls. xlv–lxx. Zürich.
- 1877 A. Beiträge zur fossilen Flora Spitzbergens, mit einem Anhang—Uebersicht der Geologie des Eisfjordes und des Bellsundes von Prof. A. E. Nordenskiöld: *Flora Fossilis Arctica*, vol. 4, pt. 1, pp. 1–141, pls. i–xxxii.
- 1878. Beiträge zur fossilen Flora Sibiriens und des Amurlandes: *Mém. Acad. Imp. Sci. St. Pétersbourg*, vol. 25, no. 6, pp. 1–58, pls. i–xv. [*Flora Fossilis Arctica*, vol. 5, pt. 2.]
- 1880. Beiträge zur Miocenen Flora von Nord-Canada: *Flora Fossilis Arctica*, vol. 6, pt. 1, no. 3, pp. 1–17, pls. i–iii.
- 1880 A. Nachträge zur Jura-Flora Sibiriens: *Mém. Acad. Imp. Sci. St. Pétersbourg*, vol. 27, no. 10, pp. 1–34, pls. i–ix.
- 1880 B. Nachträge zur fossilen Flora Grönlands: *K. Svensk. Vet.-Akad. Handl.*, vol. 18, no. 2, pp. 1–17, pls. i–vi.
- 1881. *Contributions à la Flore fossile du Portugal: Comm. Serv. Geol. Portugal*, pp. xiv, 51, pls. i–xxviii & A.
- 1882. Die fossile Flora Grönlands, I: *Flora Fossilis Arctica*, vol. 6, pt. 2, pp. 1–112, pls. i–xlvi.
- 1883 A. Die fossile Flora Grönlands, II: *Flora Fossilis Arctica*, vol. 7, pp. 1–275, pls. xlviii–cix & map [see Section 1, Die Flora der Patootschichten, pp. 1–46].

- HEER, O.—1883 B. *Flora fossilis Grønlandica*. *Afbildinger af Grønlands fossile Flora; Mæddeløser om Grønland*, vol. 5, pls. i-cix & map [plates reprinted in *Flora Fossilis Arctica*, vols. 6, 7].
- HILDEBRAND, F.—1861. *Die Verbreitung der Coniferen in der Jetztzeit und in den früheren geologischen Perioden*: *Verhandl. naturhist. Ver. preuss. Rheinl.*, vol. 18, pp. 199-384, pls. i-iv.
- HILL, R. T.—1893. *Paleontology of the Cretaceous Formations of Texas—the Invertebrate Paleontology of the Trinity Division*: *Proc. Biol. Soc. Washington*, vol. 8, pp. 9-40, pls. i-viii [see p. 39, pl. i].
- 1895 A. *On Outlying Areas of the Comanche Series in Kansas, Oklahoma, and New Mexico*: *Amer. Journ. Sci.*, ser. 3, vol. 50, pp. 205-234.
- 1895 B. *Discovery of a Dicotyledonous Flora in the Cheyenne Sandstone*: *Amer. Journ. Sci.*, ser. 3, vol. 49, p. 473.
- 1901. *Geography and Geology of the Black and Grand Prairies, Texas, with detailed descriptions of the Cretaceous Formations*: *21st Ann. Rep. U.S. Geol. Surv. for 1899-1900* [see pp. 314-318 for Knowlton's Report].
- HILL, W.—1900. See JUKES-BROWNE, A. J., and HILL, W.—1900.
- HISINGER, W.—1837. *Lethæa Svecica seu Petrificata Svecie, Iconibus et Characteribus Illustrata*. Pp. 1-124, pls. i-xxxiv, & pls. xxxv-xxxix supplement. Stockholm.
- HOCHSTETTER, F. VON.—1866. See UNGER, F.—1866.
- HOFMANN, H.—1884. *Untersuchungen über fossile Hölzer*: *Zeitschr. gesamt. Naturwiss.*, vol. 57, pp. 156-195.
- HOLLIICK, A.—1892 A. *The Paleontology of the Cretaceous Formation on Staten Island*: *Trans. New York Acad. Sci.*, vol. 11, pp. 96-104, pls. i-iv.
- * — 1892 B. *Additions to the Cretaceous Flora of Staten Island*: *Proc. Nat. Sci. Assoc. Staten Island* [not seen].
- 1893. *Additions to the Paleobotany of the Cretaceous Formations on Staten Island*: *Trans. New York Acad. Sci.*, vol. 12, pp. 28-39, pls. i-iv.
- 1893 A. *Preliminary Contributions to our Knowledge of the Cretaceous Formation on Long Island and Eastwards*: *Trans. New York Acad. Sci.*, vol. 12, pp. 222-237, pls. v-vii.
- 1893 B. *A New Fossil Palm from the Cretaceous Formation of Glen Cove, Long Island*: *Bull. Torrey Bot. Club*, vol. 20, pp. 168, 169, pl. cxlix.
- 1893 C. *Plant Distribution as a Factor in the Interpretation of Geological Phenomena, with special reference to Long Island and Vicinity*: *Trans. New York Acad. Sci.*, vol. 12, pp. 189-202.
- 1893 D. *Some further Notes upon *Serenopsis Kempii**: *Bull. Torrey Bot. Club*, pp. 334-335, pl. clxvi.
- 1894 A. *Additions to the Paleobotany of the Cretaceous Formation on Long Island*: *Bull. Torrey Bot. Club*, vol. 21, pp. 49-65, pls. clxxiv-clxxx.

- HOLLICK, A.—1894 B. Fossil Salvinias, including Description of a new Species: Bull. Torrey Bot. Club, vol. 21, pp. 253-257, pl. ccv.
- 1894 c. Some further Notes on the Geology of the North Shore of Long Island: Trans. New York Acad. Sci., vol. 13, pp. 122-130, chart.
- 1894 D. A New Fossil *Nelumbo* from the Laramie Group at Florence, Colo.: Bull. Torrey Bot. Club, vol. 21, pp. 307-310.
- 1894 E. Wing-like Appendages on the Petioles of *Liriophyllum populoides*, Lesq., and *Liriodendron alatum*, Newb., with Descriptions of the latter: Bull. Torrey Bot. Club, vol. 21, pp. 467-471, pls. ccxx, ccxxi.
- 1895. Descriptions of New Leaves from the Cretaceous (Dakota Group) of Kansas: Bull. Torrey Bot. Club, vol. 22, pp. 225-228, pls. cccxxvi-cccxxvii.
- 1896 A. Martha's Vineyard Cretaceous Plants: Bull. Geol. Soc. Amer., vol. 7, pp. 12-14.
- 1896 B. Appendages to the Petioles of *Liriodendra*: Bull. Torrey Bot. Club, vol. 23, pp. 249, 250, pls. cclxix, cclxx.
- *— 1896 c. Recent Additions to the Cretaceous Fossil Flora of Staten Island: Proc. Nat. Sci. Assoc. Staten Island, vol. 6 [not seen].
- *— 1897. Notice of finding a *Moriconia cyclotoxolon* in Staten Island: Proc. Nat. Sci. Assoc. Staten Island, vol. 6, p. 30 [not seen].
- 1898 A. Notes on Block Island: Ann. New York Acad. Sci., vol. 11, pp. 55-72, pls. ii-ix.
- 1898 B. Additions to the Palaeobotany of the Cretaceous Formation on Staten Island (New York), no. II.: Ann. New York Acad. Sci., vol. 11, pp. 415-430, pls. xxxvi-xxxviii.
- 1898 c. Geological Notes. Long Island and Block Island: Trans. New York Acad. Sci., vol. 16, pp. 9-18. [See also NEWBERRY, J. S. —1898]
- 1898 D. The Cretaceous Clay Marl Exposure at Clifford, N.J.: Trans. New York Acad. Sci., vol. 16, pp. 896-7, 124-136, pls. xi-xiv.
- 1899. Some Features of the Drift on Staten Island, N.Y.: Ann. New York Acad. Sci., vol. 12, pp. 91-102, pl. i.
- 1902. Fossil Forms from the Laramie Group of Colorado: Torrey, vol. 2, pp. 145-148, pls. iii, iv.
- 1903. A fossil Petal and a fossil Fruit from the Cretaceous (Dakota Group) of Kansas: Bull. Torrey Bot. Club, vol. 30, pp. 102-105.
- 1903 A. Geological and Botanical Notes: Cape Cod and Chappaquidick Island, Mass.: Bull. New York Bot. Garden, vol. 2, pp. 381-407, pls. xl, xli.
- *— 1904. A Recent Discovery of Amber and other Fossil Plant Remains at Kreischerville: Proc. Nat. Sci. Assoc. Staten Island, vol. 9, pp. 31, 32 [not seen].

- HOLLICK, A.—1905 a. The Occurrence and Origin of Amber in the Eastern United States: *Amer. Nat.*, vol. 39, pp. 137-145, pls. i-iii.
- 1905 b. A Recent Discovery of Amber on Staten Island: *Journ. New York Bot. Garden*, vol. 6, no. 63, pp. 45-48, text-figs. 14, 15.
- 1905 c. Additions to the Palaeobotany of the Cretaceous Formation of Long Island, no. II.: *Bull. New York Bot. Garden*, vol. 3, pp. 403-418, pls. lxx-lxxix.
- * — 1905 d. Forest Fires in the Cretaceous of Staten Island: *Proc. Nat. Sci. Assoc. Staten Island*, vol. 9, pp. 35-36 [not seen].
- 1906 A. The Cretaceous Flora of Southern New York and New England: *Mon. U.S. Geol. Surv.*, no. 50, pp. 1-219, pls. i-xl.
- 1906 B. See HOLLICK and JEFFREY.—1906.
- * — 1906 c. Insect Borings in Cretaceous Lignite from Kreischerville: *Proc. Nat. Sci. Assoc. Staten Island*, vol. 10, pp. 23, 24 [not seen].
- * — 1908. Chemical Analysis of Cretaceous Amber from Kreischerville: *Proc. Nat. Sci. Assoc. Staten Island*, vol. 12, p. 34.
- and JEFFREY, E. C.—1906. Affinities of certain Cretaceous Plant-Remains commonly referred to the genera *Dammara* and *Brachyphylum*: *Amer. Nat.*, vol. 40, pp. 189-204, pls. i-iv. f
- and JEFFREY, E. C.—1909. Studies of Cretaceous Coniferous Remains from Kreischerville, New York: *Mem. New York Bot. Garden*, vol. 3, pp. 1-76, pls. i-xxix.
- HOLM, T.—1890. Notes on the Leaves of *Liriodendron*: *Proc. U.S. Nat. Mus.*, vol. 13, pp. 15-35, pls. iv-ix.
- 1895. On the Validity of some Fossil Species of *Liriodendron*: *Bot. Gazette*, vol. 20, pp. 312-316, pl. xxiii.
- 1896. Remarks upon *Paleohillia*, a problematic Fossil Plant: *Bot. Gazette*, vol. 21, pp. 207-209, pl. xvii.
- HOLZAPFEL, E.—1884. Ueber einige wichtige Mollusken der Aachener Kreide: *Zeitschr. deutsch. geol. Ges.*, vol. 36, pp. 454-484.
- 1885. Ueber die Fauna des Aachener Sandes und seine Aequivalente: *Zeitschr. deutsch. geol. Ges.*, vol. 37, pp. 595-609.
- HOSIUS, A.—1870. Beiträge zur Geognosie Westfalens.—Die westfälischen Kreideformation vorkommenden Pflanzenreste: *Neues Jahrb. f. Min.*, pp. 381, 382.
- 1870 A. Ueber einige Dicotyledonen der westfälischen Kreideformation: *Palaeontogr.*, vol. 17, pp. 89-104, pls. xii-xvii.
- and MARCK, W. von DER.—1880. Die Flora der westfälischen Kreideformation: *Palaeontogr.*, vol. 26, pp. 126-241, pls. xxiv-xliv.
- 1885. Weitere Beiträge zur Kenntniss der fossilen Pflanzen und Fische aus der Kreide Westfalens: *Palaeontogr.*, vol. 31, pp. 225-232, pls. xix, xx.
- HOVELACQUE, M.—1889. Sur la nature végétale de l'*Aachenosaurus* (résumé): *Bull. Soc. belge Géol.*, vol. 3, pp. 505, 506.
- 1890. Sur la nature végétale de l'*Aachenosaurus multident*, G. Smets: *Bull. Soc. belge Géol., Mém.*, vol. 4, pp. 59-72, pl. iii, text-figs.

- HOVELACQUE, M., and KILIAN.—1900. Album de Microphotographies de Roches sédimentaires. Pp. 1-14, pls. i-lxix.
- HUTTON, W.—1831-1837. See LINDLEY and HUTTON.
- IDDINGS, J. P.—1899. See HAGUE, A., *et alii*.—1899.
- IVES, J. C.—1861. Report upon the Colorado River of the West, explored in 1857 and 1858 by Lieutenant Joseph C. Ives. United States Senate, 36th Congress. Ex. Doc. order of Sec. of War.
- JASCHE, C. F.—1858. Die Gebirgsformationen in der Grafschaft Wernigerode am Harz, nebst Bemerkungen über die Steinkohlen-Formation in der Grafschaft Hohenstein. Pp. 1-118, pls. i-iv. Wernigerode.
- JEFFREY, E. C.—1906 B. The Wound Reactions of *Brachyphyllum*: Ann. Bot., vol. 20, pp. 383-394, pls. xxvii, xxviii.
- 1906 c. See HOLLICK and JEFFREY.—1906.
- 1907. *Araucariopitys*, a new genus of Araucarians: Bot. Gazette, vol. 44, pp. 435-444, pls. xxviii-xxx.
- 1908. On the Structure of the Leaf in Cretaceous Pines: Ann. Bot., vol. 22, pp. 207-220, pls. xiii, xiv.
- 1909. See HOLLICK and JEFFREY.—1909.
- 1910 A. On the Affinities of the genus *Yezonia*: Ann. Bot., vol. 24, pp. 767-773, 1 pl.
- 1910 B. New *Prepinus* from Martha's Vineyard: Proc. Boston Soc. Nat. Hist., vol. 34, pp. 1-6, pls. i-iv.
- 1910 c. A New *Prepinus* from Martha's Vineyard: Proc. Boston Soc. Nat. Hist., vol. 34, pp. 333-338, pl. xxxiii.
- and CHRYSLER, M. A.—1906. On Cretaceous *Pityoxyla*: Bot. Gazette, vol. 42, pp. 1-15, pls. i, ii.
- JOHNSTON, R. M.—1896. Further Contributions to the History of the Fossil Flora of Tasmania: Proc. Roy. Soc. Tasmania for 1894-5, pp. 57-63, & 5 pls.
- JONES, T. R.—1862. Trails, Tracks, and Surface-markings: Geologist, vol. 5, pp. 128-139, pl. vii.
- JUKES-BROWNE, A. J., and HILL, W.—1900. The Cretaceous Rocks of Britain.—Vol. I. The Gault and Upper Greensand of England: Mem. Geol. Surv. United Kingdom, pp. 1-499.
- KERNER, F.—1896. Kreidepflanzen von Lesina: Jahrb. k.k. geol. Reichsanst., vol. 45, pp. 37-58, pls. i-v.
- KERSHAW, E. M.—1910. A Fossil Solenostelic Fern: Ann. Bot., vol. 24, pp. 684-691, pl. lviii, text-fig.
- 1910 A. See STOPES and KERSHAW.—1910.
- KIDSTON, R., and GWYNNE-VAUGHAN, D. T.—1908. On the Fossil Osmundaceæ: Trans. Roy. Soc. Edinb., vol. 45, pp. 759-780, pls. i-vi [issued separately, 1907].
- KNOWLTON, F. H.—1889. The Fossil Wood and Lignites of the Potomac Formation: Amer. Geologist, vol. 3, pp. 99-106.

- KNOWLTON, F. H.—1889 A. Fossil Wood and Lignite from the Potomac Formation: Bull. U.S. Geol. Surv., no. 56, pp. 11-72, pls. i-vii.
- 1889 B. Description of two New Species of Fossil Coniferous Wood from Iowa and Montana: Proc. U.S. Nat. Mus., vol. 11, pp. 5-8, pls. ii, iii.
- 1889 C. See LESQUEREUX, L.—1889.
- 1890. A Revision of the Genus *Araucarioxylon* of Kraus, with compiled Descriptions and partial Synonymy of the Species: Proc. U.S. Nat. Mus., vol. 12, pp. 601-617.
- 1892. See LESQUEREUX, L.—1892.
- 1893. See WEED and KNOWLTON.—1893.
- 1893 A. Description of a New Fossil Species of *Chara*: Bot. Gazette, vol. 18, pp. 141, 142, text-figs.
- 1893 C. Bread-fruit Trees in North America: Science, vol. 21, pp. 24-25.
- 1893 D. See WEED and KNOWLTON.—1893.
- 1895. Description of a New problematical Plant from the Lower Cretaceous of Arkansas: Bull. Torrey Bot. Club, vol. 22, pp. 387-390, text-figs.
- 1895 A, in R. T. HILL.—1895 A. Report upon a small Collection of Fossil Plants from Black Hills, near Belvidere, Kansas, collected by Prof. R. T. Hill, in August, 1894: Amer. Journ. Sci., ser. 3, vol. 50, pp. 212-214.
- 1896. The Fossil Plants of the Denver Basin: Mon. U.S. Geol. Surv., no. 27, pp. 446-473. See EMMONS *et alii*.—1896.
- 1896 A. American Amber-producing Tree: Science, n. s., vol. 3, pp. 582-584, text-figs. 1-4.
- 1897. See STANTON, T. W., and KNOWLTON, F. H.—1897.
- 1898. A Catalogue of the Cretaceous and Tertiary Plants of North America: Bull. U.S. Geol. Surv., no. 152, pp. 1-247.
- 1898 A. Description of *Pityoxylon Hollicki*, n. sp. See HOLLICK, A.—1898 A.
- 1899 B. Fossil Flora of the Yellowstone National Park: Mon. U.S. Geol. Surv., no. 32, pt. ii, pp. 651-791, pls. lxxvii-cxxi. See HAGUE, A., *et alii*.
- 1900 A. Flora of the Montana Formation: Bull. U.S. Geol. Surv., no. 163, pp. 1-77, pls. i-xix.
- 1900 B. Preliminary Report on a Collection of Fossil Plants from the vicinity of Winthrop, Methow Valley, Northern Cascade Mountains, Washington, made by Prof. I. C. Russell, September 4, 1898: 20th Ann. Rep. U.S. Geol. Surv., pt. 2, pp. 117-118. See RUSSELL, I. C.—1900.
- 1901. Reports on Fossil Plants from the Woodbine Formation: 21st Ann. Rep. U.S. Geol. Surv., pt. 7, pp. 314-318, pl. xxxix.
- 1905. Fossil Plants of the Judith River Beds: Bull. U.S. Geol. Surv., no. 257, pp. 129-155, pls. xiv-xix. See also STANTON and HATCHER.—1905.

- KNOWLTON, F. H.—1907. Description of a Collection of Kootanie Plants from the Great Falls Coal Field of Montana: Smithsonian Miscell. Coll., vol. 50, pp. 105–127, pls. xi–xiv.
- 1904. Description of New Fossil Liverwort from the Fort Union of Montana: Proc. U.S. Nat. Mus., vol. 35, pp. 157–159, pl. xxv.
- 1908 A. See DILLER.—1908.
- 1909. The Stratigraphic Relations and Palæontology of the "Hill Creek Beds," "Ceratops Beds," and equivalents, and their Reference to the Fort Union Formation: Proc. Washington Acad. Sci., vol. 11, pp. 179–238.
- 1910. Succession and Range of Mesozoic and Tertiary Floras: Journ. Geol., vol. 18, pp. 105–116.
- KRASSER, F.—1889. Ueber die fossilen Pflanzenreste der Kreideformation in Mähren: Verhandl. zool.-bot. Ges. Wien, vol. 39, pp. 31–34.
- 1896. Beiträge zur Kenntniss der fossilen Kreideflora von Kronstadt in Mähren: Beitr. Paläont. Geol. Österreich-Ungarns, vol. 10, pp. 113–152, pls. xi–xvii.
- 1906 A. Ueber die fossile Kreideflora von Grünbach in Niederösterreich: Anzeig. k. Akad. Wiss. Wien, vol. 43, pp. 41–43.
- and KUBART.—1906 B. Vorläufige Bericht ueber die Fossilen Flora von Moletuin in Mähren: Anzeig. k. Akad. Wiss. Wien, vol. 43, pp. 46–47.
- KREJČÍ, J.—1853. Kounická skála: Ziva, vol. 1, p. 28, pls. i, ii.
- KRENDOWSKIJ, M.—1880. Beschreibung fossiler Bäume hauptsächlich aus dem Süden Russlands.—I, II: Arbeiten Naturf.-Ges. k. Univ. Charkow, vol. 13, pp. 263–294, pls. i, ii. [In Russian, ref. by Winkler, Bot. Centralbl., vol. 6, 1881, pp. 415–417.]
- KUBART.—1906. See KRASSER and KUBART.—1906 B.
- KURTZ, F.—1902. Contribuciones á la Palæophytologia Argentina.—III: Sobre la existencia de una Dakota-Flora en la Patagonia Austro-Occidental: Revista Mus. La Plata, vol. 10, pp. 43–59.
- LAMPE, E.—1894. Ueber neue Fundorte der subhercynischen Flora: Zeitschr. f. gesammte Naturwiss., vol. 67, pp. 193–198.
- LAMPLUGH, G. W., and WALKER, J. F.—1903. On a Fossiliferous Band at the Top of the Lower Greensand near Leighton Buzzard (Bedfordshire): Quart. Journ. Geol. Soc., vol. 59, pp. 234–265, pls. xvi–xviii.
- LANGE, T.—1890. Beiträge zur Kenntniss der Flora des Aachener Sandes: Zeitschr. deutsch. geol. Ges., vol. 42, pp. 658–676, pls. xxxii–xxxiv.
- LAURENT, L.—1899. Thèse—Flore des calcaires de Célas: Ann. Mus. d'Hist. Nat. Marseille, ser. 2, Bull., vol. 1, fasc. 2, pp. 1–148, pls. i–xiv, & 3 maps.
- 1898. See MARION, A. F., and LAURENT, L.—1898.
- 1907. Les Progrès de la Paléobotanique Angiospermiqne dans la dernière décade: Progress. Rei Bot., vol. 1, pp. 319–367.
- LERICHE, M.—1909. Sur les Fossiles de la Craie phosphatée de la Picardie, à *Actinocamax quadratus*: Assoc. Franç. Avance. Sci., sess. 37, year 1908, pp. 494–503, text-figs.

- LESQUEREUX, L.—1859. On Fossil Plants collected by Dr. John Evans at Vancouver Island and at Bellingham Bay, Washington Territory; in a letter from L. Lesquereux to J. D. Dana, dated Columbus, Ohio, May 12, 1859: also in *Amer. Journ. Sci.*, ser. 2, vol. 27, pp. 360-362.
- 1860. Note on Prof. Newberry's criticisms of Prof. Heer's determination of Species of North American Fossil Plants: *Amer. Journ. Sci.*, ser. 2, vol. 29, pp. 434-436.
- 1868. On some Cretaceous Fossil Plants from Nebraska: *Amer. Journ. Sci.* ser. 2, vol. 46, pp. 91-105.
- 1868 A. In Hayden's Notes on the Lignite Deposits of the West: *Amer. Journ. Sci.*, ser. 2, vol. 45, pp. 205-208.
- 1869. On Species of Forest Plants from the Tertiary of the State of Mississippi [Appendix on Cretaceous Plants, pp. 430-433]: *Trans. Amer. Phil. Soc.*, vol. 13, pp. 411-433, pls. xiv-xxiii.
- 1871. On the Fossil Plants of the Cretaceous and Tertiary Formations of Kansas and Nebraska: Hayden's Prelim. Rep. U.S. Geol. Surv. Wyoming, pp. 370-385.
- 1872. Fossil Flora: Hayden's Prelim. Rep. U.S. Geol. Surv. Montana, pp. 283-304.
- 1873. Lignitic Formation and Fossil Flora: Hayden's 6th Ann. Rep. U.S. Geol. Surv., pp. 317-428.
- 1874. Contributions to the Fossil Flora of the Western Territories. —I. The Cretaceous Flora: Rep. U.S. Geol. Surv. Territ., vol. 6, pp. 1-133, pls. i-xxx.
- 1876 A. New Species of Fossil Plants from the Cretaceous Formation of the Dakota Group: Bull. U.S. Geol. Surv. Territ., ser. 2, no. 5, pp. 391-400.
- 1876 B. A Review of the Cretaceous Flora of North America: Ann. Rep. U.S. Geol. Surv. Territ., for 1874, pp. 316-365, pls. i-viii.
- 1876 C. On some New Species of Fossil Plants from the Lignitic Formations: Bull. U.S. Geol. Surv. Territ., ser. 2, no. 5, pp. 363-389; also Ann. Rep. for 1874, pp. 296-315.
- 1878 A. Remarks on Specimens of Cretaceous and Tertiary Plants secured by the Survey in 1877; with a List of the Species hitherto described: Tenth Ann. Rep. U.S. Geol. Surv. Territ., for 1876, pp. 481-520.
- 1878 B. Contributions to the Fossil Flora of the Western Territories.—II. The Tertiary Flora: Rep. U.S. Geol. Surv. Territ., vol. 7, pp. 1-366, pls. i-lxv.
- 1878 C. See NEWBERRY, J. S.—1878.
- 1882. Remarks on the Cretaceous and Tertiary Flora of the Western Territories: *Amer. Nat.*, vol. 16, pp. 102-108.
- 1883. Contributions to the Fossil Flora of the Western Territories. III. The Cretaceous and Tertiary Floras: Rep. U.S. Geol. Surv. Territ., vol. 8, pp. 1-283, pls. i-lx.
- 1884. Report on the Recent Additions of Fossil Plants: Bull. Mus. Comp. Zool. Harvard, vol. 7, pp. 225-230.

- LESQUEREUX, L.—1889. Recent Determinations of Fossil Plants from Kentucky, Louisiana, Oregon, California, Alaska, Greenland, etc., with descriptions of new Species (posthumous work edited by Knowlton): Proc. U.S. Nat. Mus., vol. 11, pp. 11-38, pls. iv-xvi.
- 1892. The Flora of the Dakota Group (edited by Knowlton): Mon. U.S. Geol. Surv., no. 17, pp. 1-256, pls. i-lxvi.
- 1893. The Genus *Winchellia*: Amer. Geologist, vol. 12, pp. 209-213, pls. viii, ix.
- 1895. Cretaceous Fossil Plants from Minnesota: Minnesota Geol. and Nat. Hist. Survey, Final Rep., 1885-1892, vol. 3, pt. i (Palaeontology), pp. 1-22, pls. A, B.
- LIEBUS, A.—1901. Ueber ein fossiles Holz aus der Sandablagerung Sulawa bei Radotin: Sitzb. deutsch. naturwiss.-mediz. Verein. Böhmen, Lotos, n. s., vol. 21, pp. 15-18.
- LIGNIER, O.—1907. Végétaux fossiles de Normandie.—IV. Bois Divers. (1^{re} série): Mém. Soc. Linn. Normandie, vol. 22, pp. 239-332, pls. xvii-xxiii.
- LIMA, W. DE.—1901. Noticia sobre alguns vegetaes fosseis da flora senoniana (sensu lato) do solo Portuguez: Comm. Serv. Geol. Portugal, vol. 4, pp. 1-12.
- LINDLEY, J., and HUTTON, W.—1831-37. The Fossil Flora of Great Britain; or, Figures and Descriptions of the Vegetable Remains found in a fossil state in this Country. Vol. 1, pp. 218, pls. lxxix; vol. 2, pp. 208, pls. lxxx-clvi; vol. 3, pp. 230, pls. clvii-ccxxx. London, 1831-37.
- LORENZ, J., von LIBURNAU.—1897. Eine fossile *Halimeda* aus dem Flysch von Muntige (Monticulus) bei Salzburg: Sitzb. k. Akad. Wiss. Wien, vol. 106, pp. 174-177, pls. i, ii.
- 1901. Zur Deutung der fossilen Fucoiden-Gattungen *Tenidium* und *Gyrophyllites*: Denkschr. k. Akad. Wiss. Wien, vol. 70, pp. 523-583, pls. i-iv, text-figs.
- 1902. Ergänzung zur Beschreibung der fossilen *Halimeda Fuggeri*: Sitzb. k. Akad. Wiss. Wien, vol. 111, pp. 685-712, pl. ii.
- LORENZ, T.—1902. Geologische Studien im Grenzgebiete zwischen helvetischer und ostalpiner Facies: Bericht. Naturforsch. Ges. Freiberg, vol. 12, pp. 1-61 (34-94), pls. i-ix.
- LUDWIG, R.—1861. Fossile Pflanzen aus der ältesten Abtheilung der Rheinisch-Wetterauer Tertiär-Formation: Palaeontogr., vol. 8, pp. 39-154, pls. vi-lx.
- LYDEKKER, R.—1889. See BOULENGER, G. A., and LYDEKKER, R.—1889.
- LYELL, C.—1834. See MURCHISON and LYELL.—1834.
- MAAS, G.—1895. Die untere Kreide des subhercynen Quadersandstein Gebirges: Zeitschr. deutsch. geol. Ges. vol. 47, pp. 227-302, pls. v-ix. [For plants in Neocomian and Gault, see pp. 275 and 282.]
- MACGIE, S. J.—1862 A. Some Fossil Fruits from the Chalk: Geologist, vol. 5, pp. 1-4, pl. i.

- MACKIE, S. J.—1862 B. The "Dragon-Tree" of the Kentish Rag: *Geologist*, vol. 5, pp. 401-404, pl. xxii.
- MAILLARD, G.—1887. Considérations sur les fossiles décrits comme algues: *Mém. Soc. paléont. Suisse*, vol. 24, pp. 2-40, pls. i-v.
- MALLOIZEL, G.—1887. Oswald Heer, Bibliographie et Tables Iconographiques. Précédé d'une Notice biographique par R. Zeiller. Pp. 176, and portrait. Stockholm.
- MANTELL, G. A.—1822. The Fossils of the South Downs, or Illustrations of the Geology of Sussex. Pp. 327, pls. xlii, & map. London.
- 1827. Illustrations of the Geology of Sussex: containing a general view of the geological relations of the South-Eastern Part of England; with Figures and Descriptions of the Fossils of the Tilgate Forest. Pp. 92, pls. xx, & map. London.
- 1833. The Geology of the South-East of England. Pp. 415, pls. v, & map. London.
- 1835. A Tabular Arrangement of the Organic Remains of the County of Sussex: *Trans. Geol. Soc.*, ser. 2, vol. 3, pp. 201-216 [Issued separately in 1829.]
- 1843. Description of some Fossil Fruits from the Chalk Formation of the South-East of England: *Proc. Geol. Soc.*, vol. 4, pp. 34, 35.
- 1844. The Medals of Creation; or first lessons in Geology and in the Study of Organic Remains. 2 vols. Pp. 1016, pls. vi. London.
- 1846. Description of some Fossil Fruits from the Chalk Formation of the South-East of England: *Quart. Journ. Geol. Soc.*, vol. 2, pp. 51-54, pl. ii.
- 1847. Geological Excursions round the Isle of Wight and along the adjacent Coast of Dorsetshire. Pp. 428, text-illustr. London.
- 1851. Petrifications and their Teachings; or, a Handbook to the Gallery of Organic Remains of the British Museum. Pp. 496, text-illustr. London.
- 1854. The Medals of Creation, ed. 2: 2 vols. Pp. 930. London.
- MARCK, W. VON DER.—1864. Fossile Fische, Krebse und Pflanzen aus dem Plattenkalk der jüngsten Kreide in Westphalen: *Palæontogr.*, vol. 11 [plants, pp. 77-83, pl. xiii].
- MARIK, V.—1901. Príspevek K florě českého Cenomanu: *Rospravy Ceske Akad. Cisare Praze*, vol. 10, pt. 3, pp. 1-16, pls. i, ii.
- MARION, A. F.—1873. See SAPORTA and MARION.—1873.
- 1878. See SAPORTA and MARION.—1878.
- 1885. See SAPORTA and MARION.—1885.
- 1890. Sur la Flore turonienne des Martigues (Bouches-du-Rhône): *Comptes Rendus Acad. Sci. Paris*, vol. 110, pp. 1052-1055.
- 1890 A. See VASSEUR, G.—1890.
- and LAURENT, L.—1898. Examen d'une Collection de Végétaux fossiles de Roumanie: *Anuar. Mus. Geol. Paleont. Bucharest* for 1895, pp. 187-227, pls. i, ii.
- MARTIN, K.—1890. Untersuchungen über den Bau von *Orbitolina* von Borneo: *Samml. geol. Reichsmus. Leiden*, ser. 1, vol. 4, pp. 211-231, pls. xxiv, xxv.

- MARTIN, K.—1901. *Lithothamnium* in cretaceischen und jüngeren Ablagerungen tropischer Inseln: Centralbl. f. Min., 1901, pp. 161-165.
- MARTIUS, C.—1822. De plantis nonnullis antediluvianis ope specierum inter tropicos viventium illustrandis: Denkschr. k. bot. Ges. Regensburg, vol. 2, pp. 121-147, pl. ii.
- MASSALONGO, A.—1855. Zoophycos, novum genus plantarum fossilium. Pp. 52, pls. iii. Verona.
- 1856. Studii Paleontologici. Pp. 53, pls. vii. Verona.
- 1857. Vorläufige Nachricht über die neueren paläontologischen Entdeckungen am Monte Bolca: Neues Jahrb. f. Min., pp. 775-778.
- McBRIDE, T. H.—1893. A New Cycad: Amer. Geologist, vol. 12, pp. 248-250, pl. xi.
- MEEK, F. B., and HAYDEN, F. V.—1859. Remarks on the Lower Cretaceous Beds of Kansas and Nebraska, together with Descriptions of some New Species of Carboniferous Fossils from the Valley of Kansas River [with letter by Heer]: Proc. Acad. Nat. Sci. Philadelphia, 1858, pp. 256-266.
- 1859 b. On the so-called Triassic Rocks of Kansas and Nebraska. Amer. Journ. Sci., ser. 2, vol. 27, pp. 31-35.
- 1859 c. Remarks on the Lower Cretaceous Beds of Kansas and Nebraska: Amer. Journ. Sci., ser. 2, vol. 27, pp. 219-227, text-figs.
- MENZEL, P.—1908. Fossile Koniferen aus der Kreide- und Braunkohlenformation Nordböhmens: Abhandl. naturwiss. Ges. Isis, Dresden, pp. 27-32, pl. ii.
- 1909. Fossile Pflanzenreste aus den Mungo-Schichten in Kamerun: Abhandl. k. preuss. geol. Landesanst., vol. 62, pp. 399-404, pl. ii.
- MERCKLIN, C. E. VON.—1852. Prospectus der paläontologischen Pflanzenüberreste in Russland, so wie ihrer Erforschung: Bull. Acad. Imp. Sci. St. Pétersbourg, vol. 10, pp. 373-384.
- 1853. Verzeichniss aller in Russland bis jetzt (November 1852) aufgefundenen, beschriebenen, unbeschriebenen oder zweifelhafter fossiler Pflanzen: Bull. Acad. Imp. Sci. St. Pétersbourg, vol. 11, pp. 300-305.
- 1855. Palæodendrologikon Rossicum—Vergleichende anatomisch-mikroskopische Untersuchungen fossiler Hölzer aus Russland: Preisschrift k. Akad. Wiss. S. Petersburg. Pp. 99, with Atlas, pls. xx.
- MERRILL, G. P.—1907. Catalogue of the Type and Figured Specimens of Fossils, Minerals, Rocks and Ores in the Department of Geology, United States National Museum: Bull. U.S. Nat. Mus., no. 53 [for plants see pp. 87-340].
- MESCHINELLI, A.—1892. Fungi fossiles: in P. A. Saccardo's Sylloge Fungorum omnium hucusque cognitorum, vol. 10, Suppl. univ., pp. 741-808. Patavii.
- 1895. Fungi fossiles: in Saccardo, *op. cit.*, vol. 11, pp. 657-659. Patavii.

- MESCHINELLI, A.—1902. *Fungorum fossilium omnium hucusque cognitorum Iconographia*. Pp. 144, pls. xxxi. Vicetiae.
- MILLER, S. A.—1881. *North American Mesozoic and Cænozoic Geology and Palæontology* [reprinted from *Proc. Cincinnati Soc. Nat. Hist.*]. Cincinnati.
- MIQUEL, F. A.—1853. *De fossiele Planten van het Krijt in het Hertogdom Limburg: Verhandl. geol. Kaart. Nederl.*, vol. 1, pp. 35-56, pls. i-vii.
- 1861. *Prodromus Systematis Cycadearum*. Pp. 36. Amsterdam.
- MORRIS, J.—1854. *A Catalogue of British Fossils*. Ed. 2. Pp. 372 [plants, pp. 1-26]. London.
- MORTON, S. G.—1832. *On the Analogy which exists between the Marl of New Jersey and the Chalk Formation of Europe: Amer. Journ. Sci.*, vol. 22, pp. 90-95.
- MOUGEOT, A.—1844. See SCHIMPER and MOUGEOT.—1844.
- MOURLON, M.—1881. *Géologie de la Belgique*. Vol. 2. Pp. 384. Paris and Bruxelles.
- MUNIER-CHALMAS, M.—1877. *Observations sur les Algues calcaires appartenant au groupe des Siphonées verticillées et confondues avec les Foraminifères: Comptes Rendus Acad. Sci. Paris*, vol. 85, pp. 814-817, text-figs.
- 1879. *Observations sur les Algues calcaires confondues avec les Foraminifères et appartenant au groupe des Siphonées dichotomes: Bull. Soc. géol. France*, ser. 3, vol. 7, pp. 661-670.
- MURCHISON, R. I.—1835. See SEDGWICK, A., and MURCHISON, R. I.—1835.
- NATHORST, A. G.—1874. *Om några fönnodade västfossilier: Öfvers. K. Svensk. Vet.-Akad. Förhandl.*, year 30, no. 9, pp. 25-52, pls. xv-xix.
- 1881 A. *Berättelse, afgifven till Kongl. Vetenskaps-Akademien, om en med understöd af allmänna medel utförd vetenskaplig resa till Schweiz och Tyskland: Öfvers. K. Svensk. Vet.-Akad. Förhandl.*, year 38, no. 1, pp. 61-84, pl. i.
- 1881 B. *Om Spår af Några Evertebrerade djur m. m. och deras Paleontologiska Betydelse.—French resumé—Mémoire sur quelques traces d'animaux sans vertèbres etc. et de leur portée paléontologique: K. Svensk. Vet.-Akad. Handl.*, vol. 18, pp. 1-104, pls. i-xi.
- 1883. *Contributions à la Flore fossile du Japon: K. Svensk. Vet.-Akad. Handl.*, vol. 20, no. 2, pp. 1-92, pls. i-xvi.
- 1885. *Palæontologiska Forskningar vid Wajgattet och Sofias Färd till Kap York: extract from Nordenskiöld's Den Andra Dicksonska Expeditionen till Grönland*, pp. 250-248, & map. Stockholm.
- 1886 A. *Nouvelles Observations sur des Traces d'Animaux et autres Phénomènes d'origine purement mécanique décrits comme Algues fossiles: K. Svensk. Vet.-Akad. Handl.*, vol. 21, no. 14, pp. 1-58, pls. i-v.

- NATHORST, A. G.—1886 B. Ueber die Benennung fossiler Dikotylenblätter : Bot. Centralbl., vol. 25, pp. 21-25; 52-55; 89-91.
- 1890. Beiträge zur mesozoischen Flora Japans: Denkschr. k. Akad. Wiss. Wien, vol. 57, pp. 43-60, pls. i-vi, & map.
- 1890 A. Ueber die Reste eines Brotfruchtbaums, *Artocarpus dicksoni*, n. sp., aus den cenomanen Kreideablagerungen Grönlands: K. Svensk. Vet.-Akad. Handl., vol. 24, no. 1, pp. 1-10, pl. i.
- 1891. Ueber das angebliche Vorkommen von Geschieben des Hörsandsteins in der norddeutschen Diluvialablagerungen: Archiv Ver. Freunde Natur, Mecklenburg, vol. 44, pp. 17-40, pl. i.
- 1893. Review of "Types of the Laramie Flora" by Lester F. Ward: Neues Jahrb. f. Min., vol. 2, pp. 219-222.
- 1893 A. See FELIX and NATHORST.—1893.
- 1894. Sveriges Geologi allmänföretligt framställt med en inledande historik om den geologiska forskningen i Sverige jemte en kort öfversigt af de geologiska systemen. Pp. 336, text illust. Stockholm.
- 1902. Beiträge zur Kenntniss einiger mesozoischen Cycadophyten: K. Svensk. Vet.-Akad. Handl., vol. 36, no. 4, pp. 1-28, pls. i-iii.
- 1906. Ueber *Thaumatopteris Schenki*, Nath.: K. Svensk. Vet.-Akad. Handl., vol. 40, no. 3, pp. 1-9, pls. i, ii.
- 1907. Paläobotanische Mittheilungen.—1. *Pseudocycas*, eine neue Cycadophytengattung aus den cenomanen Kreideablagerungen Grönlands: K. Svensk. Vet.-Akad. Handl., vol. 42, no. 5, pp. 1-20, pls. i-iii.
- 1908. Paläobotanische Mittheilungen.—4. Ueber die Untersuchungen kutinisirter fossiler Pflanzenteile.—5. Ueber *Nathorstia*, Heer: K. Svensk. Vet.-Akad. Handl., vol. 43, no. 1, pp. 1-19, pls. i-iii.
- NEUMANN, R.—1907. Beiträge zur Kenntniss der Kreideformation in Mittel-Peru: Neues Jahrb. f. Min., vol. 24, pp. 69-132, pls. i-v.
- NEWBERRY, J. S.—1859. Fossil Plants from the Cretaceous of Kansas and Nebraska; letter in MEEK & HAYDEN, 1859 B, p. 33.
- 1860. Notes on the Ancient Vegetation of North America: Amer. Journ. Sci., ser. 2, vol. 29, pp. 208-218.
- 1861. [Cretaceous Plants in Ives' Report on the Colorado River, pp. 129-132, pl. iii.] See IVES, J. C.—1861.
- 1863. Descriptions of the Fossil Plants collected by Mr. George Gibbs, Geologist to the United States North-West Boundary Commission, under M. Archibald Campbell, United States Commissioner: Boston Journ. Nat. Hist., vol. 7, pp. 506-524.
- 1868. Notes on the Later Extinct Floras of North America: Amer. Journ. Sci., ser. 2, vol. 46, pp. 401-407.

- NEWBERRY, J. S.—1869. [Report on the Cretaceous and Tertiary Plants in Reynolds' Exploration of the Yellowstone and Missouri Rivers, pp. 145-174.] See HAYDEN, F. V.—1869.
- 1870. Notes on the Later Extinct Floras of North America, with Descriptions of some New Species of Fossil Plants from the Cretaceous and Tertiary Strata: Ann. Lyceum Nat. Hist. New York, vol. 9, pp. 1-76.
- 1874. On the Lignites and Plant-Beds of Western America: Amer. Journ. Sci., ser. 3, vol. 7, pp. 399-404.
- 1878. Illustrations of Cretaceous and Tertiary Plants of the Western Territories of the United States [names by Lesquereux]: U.S. Geol. Surv. Territ., pls. i-xxvi [used by NEWBERRY.—1898].
- 1881. American Cretaceous Flora: Nature, vol. 24, pp. 191-192.
- 1883. Brief Descriptions of Fossil Plants, chiefly Tertiary, from Western North America: Proc. U.S. Nat. Mus. vol. 5, pp. 502-514.
- 1886 a. The Flora of the Amboy Clays [abstract]: Bull. Torrey Bot. Club, vol. 13, pp. 33-37.
- 1886 b. *Bauhinia cretacea*, n. sp.: Bull. Torrey Nat. Club, vol. 13, pp. 77-78, pl. lvi.
- 1886 c. The Cretaceous Flora of North America: Trans. New York Acad. Sci., vol. 5, pp. 133-137.
- 1887. The Ancestors of the Tulip-Tree: Bull. Torrey Bot. Club, vol. 14, pp. 1-7, pls. lxi, lxii.
- 1889. Notes on Flora of Puget Sound Strata: in C. A. White, Bull. U.S. Geol. Surv., no. 51, p. 51.
- 1890. The Laramie Group—its Geological Relations, its Economic Importance, and its Fauna and Flora: Trans. New York Acad. Sci., vol. 9, pp. 27-32.
- 1891. The Flora of the Great Falls Coal-Field, Montana. Amer. Journ. Sci., ser. 3, vol. 41, pp. 191-201, pl. xiv.
- 1895. The Flora of the Amboy Clays [posthumous work edited by A. Hollick]: Mon. U.S. Geol. Surv., no. 26, pp. 1-137, pls. i-lviii.
- 1898. The Later Extinct Floras of North America [posthumous work edited by A. Hollick]: Mon. U.S. Geol. Surv., no. 35, pp. 1-151, pls. i-lxviii.
- NEWBOLD, T. J.—1848. On the Geological Position of the Silicified Wood of the Egyptian and Libyan Deserts, with a Description of the Petrified Forest near Cairo: Quart. Journ. Geol. Soc., vol. 4, pp. 349-357.
- NEWTON, R. B.—1909. On some Fossils of the Nubian Sandstone Series of Egypt: Geol. Mag., dec. 5, vol. 6, pp. 352-359, 388-397, pls. xix-xxi [see pp. 335-359].
- NILSSON, S.—1824. Underrättelse om fossila landväxter som finnas tillsammans med hafsmusslor, snäckor m. m. i den Skånska Grönlandskalken: K. Svensk. Vet.-Akad. Handl., 1824, pp. 143-148, pl. ii.
- 1832. Fossila Växter funna i Skåne och beskrifne: K. Svensk. Vet.-Akad. Handl. 1831, pp. 340-351, pls. i, ii.

- NORDENSKIÖLD, A. E.—1872. Account of an Expedition to Greenland in the Year 1870: Geol. Mag., vol. 9, pp. 289-306, pl. vii; pp. 355-368, pl. viii; pp. 409-427; pp. 449-463; pp. 516-524.
- NOVAK, J.—1907. Kopalna flora senónska z Potylicza (Die fossile Senonflora von Potylicz): Bull. Intern. Acad. Sci. Cracovie, pp. 45-56, pls. i, ii.
- OOSTER, W. A.—1869. Die organischen Reste der Zoophycos-Schichten der Schweizer-Alpen: Mittheil. Berlin. Mus. Naturgeschichte, vol. 1, pt. ii, pp. 15-35, pls. iii-xi, & one unnumb. pl.
- 1871. Beitrag zur Kenntniss der Fauna der obersten Kreideschichten am Nord-Ufer des Thuner-See's (Berner-Alpen): Protozoe Helv., vol. 2, pp. 43-72, pls. ix-xi.
- ORBIGNY, A. D'.—1852. Cours élémentaire de Paléontologie et de Géologie stratigraphiques, vol. 2, pp. 260-347, text-figs. Paris.
- OSWALD, F.—1906. A Treatise on the Geology of Armenia. Pp. 516. Beeston.
- OTTO, E. von.—1852-4. Additamente zur Flora des Quadersteingebirges in der Gegend um Dresden und Dippoldiswalde, enthaltend meist noch nicht oder wenig bekannte fossile Pflanzen. Vol. 1 (1852), pp. 27, pls. vii. Vol. 2 (1854), pp. x, 53, pls. ix. Dippoldiswalde.
- PELOURDE, F., 1908.—Recherches sur la position systématique des plantes fossiles dont les tiges ont été appelées *Psaronius*, *Psaroniocalen*, *Caulopteris*: Bull. Soc. bot. France, vol. 8, pp. 88-119, pls. iii, iv.
- PENHALLOW, D. P.—1900. Notes on the North American Species of *Dadoxylon*, with Special Reference to Type material in the Collections of the Peter Redpath Museum, McGill University: Proc. & Trans. Roy. Soc. Canada, ser. 2, vol. 6, pp. 51-79, pls. lviii-xcvii.
- 1902 A. *Osmundites skidegatensis*, n. sp.: Proc. & Trans. Roy. Soc. Canada, ser. 2, vol. 8, pp. 1-18, pls. i-vi.
- 1902 B. Notes on Cretaceous and Tertiary Plants of Canada: Proc. & Trans. Roy. Soc. Canada, ser. 2, vol. 8, pp. 31-72, pls. vii-xvi.
- 1907. A Report on Fossil Plants from the International Boundary Survey for 1903-05, collected by Dr. R. A. Daly: Trans. Roy. Soc. Canada, ser. 3, vol. 1, sect. 4, pp. 287-351, pls. i-ix, text-figs.
- 1907 A. A Manual of the North American Gymnosperms, exclusive of the Cycadales but together with certain exotic species. Pp. 374, pls. lv. Boston.
- 1908. Report on a Collection of Fossil Woods from the Cretaceous of Alberta: Ottawa Naturalist, vol. 22, pp. 82-85.
- 1908 A. Report on the Fossil Plants of the International Boundary Survey: Trans. Roy. Soc. Canada, ser. 3, vol. 2, sect. 4, pp. 287-352, pls. i-ix.
- PERUZZI, D. G.—1881. Osservazioni sui generi *Paleodictyon* e *Paleomeandron* dei terreni cretacei ed eocenici dell' Appennino sett. e centrale: Atti Soc. Tosc. Sci. Nat., vol. 5, pp. 3-8, pl. i.

- PHILLIPS, J.—1864. Oxford Fossils: Geol. Mag., vol. 2, pp. 292, 293, pl. ix.
- PINTO, A. C. O.—1908. Un remarquable cas de silicification: Bull. Soc. Portug. sci. nat., vol. 1, pp. 174–176, pl. viii.
- PIRSSON, L. V.—1898. See WEED and PIRSSON.—1898.
- POGGIO, E. DI.—1906. Le Piante fossili. Elementi di Paleofitologia. Pp. 200, text-figs. Turin.
- POLLARD, C. L.—1894. Note on Fossil Leaves from Great Neck, Long Island, New York: Trans. New York Acad. Sci., vol. 13, pp. 180, 181.
- POMEL, A.—1849. Ueber die fossile Flora des französischen Jura und der französischen Kreide: Amtl. Bericht deutsch. naturf. Versamml. 25 Aachen, pp. 331–354.
- POTONÉ, H.—1897.—Bennettitaceæ: in Engler & Prantl, Die natürlichen Pflanzenfamilien, Nachträge, pp. ii–iv, pp. 14–17. Leipzig.
- 1899. Lehrbuch der Pflanzenpaläontologie mit besonderer Rücksicht auf die Bedürfnisse des Geologen. Pp. 402, pls. iii, 355 text-figs. Berlin.
- 1900. See BORNHARDT, W.—1900.
- 1902. Fossile Hölzer aus der oberen Kreide Deutsch-Ostafrikas: Mittheil. deutsch. Schutzgebiet., vol. 15, pp. 227–229, pl. ii.
- 1903–1910. Abbildungen und Beschreibungen fossiler Pflanzenreste der paläozoischen und mesozoischen Formationen. [Published by K. preuss. geol. Landesanst., Berlin.]
- PUSCH, G. G.—1837.—Polens Paläontologie. Pp. 218, pls. xvi [records Cret. Algæ, pp. 3–4]. Stuttgart.
- RACIBORSKI, M.—1893. *Cycadeoidea Niedzwiedz-kii*, nov. sp.: Rozprawy Akad. Umiejętności, ser. 2, vol. 6, pp. 301–310, pls. vii, viii [resumé in Bull. Acad. Intern. Sci. Cracovie, 1892, pp. 355–359].
- 1909. *Rhizodendron* w opoce lwowskiej (*Rhizodendron* in den senonen Mergeln der Umgebung von Lemberg): Kosmos, vol. 34, pp. 845–848, text-fig.
- RAUFF, H.—1895. Ueber *Porocystis pruniformis*, Cragin (= ? *Araucarites Wardi*, Hill) aus der unteren Kreide in Texas: Neues Jahrb. f. Min., vol. 1, pp. 1–15, pl. i.
- RAYNOLDS, W. F.—1869. See HAYDEN, F. V.—1869.
- REID, C.—1902. Palaeobotany—Tertiary: Encyclop. Brit., ed. 10, vol. 31, pp. 432–440.
- REINSCH, P. F.—1905. Die Palinosphären, ein mikroskopischer vegetabler Organismus in der Mukronatenkreide: Centralbl. f. Min., no. 13, pp. 402–407.
- REIS, O. M.—1909. Zur Fucoidenfrage: Jahrb. k.k. geol. Reichsanst., vol. 59, pp. 615–638, pl. xvii.
- REISS, K.—1907. Untersuchung über fossile Hölzer aus Japan: Inaug.-Diss. Univ. Leipzig. Pp. vi, 224, pl. i.
- RENAULT, B.—1883. Cours de Botanique fossile fait au Muséum d'Histoire naturelle: Fougères. Pp. 241, pls. A, xxxv. Paris.
- 1888. Les Plantes fossiles. Pp. 309, text-figs. Paris.

- RENGER, K.—1866. *Předvěká rostlinstvo křídového útvaru českého* (Die vorweltliche Flora der böhmischen Kreideformation): Ziva, pp. 113–141, pl. i.
- 1866 a. *Stromovité kapradiny v křídovém útvaru českém* (Die Baumfarne in der Kreideformation Böhmens): Ziva, pp. 36–45.
- REUSS, A. E.—1844. *Geognostische Skizzen aus Böhmen*. Vols. 1 & 2. Prague.
- 1846. *Die Versteinerungen der böhmischen Kreideformation*. Pt. i, pp. 58, pls i–xiii; pt. ii, pp. 140, pls. xiv–li. [Cret. plants by Corda, pt. 2, pp. 81–96, pls. xlv–li.] Stuttgart.
- 1854. *Beiträge zur geognostischen Kenntniss Mährens*: Jahrb. k.k. geol. Reichsanst., 1854, pp. 659–765. [Cret. plants, p. 740.]
- RICHARDSON, J.—1873. *Report on the Coal-fields of Vancouver and Queen Charlotte Island, with a map of the distribution of the former*: Geol. Surv. Canada, Rep. Progress for 1872–73, pp. 32–86.
- RICHTER, P. B.—1899 a. *Neocompflanzen der Kelb'schen Sandgrube bei Quedlinburg*: Zeitschr. deutsch. geol. Ges., vol. 51, pp. 39–41.
- 1899 b. *Ueber Quedlinburger Kreide-Coniferen, insbesondere über solche, welche an Geinitzien und Sequoien erinnern*: Zeitschr. deutsch. geol. Ges., vol. 51, pp. 43–44.
- 1901. *Pflanzen aus dem Neocom des Langenberges bei Quedlinburg*: Zeitschr. deutsch. geol. Ges., vol. 53, pp. 20–21.
- 1904. *Ueber die Kreidepflanzen der Umgebung Quedlinburgs*: Beitr. Programm k. Gymn. Quedlinburg, pp. 40, pls. ii.
- 1905. *Beiträge zur Flora der oberen Kreide Quedlinburgs und seiner Umgebung*.—I. Die Gattung *Credneria* und einige seltenere Pflanzenreste. Pp. 18, pls. vi. Leipzig.
- 1906. *Beiträge zur Flora der unteren Kreide Quedlinburgs*.—I. Die Gattung *Hausmannia*, Dunker, und einige seltenere Pflanzenarte. Pp. 27, pls. vii. Leipzig.
- 1909. *Beiträge zur Flora der unteren Kreide Quedlinburgs*.—II. Die Gattung *Nathorstiana*, P. Richter, und *Cylindrites spongioides*, Goeppert. Pp. 12, pls. viii–xiii. Leipzig.
- 1910. *Ueber Nathorstiana P. Richter und Cylindrites spongioides Goepp.*: Monatsb. deutsch. geol. Ges., vol. 62, pp. 278–284.
- RODR, E.—1878. [Ueber einige Stämme aus der böhmischen Kreideformation]: *Vesmir* (Prague), vol. 7, pp. 183–185, 194–196.
- ROEMER, F. A.—1839. *Die Versteinerungen des norddeutschen Oolithen-Gebirges*: ein Nachtrag. Pp. iv, 59, pls. A, xvii–xx. Hanover.
- 1841. *Die Versteinerungen des norddeutschen Kreidegebirges*. Pp. 145, pls. xvi. Hanover.
- ROEMER, F.—1852. *Die Kreidebildungen im Texas und ihre organischen Einschlüsse*. Pp. 109, pls. xi [notes on Cret. plants by Unger, pp. 94, 95]. Bonn.
- 1854 c. [Letter to Prof. Bronn on Cretaceous of Aix]: *Neues Jahrb. f. Min.*, pp. 167–169.

- ROEMER, F.—1855. Bemerkungen über die Kreidebildungen der Gegend von Aachen, gegründet auf Beobachtungen im Jahre 1853: *Zeitschr. deutsch. geol. Ges.*, vol. 7, pp. 534-546.
- 1856. See BRONN, H. G., and ROEMER, F.—1856.
- 1870. Geologie von Oberschlesien. Pp. 587, pls. i-l. [*Cret.* pp. 276-357.] Breslau.
- 1887. Ueber die Auffindung einer fossilen Flora in Thonen der Kreide-Formation bei Bunzlau: 64 *Jahresber. Schles. Ges. Vaterl. Cult.*, p. 117.
- 1889. Ueber Blattabdrücke in senonen Thonschichten bei Bunzlau in Niederschlesien: *Zeitschr. deutsch. geol. Ges.*, vol. 41, pp. 139-147, pl. xii.
- ROTHPLETZ, A.—1891 A. Das Verhältniss der fossilen zu den lebenden *Lithothamnium*-Arten: *Bot. Centralbl.*, vol. 45, pp. 235-36.
- 1891 B. Fossile Kalkalgen aus den Familien der Codiaceen und der Corallineen: *Zeitschr. deutsch. geol. Ges.*, vol. 43, pp. 295-322, pls. xv-xvii.
- 1896. Ueber die Flysch-Fucoiden und einige andere fossile Algen sowie über liasische Diatomeen-führende Hornschwämme: *Zeitschr. deutsch. geol. Ges.*, vol. 48, pp. 854-914, pls. xxii-xxiv.
- RUEDEMANN, R.—1909. Some Marine Algæ from the Trenton Limestone of New York: *Bull. New York State Mus.*, no. 133, pp. 194-216, pls. 1-3.
- RUSSELL, I. C.—1900. A Preliminary Paper on the Geology of the Cascade Mountains in Northern Washington: 20th *Ann. Rep. U.S. Geol. Surv.*, pp. 89-210, pls. viii-xx.
- SACCARDO, P. A.—1892. See MESCHINELLI, A.—1892.
- SACCO, F.—1888. Note di Paleoenologia Italiana.—II: *Atti Soc. Ital. Sci. Nat.*, vol. 31, pp. 151-192, pls. i, ii.
- SAPORTA, G. DE.—1867. Note sur une collection de plantes fossiles provenant de la Craie à *Belemnites mucronatus* de Haldem en Westphalie: *Bull. Soc. géol. France*, ser. 2, vol. 24, pp. 33-36.
- 1868. Prodrome d'une Flore Fossile des Travertins anciens de Sézanne: *Mém. Soc. géol. France*, ser. 2, vol. 8, pp. 289-436, pls. i-xv.
- 1873-75. *Paléontologie Française, ou Description des Fossiles de la France.*—2nd sér. *Végétaux. Plantes Jurassiques.* Vol. 1, pp. 501, pls. lxx; vol. 2, pp. 339, pls. lviii. Paris.
- 1879. *Le Monde des Plantes avant l'Apparition de l'Homme.* Pp. 416, pls. xiii, & text-figs. Paris.
- 1880. Notice sur les Végétaux fossiles de la Craie inférieure des Environs du Havre: *Exposit. géol. 1877 Résumés, etc.*, sur la Géol. Normandie, forming vol. 6 of the *Bull. Soc. Géol. Normandie*, pp. 640-661, pls. i-iv.
- 1882. Sur quelques Types de Végétaux récemment observés à l'état fossile: *Comptes Rendus Acad. Sci. Paris*, vol. 94, pp. 922-924; 1020-1022.

- SAROTA, G. DE.—1882A. A propos des Algues fossiles. Pp. 82, pls. ix. Paris.
- 1884. Les Organismes problématiques des anciennes Mers. Pp. 102, pls. xii. Paris.
- 1888. Sur les Dicotylées prototypiques du Système infracrétacé du Portugal: Comptes Rendus Acad. Sci. Paris, vol. 106, pp. 1500-1504.
- 1890. Le *Nelumbium Provinciale* des Lignites Crétacés de Fuveau en Provence: Mém. Soc. géol. France, Paléont., no. 5, pp. 1-10, pls. xii-xiv.
- 1890A. Sur de nouvelles Flores fossiles, observées en Portugal, et marquant le Passage entre les Systèmes jurassique et infracrétacé: Comptes Rendus Acad. Sci. Paris, vol. 111, pp. 812-815.
- 1890B. Revue des Travaux de Paléontologie Végétale parus en 1888 ou dans le cours des Années précédentes: Rev. gén. Bot., vol. 12, pp. 176-184.
- 1891. Sur les plus anciennes Dicotylées européennes observées dans le Gisement de Cercal, en Portugal: Comptes Rendus Acad. Sci. Paris, vol. 113, pp. 249-253.
- 1891A. Paléontologie Française, ou Description des Fossiles de la France continuée par une réunion des Paléontologistes sous la direction d'un comité spécial.—Vol. 4. Plantes Jurassiques. Pp. 547, pls. lxxiv. Paris.
- 1893. Revue des Travaux de Paléontologie Végétale parus en France dans le cours des Années 1889-1892: Rev. gén. Bot., vol. 15, pp. 1-54, pls. iii-vi.
- 1894. Flore fossile du Portugal, nouvelles contributions à la Flore mésozoïque: Direct. Trav. géol. Portugal. Pp. 288, pls. xxxix.
- 1894A. Nouveaux détails concernant les Nymphéinées. Nymphéinées infracrétacées: Comptes Rendus Acad. Sci. Paris, vol. 119, pp. 835-837.
- 1894B. Étude monographique sur les *Rhizocaulon*: Rev. gén. Bot., vol. 6, pp. 241, 259; 301-309; 324-344; pls. v-xii.
- and MARION, A. F.—1873. Essai sur l'État de la Végétation de l'Époque des Marnes Heersiennes de Gelinden: Mém. Couron. Acad. roy. Belg., vol. 37, no. 6, pp. 1-94, pls. i-xii.
- — 1874. Sur les couches supérieures à la Mollasse du bassin de Théziers (Gard) et les plantes fossiles de Vaquières: Bull. Soc. géol. France, vol. 12, pp. 272-287, pls. vii-viii.
- — 1873. Révision de la Flore Heersienne de Gelinden d'après une collection appartenant au Comte G. de Looz: Mém. Couron. Acad. roy. Belg., vol. 41, no. 3, pp. 1-112, pls. i-xiv.
- — 1881. L'Évolution du Règne Végétal.—Les Cryptogames. Pp. xii, 238, text-figs. 85. Paris.
- — 1885. L'Évolution du Règne Végétal.—Les Phanerogames. Vol. 1. Pp. 249, 136 text-figs. Vol. 2. Pp. 247. Paris.
- SCHENK, A.—1871A. Beiträge zur Flora der Vorwelt.—III. Die fossilen Pflanzen der Wernsdorfer Schichten in den Nordkarpathen: Palaontogr., vol. 19 (1869), pp. 1-34, pls. i-vii.

- SCHENK, A.—1871 B. Beiträge zur Flora der Vorwelt.—IV. Die Flora der nordwestdeutschen Wealdenformation: Paläontogr., vol. 19, pp. 203-266, pls. xxii-xliii.
- 1876. Beiträge zur Flora der Vorwelt.—VI. Ueber einige Pflanzen-reste aus der Gosaufornation Nordtirols: Paläontogr., vol. 23, pp. 164-171, pls. xxviii-xxix.
- 1880 A. Ueber fossile Hölzer aus der Libyschen Wüste: Bot. Zeit., vol. 38, pp. 657-661.
- 1883. Fossile Hoelzer: Paläontogr., vol. 30, pt. 2, pp. 1-17, pls. i-v.
- 1890. Die fossilen Pflanzenreste: in the author's Handbuch der Botanik, vol. 4. Pp. 781. Breslau.
- 1890 A. See SCHIMPER and SCHENK.—1890.
- SCHUCHER, J. J.—1709-23. Herbarium diluvianum collectum. Tiguri, 1709, pp. 1-44, pls. i-x. Also Ed. nov. Lugduni Batavorum, 1723, pp. 1-119, pls. i-xiv. Leiden, Holland.
- SCHIMPER, W. P.—1869-74. Traité de Paléontologie végétale ou la Flore du monde primitif dans ses rapports avec les formations géologiques et la Flore du monde actuel. Vol. 1 (1869), pp. 738; vol. 2 (1870-72), pp. 968; vol. 3 (1874), pp. 896, pls. cx. Paris.
- and SCHENK, A.—1890. Handbuch der Paläontologie herausgegeben von Karl A. Zittel.—II. Paläophytologie. Pp. 958, text-illustr. Munich & Leipzig, 1879-1890.
- SCHLOTHEIM, E. F. VON.—1820. Die Petrefactenkunde auf ihrem jetzigen Standpunkte durch die Beschreibung seiner Sammlung versteinerter und fossiler Ueberreste des Thier- und Pflanzenreichs der Vorwelt erläutert. Pp. 437, pl. xv. Ed. 1832 with xxxvii pls. Gotha.
- 1822. Nachträge zur Petrefactenkunde. Pp. 100, pls. xxi. Gotha.
- SCHUCHERT, C.—1898. See WHITE, D., and SCHUCHERT, C.—1898.
- 1910. Palæogeography of North America: Bull. Geol. Soc. America, vol. 20, pp. 427-606, pls. xlii-ci [see pp. 587-597].
- SCHULZE, E.—1888. Ueber die Flora der subhercynen Kreide: Inaug.-Diss. Halle-Wittenberg. Pp. 33. [Reprinted in Zeitschr. f. gesammte Naturwiss., vol. 60, pp. 440-470.] Halle.
- SCHUSTER, J.—1910. Ueber Nicolien und Nicolien-ähnliche Hölzer: K. Svensk. Vet.-Akad. Handl., vol. 45, no. 6, pp. 1-18, pls. i-iii.
- SCOTT, D. H.—1907. The Flowering Plants of the Mesozoic Age, in the Light of Recent Discoveries: Journ. R. Microsc. Soc., pp. 129-141, pls. vi-ix.
- SEDGWICK, A., and MURCHISON, R. I.—1835. A Sketch of the Structure of the Eastern Alps; with Sections through the Newer Formations on the Northern Flanks of the Chain, and through the Tertiary Deposits of Styria, etc., etc.: Trans. Geol. Soc., ser. 2, vol. 3, pp. 301-420.
- SEWARD, A. C.—1894 A. Catalogue of the Mesozoic Plants in the Department of Geology, British Museum. The Wealden Flora.—Pt. I. Thallophyta—Pteridophyta. Pp. 1-179, pls. xi. London.

- SEWARD, A. C.—1894 B. *Algae as Rock-Building Organisms*: Sci. Progress, vol. 2, pp. 10-26.
- 1895. *Catalogue of the Mesozoic Plants in the Department of Geology, British Museum. The Wealden Flora.—Pt. II. Gymnospermæ*. Pp. 259, pls. xx. London.
- 1896 A. *Notes on the Geological History of Monocotyledons*: Proc. Phil. Soc. Cambridge, vol. 9, pp. 110-111.
- 1896 B. *Notes on the Geological History of Monocotyledons*. Ann. Bot., vol. 10, pp. 205-220, pl. xiv.
- 1898. *Fossil Plants, for Students of Botany and Geology*. Vol. 1, pp. 452, text-figs. Cambridge.
- 1899. *On the Structure and Affinities of Matonia pectinata*, R. Br., with Notes on the Geological History of the Matouineæ. Phil. Trans. Roy. Soc. London, vol. 191 B, pp. 171-209, pls. xvii-xx.
- 1900. *La Flore Wealdienne de Bernissart*: Mém. Mus. Roy. d'Hist. Nat. Belgique, vol. 1, pp. 1-37, pls. i-iv, text-figs.
- 1903. *Fossil Floras of Cape Colony*: Ann. S. African Mus., vol. 4, pp. 1-122, pls. i-xiv.
- 1904. *Floras of the Past; their Composition and Distribution*: Rep. Brit. Assoc. (Southport, 1903), pp. 824-849.
- 1907. *Fossil Plants from Egypt*: Geol. Mag., dec. 5, vol. 4, pp. 253-257, text-figs.
- 1910. *Fossil Plants, a text-book for Students of Botany and Geology*. Vol. 2, pp. 624, text-figs. Cambridge.
- & FORD, S. O.—1905. *The Anatomy of Todea*, with Notes on the Geological History and Affinities of the Osmundaceæ: Trans. Linn. Soc., Botany, ser. 2, vol. 6, pp. 237-260, pls. xxvii-xxx.
- 1906. *The Araucariæ, Recent and Extinct*: Phil. Trans. Roy. Soc. London, vol. 198 B, pp. 305-411, pls. 23, 24.
- & GOWAN, J.—1900. *The Maidenhair Tree (Ginkgo biloba, L.)*: Ann. Bot., vol. 14, pp. 109-154, pls. viii-x.
- SHIRLEY, J.—1898. *Additions to the Fossil Flora of Queensland*: Bull. Geol. Surv. Queensland, no. 7, pp. 1-25, pls. i-xxvii.
- 1902. *Notes on Fossil Plants from Duaringa, Ipswich, Dawson River, and Stanwell*: Bull. Geol. Surv. Queensland, no. 18, pp. 1-13, pls. i-viii.
- SINNOTT, G. W.—1909. *Paracedroxylon*, a New Type of Araucarian Wood: Rhodora, vol. 11, pp. 165-173, pls. lxxx-lxxxi.
- *SMETS, G.—1887. *Un Reptile nouveau des Sables d'Aix-la-Chapelle*. Pp. 133. Louvain.
- *— 1888. *Achenosaurus multidentis*, Reptile fossile des Sables d'Aix-la-Chapelle: Mém. Congrès sav. Cathol. Paris, pp. 23, pl. i.
- SMITH, E. A.—1894. *Report on the Geology of the Coastal Plain of Alabama (Cretaceous, Tertiary, and Post-Tertiary Formations)*: Geol. Surv. Alabama, pp. 1-759, pls. i-xxix [list of fossil plants, p. 348].
- SOLMS-LAUBACH, H. GRAF ZU.—1887. *Einleitung in die Paläophytologie vom botanischen Standpunkt aus*. Pp. 416. Leipzig. [English translation published by Oxford Univ. Press, 1891.]

- SOLMS-LAUBACH, H. GRAF ZU.—1890. Ueber die Fructification von *Bennettites Gibsonianus*, Carr.: Bot. Zeit., vol. 48, pp. 789-798; 805-816; 821-833; 843-847; pls. ix, x [translated Ann. Bot., vol. 5, pp. 419-452, pls. xxv, xxvj].
- 1892. See CAPELLINI, G., and SOLMS-LAUBACH.—1892.
- 1901. Monograph of the Acetabulariæ: Trans. Linn. Soc., Botany, ser. 2, vol. 5, pp. 1-39, pls. i-iv. [1895.]
- SQUINABOL, S.—1890. Alghe e Pseudoalghe fossili italiane: Atti Soc. Ligust. Sci. Nat., vol. 1, pp. 29-49; 166-199, pls. v-xii.
- STACHE, G.—1880. Die Liburnische Stufe: Verhandl. k.k. geol. Reichsanst., 1880, pp. 194-209.
- 1889. Die Liburnische Stufe und deren Grenz-Horizonte. Eine Studie über die Schichtenfolgen der Cretacisch-Eocänen oder Protocänen Landbildungsperiode im Berichte Küstenländer von Österreich-Ungarn: Abhandl. k.k. geol. Reichsanst., vol. 13, pp. 1-170, pls. i-vi, & map.
- STANTON, T. W.—1894. See DILLER and STANTON.—1894.
- 1895. Contributions to the Cretaceous Palæontology of the Pacific Coast.—The Fauna of the Knoxville Beds: Bull. U.S. Geol. Surv., no. 133, pp. 1-85, pls. i-xx [notes on plants of Horsetown age by Fontaine, p. 22].
- 1899. See HAGUE, A., et alii.—1899.
- 1909. The Age and Stratigraphic Relations of the "Ceratops Beds" of Wyoming and Montana: Proc. Washington Acad. Sci., vol. 11, pp. 239-293.
- and KNOWLTON, F. H.—1897. Stratigraphy and Palæontology of the Laramie and related Formations in Wyoming: Bull. Geol. Soc. America, vol. 8, pp. 127-156.
- and HATCHER, J. B.—1905. Geology and Palæontology of the Judith River Beds, with a chapter on the Fossil Plants by F. H. Knowlton: Bull. U.S. Geol. Surv., no. 257, pp. 1-174, pls. i-xix.
- STAUB, M.—1888. Stand der phytopaläontologischen Sammlung der königl. ungarischen geologischen Anstalt am Ende des Jahres 1885.—II: Jahrb. k. ungar. geol. Anst., pp. 203-234.
- 1905. Die Geschichte des Genus *Cinnamomum*. Pp. 138, pls. xxvi, 2 maps. Budapest.
- STEFANI, C. DE.—1881. Le Alghe fossili nelle rocce delle Alpi Apuane: Atti Soc. Tosc. Sci. Nat., vol. 2, pp. 280-282.
- STEINMANN, G.—1880. Zur Kenntniss fossiler Kalkalgen (Siphonocæen): Neues Jahrb. f. Min., vol. 2, pp. 130-140, pl. v.
- 1899. Ueber fossile Dasycladaceen vom Cerro Escamela: Beitr. Geol. Palæont. Republ. Mexico, von Felix & Lenk, pt. ii, pp. 189-204, text-figs. Leipzig.
- 1901. Ueber *Bouëina*, eine fossile Alge aus der Familie der Codiaceen: Bericht. naturf. Ges. Freiburg, vol. 11, pp. 62-72, text-figs. 1-13.
- 1903. *Tetraplopora Ramsi*, eine neue Dasycladacea aus dem Tithon von Stramberg: Beitr. Palæont. Geol. Oesterreich-Ungarns, vol. 15, pp. 45-54, text-figs. 1-11.

- STENZEL, K. G.—1872. Ueber fossile Palmenhölzer: Jahresb. schles. Ges. vaterl. Cultur, 1871, pp. 71-72.
- 1886. *Rhizodendron Oppoliense*, Goeppl.: Jahresb. schles. Ges. vaterl. Cultur, Ergänzungsheft 63 Jahr, pp. 1-30, pls. i-iii.
- 1897. Verkieselte Farne von Kamenz in Sachsen.—II. *Rhizocaulon antiquense*, n. sp.: Mitteil. k. min.-geol. Mus. Leipzig, vol. 13, pp. 1-24, pls. i-iii.
- 1904. Fossile Palmenhölzer: Beitr. Paläont. Geol. Oesterreich-Ungarns, vol. 16, pp. 107-287, pls. iii-xxiv.
- STERNBERG, K. von.—1820-38. Versuch einer geognostisch-botanischen Darstellung der Flora der Vorwelt. Pt. 1 (1820), pp. 24, pls. xiii; pt. 2 (1821), pp. 33, pls. xiv-xxvi; pt. 3 (1823), pp. 40, pls. xxvii-xxxix; pt. 4 (1825), pp. i-xlii and index, pls. xl-lix, A-E; pts. 5, 6 (1833), pp. 80, pls. i-xxvi; pts. 7, 8 (1838), pp. 81-200, pls. xxvii-lxviii. Leipzig & Prag.
See also CORDA.—1838.
- STIEHLER, A. W.—1854. Ueber fossile Pflanzen aus der Kreideformation von Quedlinburg: Zeitschr. deutsch. geol. Ges., vol. 6, pp. 659-662.
- 1857. Die Flora im Quadersandstein des Langenberges bei Quedlinburg: Zeitschr. f. gesamt. Naturwiss. Sachsen, vol. 9, pp. 452-455.
- 1858. Beiträge zur Kenntniss der vorweltlichen Flora des Kreidegebirges im Harze: Paläontogr., vol. 5, pp. 47-70, 71-80, pls. ix-xi, xii-xv.
- ? [unsigned] 1859. Ueber *Credneria*, Zenker: Ber. naturwiss. Ver. Harzes, p. 22.
- STOKES and WEBB.—1824. Description of some Fossil Vegetables of the Tilgate Forest in Sussex [no author's name given]: Trans. Geol. Soc., ser. 2, vol. 1, pp. 421bis-424, pls. xlv-xlvii.
- STOPES, M. C.—1909. Plant-containing Nodules from Japan, considered structurally in their Relation to the "Coal-Balls" and "Roof-Nodules" of the European Carboniferous: Quart. Journ. Geol. Soc., vol. 65, pp. 195-205, pl. ix.
- 1910 A. The Internal Anatomy of *Nilssonia orientalis*: Ann. Bot., vol. 24, pp. 389-393, pl. xxvi.
- 1910 B. Further Observations on the Fossil Flower, *Cretovarium*: Ann. Bot., vol. 24, pp. 679-681, pls. lvi-lvii.
- and FUJII, K.—1909. Studies on the Structure and Affinities of Cretaceous Plants: Proc. Roy. Soc. London, vol. 81 B, pp. 559-561.
- — 1910. Studies on the Structure and Affinities of Cretaceous Plants: Phil. Trans. Roy. Soc. London, vol. 201 B, pp. i-90, pls. i-ix.
- and KERSHAW, E. M.—1910. The Anatomy of Cretaceous Pine Leaves: Ann. Bot., vol. 24, pp. 395-402, pls. xxvii-xxviii.
- STRUB, D.—1863. Bericht über die geologische Uebersichtsaufnahme des südwestlichen Siebenbürgen im Sommer 1860: Jahrb. k.k. geol. Reichsanst., vol. 13, pp. 33-120.
- 1871. Geologie der Steiermark. Pp. 654. Graz.

- STUR, D.—1873. Vorkommen einer Palmenfrucht-Hülle = *Lepidocaryopsis Westphaleni*, n. g. et sp., im Kreide-Sandstein der Peruzer-Schichten bei Kaunitz in Böhmen: Verhandl. k. k. geol. Reichsanst., 1873, pp. 1-3.
- SUZUKI, Y.—1910. On the Structure and Affinities of Two New Conifers and a New Fungus from the Upper Cretaceous of Hokkaidō (Yezo): Bot. Mag. Tokyo, vol. 24, no. 284, pp. 181-196, pl. vii.
- TOMMASI, A.—1892. Contribuzione allo Studio della Fauna Cretacea del Friuli.—I Fossili Senoniani di Vernasso presso S. Pietro al Natisone: Atti Istit. Veneto Sci. (vol. 38), vol. 2, pp. 1089-1122, pl. unnumb.
- TOULA, F.—1884. Geologische Untersuchungen im westlichen Theile des Balkan und in den angrenzenden Gebieten: Sitzb. k. Akad. Wiss. Wien, vol. 88, pp. 1279-1348, pls. i-ix.
- TRAUTSCHOLD, H.—1876. Der Klin'sche Sandstein in Russland: Nouv. Mém. Soc. Imp. Nat. Moscou, vol. 13, pp. 191-236, pls. xviii-xxii.
- TRIEBEL, R.—1889. See CASPARY.—1889.
- TROOST, G.—1821. Description of a Variety of Amber and of a Fossil Substance supposed to be the Nest of an Insect, discovered at Cape Sable, Magothy River, Anne Arundel County, Maryland: Amer. Journ. Sci., vol. 3, pp. 8-15.
- TUZSON, J.—1908. Adatok Magyarországi fosszilis flórájához (Beiträge zur fossilen Flora Ungarns).—Additamenta ad floram fossilem Hungariæ. II: Beiblatt Növénytani Közlemények, Budapest, 1908, pp. 1-4 and (1)-(4), pls. i-iii.
- UDDEN, J. A.—1908. A Cycad from the Upper Cretaceous in Maverick County, Texas: Science, vol. 28, pp. 159-160.
- UNGER, F.—1842. In Endlicher's Genera Plantarum, suppl. ii, pp. 1-114.
- 1845. Synopsis Plantarum fossilium. Pp. 330. Leipzig.
- 1847. Chloris protogaea.—Beiträge zur Flora der Vorwelt. Pp. xxii, 149, pls. l. Leipzig.
- 1849. Botanische Beobachtungen.—VII. Einige interessante Pflanzenabdrücke aus der königl. Petrefactensammlung in München: Bot. Zeit., vol. 7, pp. 345-353.
- 1850 a. Genera et Species Plantarum fossilium. Pp. 627. Vienna.
- 1851. Die fossile Flora von Sotzka: Denkschr. k. Akad. Wiss. Wien, vol. 2, pp. 131-197, pls. xxii-lxviii.
- 1852. Beschreibung fossiler Hölzer aus Texas: in ROEMER, F.—1852, pp. 94-95.
- 1853. Iconographia Plantarum fossilium.—Abbildungen und Beschreibungen fossiler Pflanzen: Denkschr. k. Akad. Wiss. Wien, vol. 4, pp. 73-118, pls. xxiv-xlv.
- 1858. Ueber fossile Pflanzen des Süßwasser-Kalkes und Quarzes: Denkschr. k. Akad. Wiss. Wien, vol. 14, pp. 1-12, pls. i-iii.

- UNGER, F.—1859. Der versteinerte Wald bei Cairo und einige andere Lager verkieselten Holzes in Ägypten: Sitzb. k. Akad. Wiss. Wien, vol. 33, pp. 209–232, pls. i–iii.
- 1860. Sylloge Plantarum Fossilium.—Sammlung Fossiler Pflanzen besonders aus der Tertiärformation: Denkschr. k. Akad. Wiss. Wien, vol. 19, pp. 1–48, pls. i–xxi.
- 1864. Sylloge Plantarum Fossilium, pugillus secundus.—Sammlung Fossiler Pflanzen besonders aus der Tertiärformation: Denkschr. k. Akad. Wiss. Wien, vol. 22, pp. 1–36, pls. i–xii.
- 1865. Über einige fossile Pflanzenreste aus Siebenbürgen und Ungarn: Sitzb. k. Akad. Wiss. Wien, vol. 51, pp. 373–380, 1 pl.
- 1866. Fossile Pflanzenreste aus Neu-Seeland: in Reise der Oesterreichischen Fregatte Novara . . . Geologischer Theil, Bd. I, Abt. 2, Paläontologie von Neu-Seeland. Beiträge zur Kenntniss der fossilen Flora und Fauna der Provinzen Auckland und Nelson . . . Redigirt von F. von HOCHSTETTER, M. HÖRNES, und F. von HAUER. Pp. 1–13, pls. i–v. Vienna.
- 1866 A. Notiz über fossile Hölzer aus Abyssinien: Sitzb. k. Akad. Wiss. Wien, vol. 54, pp. 289–297, pl. i.
- 1867. Kreidepflanzen aus Oesterreich: Sitzb. k. Akad. Wiss. Wien, vol. 55, pp. 642–654, pls. i, ii.
- 1869. Die fossile Flora von Radoboj in ihrer Gesamtheit und nach ihrem Verhältnisse zur Entwicklung der Vegetation der Tertiärzeit: Denkschr. k. Akad. Wiss. Wien, vol. xxix, pp. 1–46, pls. i–v.
- 1870 A. Über Lieschkolben (*Typha*) der Vorwelt: Sitzb. k. Akad. Wiss. Wien, vol. 61, pt. 1, pp. 94–116, pls. i–iii.
- UPHAM, W.—1888. See WINCHELL, N. H., and UPHAM, W.—1888.
- VANHÖFFEN, E.—1897. Die Fauna & Flora Grönlands: in Grönland-Expedition der Gesellschaft für Erdkunde zu Berlin 1891–1893 unter Leitung von Erich von Drygalski, pp. 359–372, text-figs. Berlin.
- VASSEUR, G.—1890. Découverte d'une Flore turonienne dans les Environs des Martigues (Bouches-du-Rhône): Comptes Rendus Acad. Sci. Paris, vol. 110, pp. 1086–1089.
- VATER, H.—1884. Die fossilen Hölzer der Phosphoritlager des Herzogthums Braunschweig: Zeitschr. deutsch. geol. Ges., vol. 36, pp. 783–853, pls. xxvii–xxix.
- VELENOVSKY, J.—1882 A. Vorläufiger Bericht über die dicotyledonen Pflanzen der böhmischen Kreideformation: Sitzb. k. böhm. Ges. Wiss., 1881, pp. 212–219.
- 1882 B. Die Flora der böhmischen Kreideformation.—I. Crenneriaceae und Araliaceae: Beitr. Palaeont. Oesterreich-Ungarns, vol. 2, pp. 8–32, pls. iii–viii (v–x).
- 1882 C. Die Flora aus den Ausgebrannten tertiären Letten von Vrsovic bei Laun: Abhandl. k. böhm. Ges. Wiss., vol. 11, pp. 1–54, pls. i–x.

- VELENOVSKY, J.—1883. Die Flora der böhmischen Kreideformation.—II. Proteaceæ, Myricaceæ, Cupuliferæ, Moreæ, Magnoliaceæ, Bombaceæ: Beitr. Palæont. Oesterreich-Ungarns, vol. 3, pp. 26-47, pls. ix-xv.
- 1885. Die Gymnospermen der böhmischen Kreideformation. Pp. 34, pls. xiii. Prague.
- 1886. Die Flora der böhmischen Kreideformation.—III.: Beitr. Palæont. Oesterreich-Ungarns, vol. 4, pp. (1-14) 48-61, pls. (i-viii) xvi-xxiii.
- 1887. Die Flora der böhmischen Kreideformation.—IV.: Beitr. Palæont. Oesterreich-Ungarns, vol. 5, pp. (1-14) 62-75, pls. (i-viii) xxiv-xxxi.
- 1887 A. Neue Beiträge zur Kenntnis der Pflanzen des böhmischen Cenomans: Sitzb. k. böhm. Ges. Wiss., 1886, pp. 633-645, & 1 unnumb. pl.
- 1888 A. Ueber einige neue Pflanzenformen der böhmischen Kreideformation: Sitzb. k. böhm. Ges. Wiss., 1887, pp. 590-598, & 1 unnumbered pl.
- 1888 B. Die Farne der böhmischen Kreideformation: Abhandl. k. böhm. Ges. Wiss., [7] vol. 2, pp. 1-32, pls. i-vi.
- 1889. Kvetena českého Cenomanu: Abhandl. k. böhm. Ges. Wiss., [7] vol. 3, pp. 1-75, pls. i-vi.
- VREDENBURG, E. W.—1908. Pseudo-Fucoids from the Pab Sandstones at Fort Munro, and from the Vindhyan Series: Rec. Geol. Surv. India, vol. 36, pp. 241-253, pls. xxxi-xxxiv.
- WAAGEN, W.—1896. See KRASSER.—1896.
- WALCOTT, C. D.—1899. See HAGUE, A., *et alii*.—1899.
- WALKER, J. F.—1903. See LAMPLUGH and WALKER.—1903.
- WARD, L. F.—1884. On Mesozoic Dicotyledons: Amer. Journ. Sci., ser. 3, vol. 27, pp. 292-303.
- 1885. Synopsis of the Flora of the Laramie Group: 6th Ann. Rep. U.S. Geol. Surv., pp. 405-557, pls. xxxi-lxv.
- 1885 A. Sketch of Paleobotany: 5th Ann. Rep. U.S. Geol. Surv., pp. 363-452, pls. lvi-lviii.
- 1887. Types of the Laramie Flora: Bull. U.S. Geol. Surv., no. 37, pp. 1-115, pls. i-lvii.
- 1888. Evidence of the Fossil Plants as to the Age of the Potomac Formation: Amer. Journ. Sci., ser. 3, vol. 36, pp. 119-131.
- 1889. The Geographical Distribution of Fossil Plants: 8th Ann. Rep. U.S. Geol. Surv., pp. 669-960, & map.
- 1889 A. The Palæontological History of the Genus *Platanus*: Proc. U.S. Nat. Mus., vol. 11, pp. 39-42, pls. xvii-xxii.
- 1894. The Cretaceous Rim of the Black Hills: Journ. Geol., vol. 2, pp. 250-266.
- 1894 A. Recent Discoveries of Cycadean Trunks in the Potomac Formation of Maryland: Bull. Torrey Bot. Club. vol. 21, pp. 291-299.
- 1894 B. Fossil Cycadean Trunks of North America, with a Revision of the Genus *Cycadeoidea*, Buckland: Proc. Biol. Soc. Washington, vol. 9, pp. 75-87.

- WARD, L. F.—1895 a. The Potomac Formation: 15th Ann. Rep. U.S. Geol. Surv., pp. 313-397, pls. ii, iii, iv.
- 1895 b. The Mesozoic Flora of Portugal compared with that of the United States: Science, n.s., vol. 1, pp. 337-346.
- 1896. Some Analogies in the Lower Cretaceous of Europe and America: 16th Ann. Rep. U.S. Geol. Surv., pp. 463-540, pls. xcvii-cvii.
- 1897 a. A New Species of *Eucalyptus* from the Dakota Group of South-Western Kansas: Bull. Torrey Bot. Club, vol. 24, pp. 576-577, text-fig.
- 1897 b. Professor Fontaine and Dr. Newberry on the Age of the Potomac Formation: Science, n. s., vol. 5, pp. 411-423.
- 1897 c. Descriptions of the Species of *Cycadeoidea*, or Fossil Cycadean Trunks, thus far discovered in the Iron Ore Belt, Potomac Formation, of Maryland: Proc. Biol. Soc. Washington, vol. 11, pp. 1-17.
- 1899 a. Descriptions of the Species of *Cycadeoidea*, or Fossil Cycadean Trunks, thus far determined from the Lower Cretaceous Rim of the Black Hills: Proc. U.S. Nat. Mus., vol. 21, pp. 193-229.
- 1899 b. The Cretaceous Formation of the Black Hills as indicated by the Fossil Plants (with the collaboration of W. P. Jenney, W. M. Fontaine, and F. H. Knowlton): 19th Ann. Rep. U.S. Geol. Surv., pt. 2, pp. 521-712, pls. lvii-clxxii.
- 1900. Status of the Mesozoic Floras of the United States.—I. The Older Mesozoic: 20th Ann. Rep. U.S. Geol. Surv., pp. 217-430, pls. xxi-clxxix.
- 1900 a. Elaboration of the Fossil Cycads in the Yale Museum: Amer. Journ. Sci., ser. 4, vol. 10, pp. 327-345, pls. ii-iv.
- 1905. Status of the Mesozoic Floras of the United States (Second paper, with the collaboration of W. M. Fontaine, A. Bibbins, and G. R. Wieland): Mon. U.S. Geol. Surv., no. 48, pp. 616, pls. cxix.
- WATELET, A.—1866. Description des Plantes Fossiles du Bassin de Paris. Pp. 264, pls. ix. Paris.
- WEBER, O.—1852. Die Tertiärflora der Niederrheinischen Braunkohlenformation: Palæontogr., vol. 2, pp. 117-236, pls. xviii-xxv.
- 1856. See WESSEL and WEBER.—1856.
- WEED, W. H.—1899. See HAGUE, A., *et alii*.—1899.
- and KNOWLTON, F. H.—1893. The Laramie and the Overlying Livingston Formation in Montana, with Report on Flora: Bull. U.S. Geol. Surv., no. 105, pp. 1-68, pls. i-vi.
- and PIRSSON, L. V.—1898. Geology and Mineral Resources of the Judith Mountains of Montana: 18th Ann. Rep. U.S. Geol. Surv., pt. 3 [fossil plants by Fontaine, pp. 481, 482].
- WELSCH, J.—1897. Sur l'Age sénonien des grès à *Sabalites andegavensis* de l'ouest de la France: Comptes Rendus Acad. Sci. Paris, vol. 125, pp. 667-669.

- WESSEL, P., and WEBER, O.—1856. Neuer Beitrag zur Tertiärflora der niederrheinischen Braunkohlenformation: *Palaontogr.*, vol. 4, pp. 111-178, pls. xx-xxx.
- WHITE, C.A.—1889. See NEWBERRY, J. S.—1889.
- 1891. Correlation Papers.—Cretaceous: *Bull. U.S. Geol. Surv.*, no. 82, pp. 1-273, & maps.
- WHITE, D.—1890 A. On Cretaceous Plants from Martha's Vineyard: *Amer. Journ. Sci.*, ser. 3, vol. 39, pp. 93-101, pl. ii.
- 1890 B. Cretaceous Plants from Martha's Vineyard (Abstract): *Bull. Geol. Soc. America*, vol. 1, pp. 554-555.
- and SCHUCHERT, C.—1898. Cretaceous Series of the West Coast of Greenland: *Bull. Geol. Soc. America*, vol. 9, pp. 343-368, pls. xxv, xxvi.
- WIELAND, G. R.—1899 c. A Study of American Fossil Cycads.—III. The Female Fructifications of Cycadeoidea: *Amer. Journ. Sci.*, ser. 4, vol. 7, pp. 383-391, pls. viii-x.
- 1900. The Yale Collection of Fossil Cycads: *Yale Sci. Monthly*, vol. 6, pp. 211-221, pl. i.
- 1901. A Study of some American Fossil Cycads.—IV. On the Microsporangiate Fructification of *Cycadeoidea*: *Amer. Journ. Sci.*, ser. 4, vol. 11, pp. 423-436.
- 1903 A. Notes on the Marine Turtle *Archelon*.—II. Associated Fossils: *Amer. Journ. Sci.*, ser. 4, vol. 15, pp. 215, 216.
- 1903 B. Polar Climate in Time: *Amer. Journ. Sci.*, ser. 4, vol. 16, pp. 401-430.
- 1905. See WARD, L. F.—1905.
- 1906. American Fossil Cycads: *Publ. Carnegie Inst.*, no. 34. Pp. 296, pls. 1. Washington.
- 1908. Historic Fossil Cycads: *Amer. Journ. Sci.*, ser. 4, vol. 25, pp. 93-101, text-fig.
- 1910. Two new Araucarians from the Western Cretaceous: *Bull. S. Dakota Geol. Surv.* for 1908, pp. 77-81, pl. i.
- WILLIAMSON, W. C.—1887. On the Morphology of *Pinites oblongus* (*Abies oblonga* of Lindley and Hutton): *Mem. Manchester Lit. Phil. Soc.*, ser. 3, vol. 10, pp. 189-194, pl. ix.
- WILLISTON, S. W.—1905. The *Hallopus*-, *Baptanodon*-, and *Atlantosaurus*-beds of Marsh: *Journ. Geol.*, vol. 13, pp. 339-350.
- WIMMER, F.—1845. Neue Beiträge zur Flora von Schlesien zur Geschichte und Geographie derselben. Pp. 225. Breslau.
- WINCHELL, N. H.—1885. The Geological and Natural History Survey of Minnesota: 13th Ann. Rep. for Year 1884, pp. 76-77.
- and UPHAM, W.—1888. The Geology of Minnesota.—Vol. II. of the Final Report: *Geol. & Nat. Hist. Surv. Minnesota* for 1882-1885, pp. 1-695, pls. xxxii-lxi, map, & lettered plates. St. Paul.
- WINKLER, C.—1881. Review of Krendowskij, Beschreibung fossiler Bäume hauptsächlich aus dem Süden Russlands: *Bot. Centralbl.*, vol. 6, pp. 415-417.

- WOODWARD, H.—1885. Notes on some Mesozoic Plant-Remains from South Australia: *Geol. Mag.*, dec. 3, vol. 2, pp. 289-293, pl. vii.
- WOODWARD, H. B.—1887. The Geology of England and Wales; with Notes on the Physical Features of the Country, ed. 2. Pp. 670, with map. London.
- WORSDELL, W. C.—1903. The Affinities of the Mesozoic Fossil, *Bennettites Gibsonianus* Carr.: *Ann. Bot.*, vol. 14, pp. 717-721.¹
- YOKOYAMA, M.—1889. Jurassic Plants from Kaga, Hida, and Echizen: *Journ. Coll. Sci. Imp. Univ. Tokyo*, vol. 3, pp. 1-66, pls. i-xiv.
- 1894. Mesozoic Plants from Kōzuke, Kii, Awa, and Tosa: *Journ. Coll. Sci. Imp. Univ. Tokyo*, vol. 7, pp. 201-231, pls. xx-xxviii.
- 1906. Mesozoic Plants from China: *Journ. Coll. Sci. Imp. Univ. Tokyo*, vol. 21, art. 9, pp. 1-39, pls. i-xii.
- ZEILLER, R.—1885. Sur les affinités du genre *Laccopteris*: *Bull. Soc. bot. France*, vol. 32, pp. 21-25, text-fig.
- 1897. Revue des Travaux de Paléontologie végétale publiés dans le cours des années 1893-1896: *Rev. gén. Bot.*, vol. 9, pp. 459-462.
- 1900. *Éléments de Paléobotanique*. Pp. 421, text-figs. Paris.
- 1903. Revue des Travaux de Paléontologie végétale publiés dans le cours des années 1897-1900: *Rev. gén. Bot.*, vol. 15, pp. 330-336.
- 1905. Sur quelques Empreintes végétales de la Formation charbonneuse supracrétacée Balkans: *Ann. Mines*, ser. 10, vol. 7, pp. 326-349, pl. vii.
- 1908. Les Progrès de la Paléobotanique de l'ère des Gymnospermes: *Progress. Rei Bot.*, vol. 2, pp. 171-226.
- 1909. Revue des Travaux de Paléontologie végétale publiés dans le cours des années 1901-1906: *Rev. gén. Bot.*, vol. 20 [see pp. 167-173, 208-210].
- ZENKER, J. C.—1833. Beiträge zur Naturgeschichte der Urwelt. Pp. 67, pls. vi. Jena.
- ZIGNO, A. DE.—1864. Ueber die geognostische Zusammensetzung der Euganäischen Berge: *Zeitschr. deutsch. geol. Ges.* vol. 16, pp. 520-529.
- ZITTEL, K. A.—1890. See SCHIMPER and SCHENK.—1890.

Note on Brongniart's "Histoire des Végétaux Fossiles"

(see p. 5).

WHEN the MS. of this Literature was completed, I followed the usual British custom, and supposed the date on the title-page, 1828, was sufficient to quote for Brongniart's "Histoire." Since then some detailed work on the Carboniferous Flora has revealed to me that for nomenclatorial purposes this is insufficient. Mr. C. Davies Sherborn has kindly devoted his expert knowledge to the task of settling, so far as possible, the exact dates of the various parts, and has now fixed definitely three dates which even M. Zeiller's admirable bibliography * left uncertain. In the text throughout my volume, where the book has been quoted as 1828, reference to the following list for the page quoted, will give the exact date. I believe in no case in the present volume does the difference in date prove important.

BRONGNIART, A.—1828-1833, "Histoire des Végétaux Fossiles" etc. Paris and Amsterdam.

- | | |
|-----------------|--|
| Vol. I. part 1. | 1828, pp. xii, 1-80; pls. 1-9, 11, 13, 14, 16-18. |
| " " | 2. 1828, pp. 81-136; pls. 9 bis, 10, 12, 15, 19-27. |
| " " | 3. 1829, pp. 137-168; pls. 28, 30-36, 38-41, 44, 45. |
| " " | 4. 1829, pp. 169-208; pls. 29, 42, 43, 46-49, 51, 52, 54-56, 61, 63. |
| " " | 5. 1830, pp. 209-248; pls. 50, 53, 57, 58, 61 bis, 62, 64, 65, 67, 68, 70, 71, 73, 76. |
| " " | 6. 1831, pp. 249-264; pls. 59, 60, 63, 69, 72, 74, 75, 77-82. |
| " " | 7. 1833, pp. 265-288; pls. 83-97. |
| " " | 8. 1833, pp. 289-312; pls. 82 A, 98-109. |
| " " | 9. 1834, pp. 313-336; pls. 110-114, 117, 118, 124, 127, 128, 130. |
| " " | 10. 1836, pp. 337-368; pls. 115, 116, 119-123, 125, 126, 129, 131-134. |
| " " | 11. 1836, pp. 369-416; pls. 135-146. |
| " " | 12. 1836, pp. 417-488; pls. 37, 37 bis, 82 B, 147-160. |
| " " | 13. 1837, pls. 161-166. |
| Vol. II. | 13. 1837, pp. 1-24; pls. 1, 2, 14, 15, 18. |
| " " | 14. 1838, pp. 25-36; pls. 3-7, 22, 23, 26, 28, 30. |
| " " | 15. 1838, pp. 37-72; pls. 8-13, 16, 17, 19-21, 24, 25. |

Vol. II. unfinished.

* ZEILLER, R.—1903. "Flore fossile des Gites de Charbon du Tonkin." Paris, p. 306.

LIST OF SPECIES OF PLANTS

FROM THE

CRETACEOUS ROCKS

DESCRIBED IN WORKS UP TO 31ST DECEMBER, 1910.

THIS is an alphabetical list of all the specific names originally given to plant-remains from the Cretaceous formations, with a few references to plants first described from other formations and subsequently discovered in those of Cretaceous age. It is merely a list, without any critical observations as to synonymy; but when a new name is given to a species already known, a cross-reference is added. Mere names in lists and doubtful records, such as "*Pinus* sp.," are not included, unless a published figure or some other special circumstance justifies this course. The date following the author's name in each case corresponds with the entry in the Bibliography (pp. 1-47), which gives the full title of the work or paper quoted. Specific names are not amended, except in the case of obvious misprints, and no species have been re-named.

The names of the European Wealden plants (other than Angiosperms) are not included in this list, except those of a few species which range to higher horizons. The Wealden Flora has already been described by Prof. Seward in two earlier volumes of this series of Catalogues.

Aachenensaurus multidens, Smets in Dollo, 1888, p. 300. (= *Aachenoxylon*, Hovelacque, 1890, and *Nicolia Moresneti*, Hovelacque, 1890.)

Aachenoxylon, Hovelacque, 1890, p. 60, text-fig. 1, pl. iii, fig. 1.
(= *Aachenosaurus multidens*, Smets, in part.)

Upper Cretaceous; Moresnet.

Abies Benstedii, Mantell, 1843, p. 34; & 1846 (1843), p. 52, pl. ii, figs. 2, *a*, *b*, *c*; (& 1844, p. 166, pl. v, figs. 2, 2 *a*, 2 *b*, wood of same).

Lower Greensand; Maidstone.

— **Briarti** (Coemans), Schimper, 1872, p. 307. Cretaceous; Belgium.

— **calcaria**, Velenovsky, 1885, p. 33, pl. v, fig. 1, pl. vi, fig. 17.

Cenomanian; Bohemia.

— **chuchlensis**, Velenovsky, 1885, p. 34, pl. v, figs. 11-13.

Perucer Beds; Bohemia.

- Abies Linkii**, Roemer, 1829, p. 10, pl. xvii, fig. 2. (= *Abietites Linkii* (Roem.), Dunker, 1846. Recorded American Trinity div. by Fontaine, 1894.) Oolithen-gebirges; Northern Germany.
- **minor**, Velenovsky, 1885, p. 33, pl. v, figs. 14, 15, pl. viii, fig. 1. Cenomanian; Bohemia.
- **oblonga**, Lindley & Hutton, 1835, p. 155, pl. cxxxvii. (= *Abietites oblongus*, Goeppert, and *Pinites oblongus*, Williamson, 1887.) Greensand; Dorsetshire.
- **cblongata**, Buckland in Bensted, 1862, p. 337. (= *Abies oblonga* in Lindley & Hutton.) Greensand; Dorsetshire.
- (**Picea?**) **Omalii** (Coemans), Schimper, 1872, p. 307. (= *Pinus Omalii*, Coemans, 1867.) Cretaceous; Belgium.
- **Valentini**, Kurtz, 1902, p. 58. U. Cretaceous ("Dakota Beds"); Patagonia.
- Abietites acicularis**, Saporta, 1890 a, p. 814; & 1894, p. 92, pl. xvii, figs. 1, 2. Valanginian; Portugal.
- **angusticarpus**, Fontaine, 1889, p. 263, pl. cxxxiii, fig. 1. Potomac Formation; Virginia, U.S.A.
- **Benstedii** (Mantell), Goeppert. (= *Abies Benstedii*, Mantell, 1846.) Lower Greensand; Maidstone.
- **californicus**, Fontaine in Diller & Stanton, 1894, p. 450 [nomen nudum]. Horsetown Beds; California, U.S.A.
- **Chevalieri**, Fliche, 1896, p. 207, pl. ix, fig. 1. Albion; Clermont, France.
- **cretacea**, Newberry, 1898, p. 18, pl. xiv, fig. 5. Dakota Group; Dakota, U.S.A.
- **curvifolius**, Dunker, 1856, p. 18, pl. xxxiii, fig. 1. Quadersandstein; Blankenburg, Saxony.
- **ellipticus**, Fontaine, 1889, p. 263, pl. cxxxii, figs. 8, 9, pl. cxxxiii, figs. 2-4, pl. clxviii, fig. 8. Potomac Formation; Virginia, U.S.A.
- **Ernestinæ** (Stiehler), Lesquereux, 1874, p. 49, pl. i, fig. 7. (= *Pterophyllum Haydenii*, Lesquereux, in part.) Dakota Group; Nebraska, U.S.A.
- **Glueckii**, Richter, 1905, p. 7, pl. i, fig. 14. Upper Cretaceous; Blankenburg.
- **Goepperti**, Dunker, 1856, p. 180, pl. xxxii. Ibid.
- **Hartigi**, Dunker, 1856, p. 180, pl. xxxiii, fig. 2. (= *Aracurites Hartigi*, Schimper, 1872.) Ibid.
- **Linkii** (Roemer), Dunker, 1846, p. 18, pl. ix, figs. 11 a-c. (= *Abies Linkii*, Roemer, 1839.) (Recorded U.S.A. by Fontaine, 1894.) Wealden; North Germany.
- **macrocarpus**, Fontaine, 1889, p. 262, pl. cxxxii, fig. 7. Potomac Formation; Virginia, U.S.A.
- **marylandicus**, Fontaine in Ward, 1905, p. 549, pl. cxv, figs. 4, 5. Older Potomac Formation; Maryland, U.S.A.
- **oblongus**, Goeppert, 1850, p. 207. (= *Abies oblonga*, Lindley & Hutton, 1835.)
- **truncatus**, Saporta, 1867, p. 34.—Quadersandstein; Westphalia.

- Abietites Tyrrellii*, Dawson, 1886, p. 17.
Fort Pierre Group; North-West Territory, Canada.
- *Valentini*, Kurtz, 1902, p. 50.
U. Cretaceous (equivalent of Dakota Group); Patagonia.
- ? *sp.*, Fontaine in Ward, 1905, p. 262, pl. lxviii, fig. 17.
Shasta Formation; California, U.S.A.
- Abiocalis yezcensis*, Suzuki, 1910, p. 181, pl. vii, figs. 1, 2.
Upper Cretaceous; Hokkaido, Japan.
- Acaciæphyllum ellipticum*, Fontaine in Ward, 1905, p. 269, pl. lxix, fig. 18.
Shasta Formation; California, U.S.A.
- *longifolium*, Fontaine, 1889, p. 279, pl. exxxvii, fig. 6; pl. exxxviii, figs. 1-3.
Potomac Formation; Virginia, U.S.A.
- *microphyllum*, Fontaine, 1889, p. 280, pl. exxxviii, fig. 5. Ibid.
- *pachyphyllum*, Fontaine in Ward, 1905, p. 270, pl. lxix, fig. 19.
Shasta Formation; Knoxville, U.S.A.
- *spatulatum*, Fontaine, 1889, p. 280, pl. exxxviii, figs. 4, 6-9.
Potomac Formation; Virginia, U.S.A.
- *variabile*, Fontaine, 1889, p. 281, pl. clxx, fig. 7.
Potomac Formation; Maryland, U.S.A.
- Acacioxylon antiquum*, Schenk, 1883, p. 9.
Cretaceous (?); Libyan Desert.
- Acer amboyense*, Newberry, 1895, p. 106, pl. xlv, figs. 5-8 (seeds only).
Amboy Clay; Woodbridge, U.S.A.
- *antiquum*, Ettingshausen, 1867 A, p. 259, pl. iii, fig. 17.
Cenomanian; Niederschoena, Saxony.
- *caudatum*, Heer, 1883 A, p. 38, pl. lxx, figs. 1-2.
Patoot Beds; Greenland.
- ? *cretaceum*, Nilsson, 1832, p. 345, pl. i, figs. 1-2.
Greensand; Scania, Sweden.
- *edentatum*, Heer, 1883 A, p. 39, pl. lxx, fig. 3.
Patoot Beds; Greenland.
- *minutus*, Hollick, 1893, p. 35, pl. iii, fig. 6.
Raritan Formation (?); Staten Island, U.S.A.
- *obtusilobum* (?), Unger, 1847, p. 134, pl. xliii, figs. 12, 13. (Recorded American Cret. Lesquereux, 1868, p. 100.) (= *Menispermites salinae*, Knowlton, 1898 ?)
- *paucidentatum*, Hollick, 1898 D, p. 132, pl. xiv, figs. 2, 3.
Upper Cretaceous (Clay Marl); New Jersey, U.S.A.
- *Saskatchewanense*, Dawson, 1886, p. 16.
Belly River Series; North-West Territory, Canada.
- *sp.* (fruit), Hollick, 1906 A, p. 89, pl. xxxiii, figs. 12, 13.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Acerates amboyensis*, Berry, 1909, p. 263 (naming *Acerates* sp. ?,
Hollick in Newberry, 1895).
Raritan Formation; New Jersey, U.S.A.
- *arctica*, Heer, 1882, p. 82, pl. xxx, figs. 19, 20.
Atane Beds; Greenland.

97363/1

- Acerates** sp.?, Hollick in Newberry, 1895, p. 124, pl. xxxii, fig. 17;
pl. xli, figs. 4, 5. (Named *Acerates amboyensis*, Berry, 1909.)
Amboy Clays; New Jersey, U.S.A.
- Aceriphyllum aralicides**, Fontaine, 1889, p. 321, pl. clxiii, fig. 8.
Potomac Formation; Virginia, U.S.A.
- Acerites? cretaceus** (Nilsson), Brongniart, 1849 A, p. 111.
Cretaceous; Scania, Sweden.
- **Emmæ**, Stiehler, 1857, p. 58 [nomen nudum].
Quadersandstein; Blankenburg, Saxony.
- **menispermifolius**, Lesquereux, 1868, p. 101. (= *Menispermites acerifolia*, Lesquereux, 1874.) Dakota Group; Nebraska, U.S.A.
- **multiformis**, Lesquereux, 1892, p. 156, pl. xxxiv, figs. 1-9.
Dakota Group; Kansas, U.S.A.
- **pristinus**, Newberry, 1870, p. 15; & 1878, pl. v, fig. 4.
Dakota Group; Nebraska, U.S.A.
- **styracifolius**, Unger, 1850 A, p. 453 [not specifically named by
Reuss, 1846, p. 96, pl. li, figs. 4, 5].
Cenomanian (Pläner); Bohemia.
- Acrostichites Egedeanus**, Heer, 1874 A, p. 39, pl. iii, figs. 5, 5 b.
Kome Beds; Arkusak, Greenland.
- Acrostichophyllum cretaceum**, Velenovsky, 1889, p. 28, pl. ii,
figs. 22, 23. Cenomanian; Vyserovic, Bohemia.
- Acrostichopteris adiantifolia** (Fontaine), Berry, 1910 D, p. 629.
(= *Baieropsis adiantifolia*, Fontaine, 1889.)
Potomac Formation, &c.; Virginia, U.S.A.
- **cyclopteroides**, Fontaine, 1889, p. 109, pl. xciv, fig. 8. Ibid.
- **densifolia**, Fontaine, 1889, p. 107, pl. xciv, fig. 4, pl. clxx, fig. 11,
pl. clxxi, figs. 2, 6, pl. clxxii, fig. 13.
Potomac Formation; Maryland, U.S.A.
- **fimbriata**, Knowlton, 1907, p. 110, pl. xi, figs. 3, 3 A.
Kootanie Formation; Montana, U.S.A.
- **longipennis**, Fontaine, 1889, p. 107, pl. clxx, fig. 10, pl. clxxi,
figs. 1, 5, 7. Potomac Formation; Maryland, U.S.A.
- **parcelobata**, Fontaine, 1889, p. 108, pl. xciv, figs. 6, 7, 11, 14.
Potomac Formation; Virginia, U.S.A.
- **parvifolia**, Fontaine, 1889, p. 108, pl. xciv, figs. 5, 9, 10, 12,
pl. clxxi, figs. 3, 4, pl. clxxii, fig. 14.
Potomac Formation; Maryland and Virginia, U.S.A.
- **pluripartita** (Fontaine), Berry, 1910 D (re-naming *Baieropsis pluripartita*, *B. pluripartita minor*, and *B. longifolia* of Fontaine, 1889).
Potomac Formation, etc.; Virginia, U.S.A.
- Acrostichum crassifolium**, Fontaine, 1889, p. 105, pl. xvi, fig. 7.
Potomac Formation; Virginia, U.S.A.
- **cretaceum**, Velenovsky, 1889, p. 5, pl. ii, figs. 22, 23.
Cenomanian; Vyserovic, Bohemia.
- **Haddeni**, Hollick, 1902, p. 146, pl. iv, figs. 3-6.
Laramie Formation; Colorado, U.S.A.

- Acrostichum primordiale*, Ettingshausen, 1893, p. 147; & 1895,
p. 10, pl. i, fig. 1. Cretaceous; Australia.
- *tristaniaephyllum*, Bayer, 1899, p. 41, text-figs. 13, 13 a.
Perucer Beds; Kuchelbad, Bohemia.
- Adiantites cassebeeroides*, Debey & Ettingshausen, 1859 B, p. 195,
pl. ii, figs. 12-17. Senonian; Aix, Rhenish Prussia.
- *Decaisneanum*, Debey & Ettingshausen, 1859 B, p. 194, pl. ii,
fig. 10. Ibid.
- *marylandicus*, Fontaine in Ward, 1905, p. 549, pl. cxv, fig. 4, 5.
Potomac Formation; Vinegar Hill, U.S.A.
- *parvifolius*, Fontaine in Ward, 1905, p. 558, pl. cxvii, fig. 1.
Lower Potomac Formation; Virginia, U.S.A.
- *prælongus*, Dawson, 1883, p. 25, pl. v, fig. 19.
Upper Cretaceous; Vancouver Island, Canada.
- *yuasensis*, Yokoyama, 1894, p. 216, pl. xxi, fig. 15.
Neocomian; Yuasa, Japan.
- Adiantum aneimiaefolium*, Saporta, 1894, p. 82, pl. xv, fig. 21.
Valanginian; Portugal.
- *densinerve*, Heer, 1883 A, p. 2, pl. xlviii, fig. 15.
Patoot Beds; Patoot, Greenland.
- *dilaceratum*, Saporta, 1894, p. 164, pl. xxix, figs. 10-11, pl. xxxi,
fig. 7. Albian; Portugal.
- *eximium*, Saporta, 1894, p. 164, pl. xxviii, fig. 18, pl. xxxi, fig. 6.
Ibid.
- *expansum*, Saporta, 1894, p. 165, pl. xxviii, fig. 17. Ibid.
- *formosum*, Heer, 1871, p. 1181; & 1874 A, p. 35, pl. iii, figs. 1,
a, b, 2; pl. xiii, figs. 2 b, 11, 12. Kome Beds; Akrosak, Greenland.
- *formosum incisum*, Heer, 1874 A, p. 36, pl. xiii, figs. 11, 12.
Ibid.
- *montanense*, Knowlton, 1907, p. 112, pl. xii, figs. 1, 2.
Kootanie Formation; Montana, U.S.A.
- *subtilinervium*, Saporta, 1894, p. 82, pl. xviii, fig. 9.
Valanginian; Portugal.
- *tenellum*, Saporta, 1894, p. 163, pl. xxix, fig. 12, pl. xxxi, fig. 4.
Albian; Portugal.
- Adoxa præatavia*, Saporta, 1894, p. 187, pl. xxxiv, fig. 5. Ibid.
- Æcidites stellatus*, Debey & Ettingshausen, 1859, p. 212, pl. iii,
figs. 2-3. Senonian; Aix, Rhenish Prussia.
- Ægirophyllum* sp., Lampe, 1894, p. 197 [nomen nudum].
Cretaceous; Heimburg.
- Alcyonidiopsis longobardiæ*, Massalongo, 1856, p. 48, pl. vii,
figs. 1-2. Lower Cretaceous; Italy.
- Alethopteris elegans affinis*, Eichwald, 1865, p. 16.
Cretaceous; Russia.
- *elongata*, Feistmantel, 1874, p. 268.
Perucer Beds; Kounitz, Bohemia.
- *odontopteroides*, Feistmantel, 1874, p. 268.
Perucer Beds; Mscheno, Bohemia.

- Alethopteris Reichiana** (Brongniart), Presl in Sternberg, 1838, p. 146.
 (= *Pecopteris Reichiana*, Brongniart, 1828.)
 Cenomanian; Niederschoenna, Saxony.
 ----- **recentior**, Ettingshausen, 1852 c, p. 16, pl. iii, figs. 17, 18.
 Wernsdorfer Beds.
 ----- **revoluta**, Schulze, 1888, p. 10 [nomen nudum].
 Neocomian; Helmstein.
Aleurites eocenica, Lesquereux, 1873, p. 397.
 Laramie Formation; Wyoming, U.S.A.
Alisma? **reticulata**, Heer, 1882, p. 59, pl. xv, fig. 1.
 Atane Beds; Greenland.
Alismacites dakotensis, Lesquereux, 1892, p. 37, pl. ii, fig. 10.
 Dakota Group; Kansas, U.S.A.
 ----- **primævus**, Saporta, 1894, p. 96, pl. xv, fig. 31, pl. xvi, fig. 13 a.
 Valanginian; Portugal.
 ----- **primigenius**, Saporta, 1890 a, p. 814.
 Ibid.
Alnites crassus, Lesquereux, 1895, p. 13, pl. B, fig. 4.
 Dakota Group; Minnesota, U.S.A.
 ----- **? Friesii** (Nilsson), Brongniart, 1849 a, p. 111.
 Cretaceous; Scania, Sweden.
 ----- **grandifolia**, Newberry, 1870, p. 9; & 1878, pl. iv, fig. 2; & 1898,
 p. 67, pl. iv, fig. 2. Dakota Group; Nebraska, U.S.A.
 ----- **insignis**, Dawson, 1883, p. 28, pl. viii, fig. 38.
 Upper Cretaceous; Vancouver Island, Canada.
 ----- **petiolatus**, Lesquereux, 1878, p. 10 [nomen nudum].
 ----- **quadrangularis**, Lesquereux, 1874, p. 62, pl. iv, fig. 1. (= *Hamamelites quadrangularis*, Lesquereux, 1876, and *Populites quadrangularis*, Lesquereux, 1868.)
 Dakota Group; Kansas, U.S.A.
 ----- **speciosus**, Eichwald, 1865, p. 58, pl. iii, fig. 15.
 Cretaceous; Russia.
Alnus? **Friesii**, Nilsson, 1832, p. 346, pl. i, fig. 7. Greensand; Sweden.
Alnus grandifolia, Newberry, in Reynolds, 1869, p. 164 [nomen nudum].
 Nebraska, U.S.A.
 ----- **Grewiopsis**, Ward, 1887, p. 30, pl. xiv, fig. 1.
 Laramie Formation; Wyoming, U.S.A.
 ----- **kanseana**, Lesquereux, 1874, p. 62, pl. xxx, fig. 8. (= *Hamamelites Kansaseana*, 1876, and *H. Kansaseanus*, 1883, re-named *Quercus Kansana* by Knowlton, 1898.)
 Dakota Group; Kansas, U.S.A.
 ----- **protogæa**, Heer, 1883 a, p. 22, pl. iv, fig. 10.
 Patoot Beds; Greenland.
 ----- **? sp.**, Roemer, 1889, p. 144, pl. xii, figs. 7-8.
 Senonian; Bunzlau, Silesia.
Alsophilina kauniciana, Dormitzer in Krejci, 1853, p. 28, pl. i.
 (= *Oncopteris Kauniciana* (Dorm. sp.) Velenovsky, 1888.)
 Perucer Beds; Kaunitz, Bohemia.
Amelanchier Whitei, Hollick, 1906 a, p. 83, pl. xxxii, fig. 1.
 Middle Cretaceous; Martha's Vineyard, U.S.A.

- Ampelophyllum attenuatum**, Lesquereux, 1876 A, p. 396; & 1876 B, p. 354, pl. ii, fig. 3. Dakota Group; Kansas, U.S.A.
- **firmum**, Lesquereux, 1876 A, p. 396. Ibid.
- **ovatum**, Lesquereux, 1876 B, p. 355; & 1883, p. 69. (= *Populites ovata*, Lesquereux, 1868, and *Celtis ? ovata*, Lesquereux, 1874.) Dakota Group; Nebraska, U.S.A.
- Amphibennet(t)ites Bleicheri**, Fliche, 1896, p. 163, pl. xiv, fig. 1; pl. v, fig. 2, text-fig. 3. Albian; Revigny, France.
- **Renaulti**, Fliche, 1893, p. 167, pl. v, fig. 3, text-fig. 4. Albian; Ardennes, France.
- Amphibryophyllum carinatum**, Debey in Mourlon, 1881, p. 133 [nomen nudum]. Cenomanian; Limbourg, Belgium.
- **plicatum**, Debey in Mourlon, 1881, p. 133 [nomen nudum]. Ibid.
- **verticillatum**, Debey in Mourlon, 1881, p. 133 [nomen nudum]. Ibid.
- Amygdalus taurica**, Eichwald, 1865, p. 67, pl. iii, fig. 22. Neocomian; Russia.
- Anacardites amissus**, Heer, 1882, p. 99, pl. xl, fig. 8. Atane Beds; Greenland.
- **antiquus**, Lesquereux, 1892, p. 156, pl. lvii, fig. 1. Dakota Group; Kansas, U.S.A.
- Andromeda acuminata**, Lesquereux, 1876 A, p. 593. Ibid.
- **affinis**, Lesquereux, 1876 B, p. 348, pl. iii, figs. 5, 5 a. Iaramie Formation; Montana, U.S.A.
- **australiensis**, Ettingshausen, 1893, p. 150; & 1895, p. 39, pl. iii, fig. 23. Cretaceous; Australia.
- **Cookii** (Newberry), Berry, 1909, p. 261 (re-naming *Andromeda flexuosa*, Newberry, 1896). Raritan Formation; New Jersey, U.S.A.
- **cretacea**, Lesquereux, 1892, p. 117, pl. xvii, figs. 17, 18, pl. xxiv, fig. 5. Dakota Group: Kansas, U.S.A.
- **flexuosa**, Newberry, 1895, p. 121, pl. xxxiv, figs. 1-5. (Re-named *Andromeda Cookii* by Berry in 1909.) Amboy Clay; Woodbridge, U.S.A.
- **grandifolia**, Berry, 1907, p. 204, pl. xv, fig. 3 (re-naming *Andromeda latifolia*, Newberry, 1895). Middle Cretaceous; North Carolina, U.S.A.
- **latifolia**, Newberry, 1895, p. 120, pl. xxxiii, figs. 6-10; pl. xxxiv, figs. 6-11; pl. xxxvi, fig. 10. (= *Andromeda grandifolia*, Berry, 1907.) Amboy Clay; Woodbridge, U.S.A.
- **linifolia**, Lesquereux, 1892, p. 118, pl. lii, fig. 5. Dakota Group; Kansas, U.S.A.
- **novæ-cæsareæ**, Hollick in Newberry, 1895, p. 121, pl. xlii, figs. 9-12, 28-31. Amboy Clay; New Jersey, U.S.A.
- **Parlatorii**, Heer in Capellini & Heer, 1867, p. 18, pl. i, fig. 5. Dakota Group; Nebraska, U.S.A.
- **Parlatorii longifolia**, Lesquereux, 1892, p. 116, pl. lxiv, fig. 19. Dakota Group; Kansas, U.S.A.

- Andromeda Pfaffiana*, Heer, 1882, p. 79, pl. xxv, fig. 6; pl. xxxviii, figs. 5-7; pl. xlv, fig. 12. Atane Beds; Greenland.
- *Snowii*, Lesquereux, 1892, p. 117, pl. xvii, fig. 16. Dakota Group; Kansas, U.S.A.
- *tenuinervis*, Lesquereux, 1892, p. 116, pl. xxxviii, fig. 7. Ibid.
- *Wardiana*, Lesquereux, 1892, p. 119, pl. lxiv, fig. 17. Ibid.
- *sp.*, Heer, 1874 A, p. 135, pl. xxvi, fig. 6 b. Cretaceous; Greenland.
- Androcvetia carolinensis*, Berry, 1910 A, p. 183, pl. xix, figs. 1-6. Middle Cretaceous; North Carolina, U.S.A.
- *statenensis*, Hollick & Jeffrey, 1909, p. 22, pl. iii, figs. 1-5; pl. vii, figs. 1-8; pl. viii, figs. 1-12; pl. xxviii, figs. 5-8; pl. xxix, figs. 1-6. Raritan Formation; Staten Island, U.S.A.
- Aneimia perplexa*, Hollick in Newberry, 1898, p. 3, pl. xv, figs. 1, 1 a; pl. xvi, fig. 3; pl. lxiii, figs. 1-4 (re-naming *Sphenopteris elongatum*, Newberry, 1863). Laramie Formation; Orcas Island, U.S.A.
- *robusta*, Hollick, 1902, p. 145, pl. iii, fig. 1. Laramie Formation; Colorado, U.S.A.
- *stricta*, Newberry, 1895, p. 38, pl. iii, figs. 1, 2. Amboy Clay; New Jersey, U.S.A.
- *supercretacea*, Hollick, 1902, p. 145, pl. iii, figs. 6, 7. Laramie Formation; Colorado, U.S.A.
- Aneimidium lobulatum*, Saporta, 1894, p. 129, pl. xxvi, fig. 23. Urgonian; Portugal.
- *minutulum*, Saporta, 1894, p. 130, pl. xxvi, fig. 13. Ibid.
- *Schimperi*, Heer, 1874 A, p. 36, pl. ii, fig. 19; pl. xv, fig. 5 c. Kome Beds; Greenland.
- *tenerum*, Saporta, 1894, p. 130, pl. xxvi, fig. 25. Urgonian; Portugal.
- *sp.*, Heer, 1874, p. 133, pl. xvi, fig. 3 a. Cretaceous; Greenland.
- Angiopteridium auriculatum*, Fontaine, 1889, p. 113, pl. vii, figs. 8-11; pl. xxviii, fig. 1. (Re-named Berry 1910 D, *Teniopteris auriculatum*.) Potomac Formation; Virginia, U.S.A.
- *canmorense*, Dawson, 1893, p. 83, text-fig. 2. Kootanie Formation; North-West Territory, Canada.
- *densinerve*, Fontaine, 1889, p. 115, pl. xxix, fig. 4. Potomac Formation; Virginia, U.S.A.
- *dentatum*, Fontaine, 1889, p. 117, pl. xxx, figs. 6, 7. Ibid.
- *ellipticum*, Fontaine, 1889, p. 114, pl. xxix, fig. 3. Ibid.
- *nervosum*, Fontaine, 1889, p. 114, pl. xxix, fig. 2. (Re-named, Berry 1910 D, *Teniopteris nervosum*.) Ibid.
- *oregonense*, Fontaine in Stanton, 1895, p. 22 [nomen nudum]. Horsetown Beds; Oregon, U.S.A.
- *cvatum*, Fontaine, 1889, p. 115, pl. xxix, figs. 6, 7. Potomac Formation; Virginia, U.S.A.
- *pachyphyllum*, Fontaine, 1889, p. 115, pl. xxix, fig. 5. Ibid.
- *strictinerve*, Fontaine, 1889, p. 116, pl. xxix, figs. 8, 9. Ibid.
- *strictinerve*, var. *latifolium*, Fontaine, 1889, p. 116, pl. xxx, figs. 1, 5. Ibid.

- Aniscophyllum semi-alatum**, Lesquereux, 1874, p. 98, pl. vi, figs. 1-5. (= *Quercus semialata*, Lesquereux, 1868.)
Dakota Group; Nebraska, U.S.A.
- **sp.**, Dawson, 1883, p. 28, pl. viii, fig. 34.
Upper Cretaceous; Vancouver Island, Canada.
- Anomaspis hispida**, Hollick & Jeffrey, 1909, p. 50, pl. x, figs. 4, 8, 9.
Raritan Formation; New York, U.S.A.
- **tuberculata**, Hollick & Jeffrey, 1909, p. 49, pl. x, figs. 5, 6; pl. xxv, fig. 5; pl. xxvi, fig. 1. Ibid.
- Anomozamites acutiloba**, Heer, 1876 c, p. 102, pl. xxiii, fig. 1 a; pl. xxiv, figs. 1-3; pl. xxv, fig. 9; pl. xxviii, fig. 3 b. Jurassic.
Recorded for Canadian Kootanie Formation, Dawson, 1886.
- **angustifolius**, Fontaine, 1889, p. 167, pl. xxx, figs. 2, 3.
Potomac Formation; Virginia, U.S.A.
- **cretaceus**, Heer, 1874 A, p. 70, pl. xvi, figs. 19, 20.
Kome Beds; Greenland.
- **virginicus**, Fontaine, 1889, p. 168, pl. xxx, fig. 4; pl. xxxi, fig. 3.
Potomac Formation; Virginia, U.S.A.
- **sp.**, Dawson, 1893, p. 91. Kootanie Formation; British Columbia.
- Ancna cretacea**, Lesquereux, 1883, p. 77.
Dakota Group; Kansas, U.S.A.
- **robusta**, Lesquereux, 1883, p. 124, pl. xx, fig. 4.
Laramie Formation; Colorado, U.S.A.
- Anthocephale bohémica**, Bayer, 1893, pp. 32, 47, text-fig. 22; & Bayer in Fritsch, 1893, p. 132, text-fig. 193.
Senonian; Priesen, Bohemia.
- Antholithes horridus**, Dawson, 1886, p. 7.
Upper Cretaceous; North-West Territory, Canada.
- Antholithus Gaudium-Rosæ**, Ward, 1895 A, p. 355, pl. iii, fig. 7.
Potomac Formation; Virginia, U.S.A.
- **nymphæoides**, Hosius, 1870 A, p. 102, pl. xvii, figs. 35-36.
Cretaceous; Westphalia.
- Apeibopsis cyclophylla**, Lesquereux, 1892, p. 180, pl. xxv, fig. 6.
Dakota Group; Kansas, U.S.A.
- **Thomseniana**, Heer, 1882, p. 95, pl. xxxvi, fig. 5.
Atane Beds; Greenland.
- Apocynophyllum cretaceum**, Ettingshausen, 1867 A, p. 258, pl. iii, fig. 19.
Cenomanian; Niederschoena, Saxony.
- **cuneatum**, Hosius & von der Mark, 1880, p. 170, pl. xxxii, fig. 106.
Upper Senonian; Haldem, Westphalia.
- **sordidum**, Lesquereux, 1892, p. 109, pl. lxvi, fig. 11.
Dakota Group; Kansas, U.S.A.
- **subrepandum**, von der Mark, 1864, p. 79, pl. xiii, fig. 5.
Upper Senonian; Westphalia.
- **warraghianum**, Ettingshausen, 1893, pp. 138, 149; & 1895, p. 31, pl. iii, figs. 5-6.
Cretaceous; Australia.

- Aralia anisoloba*, Velenovsky, 1882 A, p. 214; & 1882 B, p. 22 (15), pl. v (iii), figs. 4-6. (= *Araliphyllum anisolum*, Velenovsky, 1889, and *Aralia subformosa anisoloba*, Ettingshausen, 1895.)
Cenomanian; Bohemia.
- *berberidifolia*, Lesquereux, 1892, p. 135, pl. xvi, fig. 11.
Dakota Group; Kansas, U.S.A.
- *Brittoniana*, Berry, 1905 B, p. 96, pl. xiv, fig. 3.
Matawan Formation; New Jersey, U.S.A.
- *calomacropa*, Saporta, 1894, p. 188, pl. xxxiv, figs. 15-16; pl. xxxv, fig. 1, 2, 4.
Albian; Portugal.
- *Chlomekensis*, Velenovsky, 1882 A, p. 214 [nomen nudum].
Cretaceous; Bohemia.
- *Chlomekiana*, Velenovsky, 1882 B, p. 20 (13), pl. v (iii), fig. 3.
Cenomanian; Bohemia.
- *concreta*, Lesquereux, 1876 A, p. 394; & 1876 B, p. 349, pl. iv, figs. 2-4. (= *A. semiorbiculata*, Lx.)
Dakota Group; Kansas, U.S.A.
- *coriacea*, Velenovsky, 1886, p. 11 (58), pl. i (xvi), figs. 1-9; pl. ii (xvii), fig. 2.
Cenomanian; Bohemia.
- *Credneriaefolia*, Velenovsky, 1882 A, p. 214 [nomen nudum].
Cretaceous; Bohemia.
- *Daphnophyllum*, Velenovsky, 1882 A, p. 214; & 1882 B, p. 30 (23), pl. vii (v), figs. 5, 6, 7, 8, 10; pl. viii (vi), figs. 1-5.
Cenomanian; Bohemia.
- *decurrens*, Velenovsky, 1886, p. 11 (51), pl. iv (xix), figs. 5-7. (= *Aralia subformosa decurrens*, Ettingshausen, 1895.)
Ibid.
- *denticulata*, Hosijs & von der Marek, 1880, p. 170, pl. xxxii, figs. 107, 107 A.
Upper Senonian; Haldem, Westphalia.
- (*Panax*) *dentifera*, Velenovsky, 1886, p. 13 (60), pl. ii (xvii), figs. 3-5.
Cenomanian; Bohemia.
- *dubia*, Fontaine, 1889, p. 314, pl. clvii, figs. 1-7. (= *Aralia Fontainei*, Knowlton, 1898.)
Potomac Formation; Virginia, U.S.A.
- *elegans*, Velenovsky, 1886, p. 13 (60), pl. iv (xix), fig. 1. (= *Aralia furcata*, Vel.)
Cenomanian; Vyserovic, Bohemia.
- *Fontainei*, Knowlton, 1898, p. 37 (re-naming *A. dubia*, Fontaine).
Potomac Formation; Virginia, U.S.A.
- *formosa*, Heer, 1869 A, p. 18, pl. viii, fig. 3. (= *Araliphyllum formosum* (Heer) Velenovsky, 1889.)
Upper Cretaceous; Moletuin, Moravia.
- *furcata*, Velenovsky, in Fritsch & Bayer, 1901, p. 152. (= *Araliphyllum furcatum*, Velenovsky.)
Peruczer Beds; Bohemia.
- *grœnlandica*, Heer, 1882, p. 84, pl. xxxviii, fig. 3; pl. xxxix, fig. 1; pl. xlvi, figs. 16-17.
Atane Beds; Greenland.
- *Kowalewskiana*, Saporta & Marion, 1878, p. 12. (= *Araliphyllum Kowalewskianum* in Velenovsky, 1889.)
Cenomanian; Bohemia.
- *Masoni*, Lesquereux, 1892, p. 133, pl. xv, fig. 4.
Dakota Group; Kansas, U.S.A.

- Aralia mattenwanensis**, Berry, 1905 a, p. 95, pl. xliii, fig. 2; pl. xlii, fig. 6. Matawan Formation; New Jersey, U.S.A.
- **microphylla**, Hosijs & von der Marek, 1880, p. 171, pl. xxxii, fig. 108. Upper Senonian; Haldem, Westphalia.
- **minor**, Velenovsky, 1882 a, p. 214; & 1882 b, p. 25 (18), pl. v (iii) fig. 9. (= *Aralia subformosa minor*, Ettingshausen, 1895.) Cenomanian; Vyserovic, Bohemia.
- **nassauensis**, Hollick, 1894 a, p. 55, pl. clxxiv, figs. 3, 7. Cretaceous; Long Island, U.S.A.
- **Newberryi**, Berry, 1907, p. 201, pl. xv, fig. 1 (re-naming *A. palmata*, Newberry). Middle Cretaceous; N. Carolina, U.S.A.
- **palmata**, Newberry, 1895, p. 117, pl. xxxix, figs. 6, 7; pl. xl, fig. 3. (= *Aralia Newberryi*, Berry, 1907.) Amboy Clay; Woodbridge, U.S.A.
- **partita**, Velenovsky, 1882 a, p. 214 [nomen nudum]. Cretaceous; Bohemia.
- **patens**, Newberry MS. in Hollick, 1894 a, p. 54, pl. clxxiv, fig. 4; & Newberry, 1895, p. 117, pl. xxviii, fig. 3. Amboy Clay; Long Island, U.S.A.
- **polymorpha**, Newberry, 1895, p. 118, pl. xxxix, figs. 1-5. Amboy Clay; Woodbridge, U.S.A.
- **propinqua**, Velenovsky, 1882 a, p. 214; & 1882 b, p. 29 (22), pl. vii (v), figs. 9-12; pl. viii (vi), fig. 6; pl. ix (vii), figs. 1-3, 6; pl. x (viii), fig. 1. Cenomanian; Vyserovic, Bohemia.
- **propinqua angustifolia**, Velenovsky, 1882 a, p. 214 [nomen nudum]. Cretaceous; Bohemia.
- **propinqua latifolia**, Velenovsky, 1882 a, p. 214 [nomen nudum]. Ibid.
- **proxima**, Saporta, 1894, p. 189, pl. xxxv, fig. 3 a. Albian; Portugal.
- **pungens**, Lesquereux, 1883, p. 123, pl. xix, figs. 3-4. Laramie Formation; Colorado, U.S.A.
- **quinquepartita**, Lesquereux, 1872, p. 302; & 1874, p. 90, pl. xv, fig. 6. Dakota Group; Kansas, U.S.A.
- **radiata**, Lesquereux, 1883, p. 64, pl. vii, figs. 2-3. Ibid.
- **Ravniana**, Heer, 1882, p. 84, pl. xxxviii, figs. 1-2. Atane Beds; Greenland.
- **rotundata**, Dawson, 1886, p. 14, pl. iv, fig. 5. Mill Creek Series; Mill Creek, Canada.
- **rotundiloba**, Newberry, 1895, p. 118, pl. xxviii, fig. 5; pl. xxxvi, fig. 9. Amboy Clay; Woodbridge, U.S.A.
- **Saportana** (also spelt **Saportanea**), Lesquereux, 1876 a, p. 394; & 1876 b, p. 350, pl. i, figs. 2, 2 a. Dakota Group; Kansas, U.S.A.
- **Saportanea deformata**, Lesquereux, 1892, p. 131, pl. xxiii, figs. 1-2. Ibid.
- **semi-orbiculata**, Lesquereux, 1876 a, p. 395. (= *A. concreta*, Lx., 1876 b.) Ibid.
- **submarginata**, Lesquereux, 1883, p. 63; & 1892, p. 133, pl. xv, fig. 3. Ibid.

- Aralia subformosa*, Ettingshausen, 1893, p. 150; & 1895, p. 40, pl. iv, fig. 4. Cretaceous; Australia.
- *subformosa angustiloba*, Ettingshausen, 1895, p. 42. (= *Aralia formosa* figured by Velenovsky, 1882, pl. v, fig. 2.)
- *subformosa anisoloba*, Ettingshausen, 1895, p. 41. (= *Aralia anisoloba*, Velenovsky, 1882.)
- *subformosa crenulata*, Ettingshausen, 1895, p. 41. Cretaceous; Australia.
- *subformosa decurrens*, Ettingshausen, 1895, p. 42. (= *Aralia decurrens*, Velenovsky, 1886.)
- *subformosa dentata*, Ettingshausen, 1895, p. 40. Cretaceous; Australia.
- *subformosa denticulata*, Ettingshausen, 1895, p. 41. (= *Aralia formosa* figured by Velenovsky, pl. vi, fig. 7.)
- *subformosa integriloba*, Ettingshausen, 1895, p. 42 (incl. *Aralia triloba*, Velenovsky, 1882.)
- *subformosa minor*, Ettingshausen, 1895, p. 42. (= *Aralia minor*, Velenovsky, 1882.)
- *subformosa pluriloba*, Ettingshausen, 1895, p. 42 (incl. *Aralia Kowalewskiana* figured Velenovsky, 1882, pl. v, fig. 1, & pl. vi, fig. 1).
- *subformosa quinqueloba*, Ettingshausen, 1895, p. 42 (incl. *Aralia Kowalewskiana* figured Velenovsky, 1882, pl. vi, figs. 1-2).
- *tenera*, Velenovsky, 1882 A, p. 214 [nomen nudum]. Cretaceous; Bohemia.
- *tenuinervis*, Lesquereux, 1883, p. 63, pl. vii, fig. 4. Dakota Group; Kansas, U.S.A.
- *Towneri*, Lesquereux, 1876 A, p. 395; & 1876 B, p. 349, pl. iv, fig. 1. Ibid.
- *transitiva*, Velenovsky, 1882 A, p. 214; & 1882 B, p. 28 (21), pl. vi (iv), figs. 8-10. (= *Araliphyllum transitivum*, Velenovsky, 1889.) Cenomanian; Kaunic, Bohemia.
- *triloba*, Velenovsky (*non* Newberry), 1882 A, p. 214; & 1882 B, p. 23 (16), pl. v (iii), figs. 7, 8. Cenomanian; Vyserovic, Bohemia.
- *tripartita*, Lesquereux, *non* Saporta, 1876 A, p. 394; & 1876 B, p. 348, pl. i, fig. 1. (= *Sterculia tripartita* [Lx.], Knowlton, 1898, p. 224.) Dakota Group; Kansas, U.S.A.
- ? *vernonsensis*, Fontaine in Ward, 1905, p. 492, pl. cvii, fig. 6. Older Potomac Formation; Virginia, U.S.A.
- *Waigattensis*, Heer, 1883 A, p. 36, pl. lx, fig. 5. Patoot Beds; Greenland.
- *washingtoniana*, Berry, 1910 II, p. 27, pl. viii, fig. 4. Magothy Formation; Columbia, U.S.A.
- *wellingtoniana*, Lesquereux, 1892, p. 131, pl. xxi, fig. 1; pl. xxii, figs. 2-3. Dakota Group; Kansas, U.S.A.
- *wellingtoniana Vaughanii*, Knowlton, 1901, p. 317. Dakota Group; Texas, U.S.A.

- Aralia Westonii**, Dawson, 1886, p. 14, pl. iv, fig. 6.
Mill Creek Series; Mill Creek, Canada
- **Wiesneri**, Krasser & Kubart, 1906, p. 47 [nomen nudum].
Cretaceous; Molettein, Moravia.
- **sp.**, Krasser, 1896, p. 136 (24), pl. xvi (vi), figs. 5, 5a.
Cretaceous; Kunstadt, Moravia.
- Araliaphyllum aceroides**, Fontaine, 1889, p. 319, pl. clvi, fig. 11;
pl. clxii, fig. 2. Potomac Formation; Virginia, U.S.A.
- **acuticlobum**, Fontaine, 1889, p. 318, pl. clxiii, fig. 2. Ibid.
- **haldemianum**, Debey, MS., in Saporta, 1873, p. 60, pl. vii,
figs. 1, 2. Upper Senonian; Haldem, Westphalia.
- **magnifolium**, Fontaine, 1889, p. 318, pl. clxix, figs. 9, 10.
Potomac Formation; Virginia, U.S.A.
- **obtusilcbum**, Fontaine, 1889, p. 317, pl. clxiii, figs. 1, 4; pl. clxiv,
fig. 3. Ibid.
- Araliphyllum anisclobum**, Velenovsky, 1889, pp. 47, 50, 54.
(= *Aralia anisoloba*, Velenovsky, 1882.) Cenomanian; Bohemia.
- **formosum** (Heer), Velenovsky, 1889, pp. 50, 54, 59. (= *Aralia*
formosa, Heer, 1869.) Ibid.
- **Kowalewskianum**, Velenovsky, 1889, pp. 50, 54, 57. (= *Aralia*
Kowalewskiana, Saporta & Marion, 1878.) Ibid.
- **transitivum**, Velenovsky, 1889, p. 119. (= *Aralia transitiva*,
Velenovsky.) Ibid.
- Araucaria acutifolia**, Corda in Reuss, 1846, p. 94, pl. xlviii, figs. 13-
15. Albian; Luschnitz, Bohemia.
- **bladenensis**, Berry, 1908 A, p. 255, pl. xii; pl. xiii; pl. xiv, figs. 1-3.
Middle Cretaceous; N. Carolina, U.S.A.
- **bohémica**, Velenovsky, 1889, p. 8, pl. i, figs. 20-24.
Cenomanian; Lipenec, Bohemia.
- **brachyphylla**, Bayer in Fritsch, 1894, p. 129, text-fig. 178; &
Bayer, 1893, pp. 5, 36, fig. 3. Senonian; Priesen, Bohemia.
- **Clarkii**, Berry, 1910 A, p. 182.
Middle Cretaceous; North Carolina, U.S.A.
- **crassifolia**, Corda in Reuss, 1846, p. 94, pl. xlviii, fig. 12.
Albian; Luschnitz, Bohemia.
- **cretacea** (Brongniart), Saporta in Schimper, 1872, p. 255.
Greensand; France.
- **epactridifolia**, Bayer in Fritsch, 1893, p. 24; & Bayer, 1893,
pp. 4, 36, fig. 2. Senonian; Priesen, Bohemia.
- **Frici**, Velenovsky in Fritsch, 1893, p. 128, text-fig. 177.
Priesener Beds; Priesen, Bohemia.
- **Hatcheri**, Wieland, 1910, p. 80, fig. 2 on plate. Wyoming, U.S.A.
- **hespera**, Wieland, 1910, p. 78, fig. 1 on plate, & pl. xxii.
S. Dakota, U.S.A.
- **insulinensis**, Fliche, 1896, p. 177, pl. v, fig. 5, text-fig. 5.
Albian; Islettes, France.
- **Jeffreyi**, Berry, 1908 A, p. 258, pl. xvi.
Middle Cretaceous; N. Carolina, U.S.A.

- Araucaria latifolia**, Bozzi, *non* Feistmantel, in Tommasi, 1892, p. 1119
[nomen nudum]. Senonian (?); Italy.
- **macrophylla**, Bozzi, 1891, p. 375, pl. xvi, figs. 1-2.
Cretaceous (?); Vernasso, Italy.
- **Miqueli**, Debey, 1877, p. 110 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **obtusifolia**, Fontaine, 1889, p. 249, pl. lxxxv, fig. 13.
Potomac Formation; Virginia, U.S.A.
- **podocarpoides**, Fontaine, 1889, p. 249, pl. lxxxvi, fig. 4. Ibid.
- **reperta**, Fliche, 1896, p. 174, pl. vi, fig. 2.
Albian; Clermont, France.
- **revigniacensis**, Fliche, 1896, p. 176, pl. v, fig. 4.
Albian; Revigny, France.
- **spatulata**, Newberry, 1868, p. 10; & 1878, pl. ii, figs. 5, 5 a.
Dakota Group; Nebraska, U.S.A.
- **Toucasii**, Saporta, 1879, p. 198, text-fig. 272.
Turonian; Toulon, France.
- **vernassiensis**, Bozzi in Tommasi, 1892, p. 1120 [nomen nudum].
Senonian (?); Italy.
- **zamioides**, Fontaine, 1889, p. 250, pl. cxxi, figs. 1-1 a.
Potomac Formation; Virginia, U.S.A.
- ? — **sp.**, Jeffrey, 1906, p. 388, pl. xxviii, figs. 14, 15.
Middle Cretaceous; New York, U.S.A.
- Araucario-cauln breveradiatum**, Lignier, 1907, p. 290, pl. xix,
figs. 33-43; pl. xxi, fig. 67, diagr. 10; pl. xxii, figs. 75-78; pl. xxiii,
figs. 79-81; text-fig. 2, p. 283. Cenomanian; Dives, France.
- Araucario-phloios breveradiatum**, Lignier, 1907, p. 291. Ibid.
- Araucariopitys americana**, Jeffrey, 1907, p. 435, pls. xxviii-xxx.
Middle Cretaceous; Staten Island, U.S.A.
- Araucarioxylon ægyptiacum**, Kraus in Schimper, 1874, p. 383; &
in Schenk, 1883, p. 3, pl. i, figs. 1-2; pl. ii, fig. 3. (= *Dadoxylon*
ægyptiacum, Unger, 1858.)
Cenomanian or Lower Turonian; Libyan Desert.
- **albianum**, Fliche, 1896 A, p. 182, pl. iv, figs. 2-4.
Albian; Aube, France.
- **armeniacum**, Gürich, 1885, p. 433. Cretaceous; Transcaucasia.
- **barremianum**, Fliche, 1900, p. 26, pl. ii, figs. 2-4.
Lower Cretaceous; Vassy, France.
- **Gardoniense**, Crié, 1890, p. 235, pl. i, figs. 1-4.
Cenomanian; Ile de Aix, France.
- **Hoppertcnæ**, Knowlton in Ward, 1899, p. 644, pls. clxiii-clix.
Dakota Group; S. Dakota, U.S.A.
- **madagascariense**, Fliche, 1900 A, p. 472, text-fig. 1.
Senonian; Madagascar.
- **noveboracense**, Hollick & Jeffrey, 1909, p. 58, pl. xxi, figs. 1-3,
5, 6. Raritan Formation; Staten Island, U.S.A.
- **Prosseri**, Penhallow, 1900, p. 77.
Comanche Cretaceous; Kansas, U.S.A.

- Araucarioxylon Tankcense**, Stopes & Fujii, 1910, p. 41, pl. iii, figs. 17-18. Upper Cretaceous; Hokkaido, Japan.
- **virginianum**, Knowlton, 1889 A, p. 50, pl. vii, figs. 2-5; & 1889 B, p. 106. Potomac Formation (Trias?); Virginia, U.S.A.
- **zeelandicum**, Crié, 1889, p. 29 [nomen nudum]. Cretaceous; New Zealand.
- Araucarites acutifolius**, Endlicher, 1847 B, p. 301. (= *Araucaria acutifolia*, Corda.) Middle Cretaceous; Bohemia.
- **adpressus**, von der Marck, 1864, p. 80, pl. xiii, figs. 10-11. Upper Senonian; Drensteinfurth, Westphalia.
- **ægyptiacus**, Goepfert, 1865 A, p. 259. (= *Araucarioxylon ægyptiacum*, Kraus, 1874.)
- **aquiensis** Fontaine, 1889, p. 264, pl. cxxiii, figs. 8-12. Potomac Formation; Virginia, U.S.A.
- **argillicola**, Eichwald in Merck, 1853, p. 304 [nomen nudum]. Neocomian; Russia.
- **crassifolius**, Endlicher, 1847 B, p. 301. (= *Araucaria crassifolia*, Corda.) Middle Cretaceous; Bohemia.
- **cuneatus**, Ward, 1899, p. 670, pl. clxiii, fig. 10. Dakota Formation; Blackhills, U.S.A.
- **Hartigi** (Dunker), Schimper, 1872, p. 252. Quadersandstein; Blankenburg, Saxony.
- **inflatus**, Krendowsky, 1881, p. 275, pl. ii, figs. 14-19. Cretaceous; Russia.
- **Nordenskiöldi**, Heer, 1874 A, p. 125, pl. xxxvii, figs. 3, 4. Cretaceous; Spitzbergen.
- **ovatus**, Hollick, 1898 A, p. 128, pl. xii, figs. 3 a, 4. Upper Cretaceous (Clay Marl); New Jersey, U.S.A.
- **patagonica**, Kurtz, 1902, p. 49 (cone scale). Upper Cretaceous; Patagonia.
- **Reichenbachi**, Geinitz, 1842, p. 98, pl. xxiv, fig. 4. (= *Sequoia Reichenbachi*, Heer, 1868. = *Geinitzia Reichenbachi*, Hollick & Jeffrey, 1909.) Quadersandstein; Saxony.
- **virginicus**, Fontaine, 1889, p. 263, pl. cxxiv, fig. 7. Potomac Formation; Virginia, U.S.A.
- **? Wardi**, Hill, 1893, p. 39, pl. i, figs. 1 a-d. (This is *Porocystis pruniformis*, Cragin, see Rauff, 1895.) Lower Cretaceous; Texas, U.S.A.
- **wyomingensis**, Fontaine in Ward, 1899, p. 669, pl. clxiii, figs. 1-9. Lower Cretaceous; Wyoming, U.S.A.
- **Zeilleri**, Berry, 1908 A, p. 252, pl. xi, fig. 3. Middle Cretaceous; New Jersey, U.S.A.
- Ardisia glcssa**, Bayer, 1893, pp. 25 & 44, text-fig. 17; & Bayer in Fritsch, 1893, p. 131, text-fig. 189. Senonian; Priesen, Bohemia.
- Arisæma cretacea**, Lesquereux, 1892, p. 38, pl. xlvi, fig. 1. Dakota Group; Kansas, U.S.A.
- **? dubia**, Hollick, 1898 A, p. 130, pl. xii, fig. 6. Upper Cretaceous (Clay Marl); New Jersey, U.S.A.
- **? mattewanse**, Hollick, 1898 A, p. 130, pl. xii, fig. 7. Ibid.

- Aristolochia Daveauana*, Saporta, 1894, p. 183, pl. xxxv, fig. 10.
Albian; Portugal.
- *tecomæcarpa*, Bayer, 1899, p. 29; pl. i, figs. 7, 8, text-figs. 10, 10a.
Perucér Beds; Vyserovic, Bohemia.
- Aristolochiaphyllum* (?) *cellulare*, Ward, 1905, p. 492, pl. cviii, fig. 5.
Potomac Formation; Virginia, U.S.A.
- Aristolochites dentata*, Heer in Capellini & Heer, 1867, p. 18, pl. ii, figs. 1, 2.
Dakota Group; Nebraska, U.S.A.
- *infundibuliformis*, Lesquereux, 1876 A, p. 393.
Dakota Group; Kansas, U.S.A.
- Aristoloiaphyllum crassinerve*, Fontaine, 1889, p. 322, pl. clx, figs. 3-6.
Potomac Formation; Virginia, U.S.A.
- Arthropitys mirabilis*, Eichwald, 1865, p. 32, pl. v, fig. 9.
Cretaceous; Russia.
- Arthrotaxis* sp., Nathorst, 1891, p. 34, pl. i, figs. 12-14.
Cretaceous (?); Königs-Wusterhausen.
- Artocarpidium cretaceum*, Ettingshausen, 1867 A, p. 251, pl. ii, fig. 4.
Cenomanian; Niederschoena, Saxony.
- *Guillemainii*, Menzel, 1909, p. 401, pl. ii, figs. 4, 5, 6.
Senonian; Kamerun.
- *pseudo-cretaceum*, Ettingshausen, 1893, p. 148; & 1895, p. 22, pl. ii, fig. 11.
Cretaceous; Australia.
- Artocarpophyllum occidentale*, Dawson, 1894, p. 60, pl. xii, fig. 51; pl. xiii, fig. 52.
Upper Cretaceous; Vancouver Island, Canada.
- Artocarpus Dicksoni*, Nathorst, 1890 A, p. 6, pl. i, figs. 1-4.
Upper Cretaceous; Greenland.
- *Lessigiana* (Lesquereux), Knowlton, 1893 c, p. 24.
Lower Laramie Formation; Colorado, U.S.A.
- *undulata*, Hosius, 1870 A, p. 100, pl. xvi, fig. 29.
Senonian; Legden, Westphalia.
- Arundinites oppelensis*, Roemer, 1870, p. 291, pl. xxvii, fig. 6.
Cenomanian; Oppeln, Upper Silesia.
- *Wohlfarthi*, Otto, 1854, p. 27, pl. iv, fig. 2; pl. vii, figs. 1-5.
(= *Cycadeites*? *Wohlfarthi* (Otto), Geinitz, 1875 b.)
Quadersandstein; Saxony.
- Arundo cretaceus*, Lesquereux, 1868, p. 92. (= *Phragmites cretaceus*, Lx., 1874.)
Dakota Group; Nebraska, U.S.A.
- *grœnlandica*, Heer, 1874 A, p. 104, pl. xxviii, figs. 8-11.
Atane Beds; Greenland.
- Asimina eccenica*, Lesquereux, 1873, p. 387; & 1878 b, p. 251, pl. xliii, figs. 7, 8.
Laramie Formation; Wyoming, U.S.A.
- Aspidiophyllum dentatum*, Lesquereux, 1883, p. 88; & 1892, p. 212, pl. xxxix, fig. 1.
Dakota Group; Kansas, U.S.A.
- *platanifolium*, Lesquereux, 1883, p. 88, pl. ii, fig. 4. Ibid.
- *trilobatum*, Lesquereux, 1876 n, p. 331, pl. ii, figs. 1, 2. Ibid.

- Aspidium angustipinnatum**, Fontaine, 1889, p. 98, pl. xvi, figs. 1, 3, 8; pl. xvii, fig. 1; pl. xix, fig. 10. (= *Dryopteris angustipinnata* (Font.), Knowlton, 1898, p. 91; re-named.)
Potomac Formation; Virginia, U.S.A.
- **angustipinnatum montanense**, Fontaine, 1893, p. 491, pl. lxxiv, figs. 1, 1a. (= *Dryopteris angustipinnata montanense* (Font.), Knowlton, 1898, p. 91; re-named.)
Kootanie Formation; Montana, U.S.A.
- **cretaceo-zeelandicum**, Ettingshausen, 1887 A, p. 174 (34), pl. vii, figs. 2, 3. Upper Cretaceous; New Zealand.
- **cystopteroides**, Fontaine, 1889, p. 99, pl. xvi, fig. 2. (= *Dryopteris cystopteroides* (Font.), Knowlton, 1898, p. 91; re-named.)
Potomac Formation; Virginia, U.S.A.
- **dentatum**, Fontaine, 1889, p. 102, pl. xxv, figs. 6, 7. (= *Dryopteris dentata* (Fontaine), Knowlton, 1898, p. 91; re-named.) Ibid.
- **Dunkeri** (Schimper), Fontaine, 1889, p. 101, pl. xxii, fig. 9; pl. xxv, figs. 11, 12; pl. xxvi, figs. 2, 8, 9, 18. (= *Cladophlebis Dunkeri* (Schimp.), Seward, 1894 A, p. 100.) Ibid.
- **ellipticum**, Fontaine, 1889, p. 95, pl. xiii, figs. 9, 10. (= *Dryopteris elliptica* (Font.), Knowlton, 1898, p. 92; re-named.) Ibid.
- **fecundum**, Heer, 1882, p. 32, pl. xxix, figs. 5, 9.
Atane Beds; Greenland.
- **Foersteri**, Renault, 1888, p. 352 [nomen nudum].
Cretaceous; France.
- **fredericksburgense**, Fontaine, 1889, p. 94, pl. xi, figs. 1-6; pl. xii, figs. 1-6; pl. xvi, fig. 9; pl. xix, figs. 6, 7. (= *Dryopteris fredericksburgense* (Font.), Knowlton, 1898, p. 92; re-named.)
Potomac Formation; Virginia, U.S.A.
- **heterophyllum**, Fontaine, 1889, p. 96, pl. xiv, figs. 1-5; pl. xv, figs. 1-5. (= *Dryopteris heterophylla* (Font.), Knowlton, 1898, p. 92; re-named.) Ibid.
- **Jenseni**, Heer, 1882, p. 31, pl. xvi, fig. 4; pl. xxix, fig. 4; pl. xxx, figs. 1-6.
Atane Beds; Greenland.
- **Kennerlyi**, Newberry, 1863, p. 513; & Hollick in Newberry, 1898, p. 11, pl. xvi, figs. 4, 5. (= *Dryopteris Kennerlyi* (Newberry) Knowlton, 1898.)
Upper Cretaceous; British Columbia.
- **macrocarpum**, Fontaine, 1889, p. 103, pl. xvii, fig. 2. (= *Dryopteris macrocarpa* (Font.), Knowlton, 1898, p. 92; re-named.)
Potomac Formation; Virginia, U.S.A.
- **microcarpum**, Fontaine, 1889, p. 103, pl. lix, figs. 2, 12; pl. lx, figs. 6, 7. (= *Dryopteris microcarpa* (Font.), Knowlton, 1898, p. 92; re-named.) Ibid.
- **monocarpum**, Fontaine, 1893, p. 490, pl. lxxxiii, figs. 4-6, 6a; pl. lxxxiv, figs. 3, 3a. (= *Dryopteris monocarpa* (Font.), Knowlton, 1898, p. 92; re-named.)
Kootanie Formation; Montana, U.S.A.
- **montanense**, Fontaine, 1893, p. 490, pl. lxxxiii, figs. 1-3; pl. lxxxiii, figs. 2, 3, 3a. (= *Dryopteris montanense* (Font.), Knowlton, 1898, p. 92; re-named.) Ibid.

- Aspidium oblongifolium**, Fontaine, 1889, p. 100, pl. xxi, fig. 5.
(*Dryopteris oblongifolia* (Font.), Knowlton, 1898, p. 92; re-named.)
Potomac Formation; Virginia, U.S.A.
- **Cerstedii**, Heer, 1882, p. 30, pl. xxxiv, figs. 3, 4. (= *Dryopteris Cerstedii* (Heer), Knowlton, 1898, p. 92; re-named.)
Atane Beds; Greenland.
- **parvifolium**, Fontaine, 1889, p. 100, pl. xxi, fig. 6; pl. xxiv, fig. 8;
pl. xxv, fig. 10; pl. xxvi, figs. 1, 14, 16, 17. (= *Dryopteris parvifolia*
(Fontaine), Knowlton, 1898; re-named.)
Potomac Formation; Virginia, U.S.A.
- **pinnatifidum**, Fontaine, 1889, p. 101, pl. xxi, fig. 15. (= *Dryopteris pinnatifida* (Fontaine), Knowlton, 1898; re-named.) Ibid.
- **Schowii**, Heer, 1882, p. 31, pl. xxxii, figs. 10, 10 b.
Atane Beds; Greenland.
- **ursinum**, Heer, 1870, p. 462, pl. xxxix, fig. 6 a. (Recorded by
Heer, 1880 b, from Kome Beds.) Miocene; Greenland.
- **virginicum**, Fontaine, 1889, p. 97, pl. xv, fig. 7; pl. xxi, fig. 14.
(= *Dryopteris virginica* (Fontaine), Knowlton, 1898; re-named.)
Potomac Formation; Virginia, U.S.A.
- Aspleniopteris adiantifolia**, Fontaine, 1889, p. 118, pl. xvi, fig. 6.
Ibid.
- **Nilssoni**, Sternberg, 1825, p. xxii, pl. xliii, figs. 3, 4, 5.
Greensand; Hör, Sweden.
- **pinnatifida**, Fontaine, 1889, p. 118, pl. xxii, figs. 1-3, 6, 7.
Potomac Formation; Virginia, U.S.A.
- Asplenites dubius**, Velenovsky, 1888 b, p. 16, pl. ii, figs. 17-19.
Chlomeka Beds; Leipa, Bohemia.
- **Trevirani**, Debey, 1848, p. 116 [nomen nudum].
Quadersandstein; Bavaria.
- **sp.**, Krasser, 1896, p. 122 (?), pl. xiv (iv), figs. 8, 8 a.
Cenomanian; Moletein, Moravia.
- Asplenium albertum**, Dawson, 1886, p. 11, pl. iii, fig. 6.
Mill Creek Series; Mill Creek, Canada.
- **Boyeaenum**, Heer, 1874 a, p. 33, pl. xi, fig. 9.
Kome Beds; Greenland.
- **Brongniarti**, Debey & Ettingshausen, 1859 b, p. 193 (13), pl. ii,
figs. 1-3. (= *Hausmannia* (?) *Brongniarti* (Debey & Ettingshausen),
Richter, 1906.) Senonian; Aix, Rhenish Prussia.
- **cænopteroides**, Debey & Ettingshausen, 1859 b, p. 194 (14), pl. ii,
figs. 8-9. Senonian; Aix, Rhenish Prussia.
- **calopteris** (Debey & Ettingshausen), Heer, 1883 a, p. 5. (= *Denizia calopteris*, Debey & Ettingshausen.) Patoot Beds; Greenland.
- **Dicksonianum**, Heer, 1874 a, p. 31, pl. i, figs. 1-5.
Kome Beds; Greenland.
- **dubium**, Fontaine, 1889, p. 109, pl. x, fig. 9.
Potomac Formation; Virginia, U.S.A.
- **Foersteri**, Debey & Ettingshausen, 1859 b, p. 193 (13), pl. ii,
figs. 4-7, 11. Senonian; Aix, Rhenish Prussia.

- Asplenium Haguei*, Knowlton, 1899 B, p. 655, pl. lxxvii, figs. 1, 2.
Laramie Formation; Yellowstone Park, U.S.A.
- *Johnstrupi* (Heer), Schimper, 1869, p. 660; & Heer, 1874 A, p. 32,
pl. i, figs. 6, 7. (= *Sphenopteris Johnstrupi*, Heer.)
Kome Beds; Greenland.
- *lapideum*, Heer, 1882, p. 3, pl. ii, figs. 3, 3 b. Ibid.
- *magnum*, Knowlton, 1896, p. 450 [nomen nudum].
Laramie Formation; Yellowstone Park, U.S.A.
- *Martinianum*, Dawson, 1886, p. 5, pl. i, fig. 1.
Kootanie Formation; Martin Creek, Canada.
- *Nauckhoffianum*, Heer, 1880 B, p. 3, pl. i, figs. 9-12.
Kome Beds; Greenland.
- *Niobrara*, Dawson, 1883, p. 20, pl. i, fig. 1.
Upper Cretaceous; Peace River, North-West Territory.
- *Nordenskiöldi*, Heer, 1874 A, p. 33, pl. ii, figs. 17 a, 17 b.
Kome Beds; Greenland.
- *Nordstroemi*, Heer, 1874 A, p. 93, pl. xxvi, fig. 6 a.
Atane Beds; Greenland.
- *Palæopteris*, Unger, 1866, p. 3, pl. i, figs. 4-8.
Lower Cretaceous; New Zealand.
- *Pringelianum*, Heer, 1883 A, p. 4, pl. xlviii, figs. 9, 9 b.
Patoot Beds; Patoot, Greenland.
- *raritanense*, Berry, 1909, p. 246, pl. xviii, fig. 1.
Raritan Formation; New Jersey, U.S.A.
- *scrobiculatum*, Heer, 1883 A, p. 4, pl. xlviii, figs. 10, 10 b.
Patoot Beds; Patoot, Greenland.
- *tenellum*, Knowlton, 1900, p. 19, pl. iii, figs. 1, 2.
Montana Formation; Wyoming, U.S.A.
- *Velenovskyi*, Marik, 1901, p. 6, pl. i, fig. 13.
Cenomanian; Bohemia.
- *wyomingense*, Knowlton, 1900 A, p. 19, pl. iii, fig. 12.
Montana Formation; Wyoming, U.S.A.
- *sp.*, Knowlton, 1900 A, p. 20, pl. iii, fig. 11. Ibid.
- Asterophyllites cretaceus*, Feistmantel, 1874, p. 267. (= *Pseudo-asterophyllites cretaceus*, Velenovsky, 1887 A.)
Perucer Beds; Bohemia.
- Asterosoma radiforme*, Otto, 1854, p. 15, pl. ii, fig. 4; pl. iii,
figs. 1, 2. Quadersandstein; Saxony.
- Astrocaryopsis Sanctæ-Manehildæ*, Fliche, 1896, p. 276, pl. xiii,
figs. 4, 5, text-fig. 12. Cenomanian; Sainte-Menehould, France.
- Athrotaxopsis expansa*, Fontaine, 1889, p. 241, pl. cxiii, figs. 5, 6,
pl. cxv, fig. 2; pl. cxvi, fig. 5; pl. cxvii, fig. 6; pl. cxxxv, figs. 15,
18, 22. Potomac Formation; Virginia, U.S.A.
- *grandis*, Fontaine, 1889, p. 240, pl. cxiv, figs. 1-3; pl. cxvi, figs. 1-4;
pl. cxxxv, fig. 10. Ibid.
- *pachyphylla*, Fontaine, 1889, p. 242, pl. cxv, figs. 1, 3; pl. cxvii,
figs. 1, 3-5. Ibid.

- Athrotaxopsis tenuicaulis**, Fontaine, 1889, p. 241, pl. cxiv, figs. 4, 5
pl. cxv, fig. 4; pl. cxvi, fig. 6; pl. cxvii, fig. 2.
Potomac Formation; Virginia, U.S.A.
- Aulacolepis rhomboidalis**, Ettingshausen, 1893, p. 147; & 1895,
p. 12, pl. i, fig. 10. Cretaceous; Australia.
- Aulacophycus pedatus**, Heer, 1877, p. 143, pl. lviii, fig. 12.
Neocomian; Switzerland.
- Baiera arctica**, Heer, 1874 A, p. 37, pl. iii, fig. 3.
Kome Beds; Greenland.
- **brevifolia**, Newberry, 1891, p. 199, pl. xiv, fig. 3.
Kootanie Formation; Montana, U.S.A.
- **cretosa**, Schenk, 1871 A, p. 5, pl. i, fig. 7. (= *Sclerophyllina*
cretosa, Heer.) Wernsdorfer Beds; Austrian Silesia.
- **dichotoma?** Heer (*non* Braun), 1876 A, p. 49, pl. xxxii, fig. 1.
Cretaceous; Spitzbergen.
- **foliosa**, Fontaine, 1889, p. 213, pl. xciv, fig. 13.
Potomac Formation; Virginia, U.S.A.
- **grandis**, Heer, 1874 A, p. 37, pl. iii, fig. 4. Kome Beds; Greenland.
- **incurvata**, Heer, 1882, p. 45, pl. xiii, fig. 6.
Atane Beds; Greenland.
- **leptopoda**, Heer, 1882, p. 46, pl. xxviii, fig. 9. Ibid.
- **longifolia** (Pomel), Heer, 1876 c, p. 39, pl. viii, fig. 6. (= *Dicro-*
pteris longifolia, Pomel, 1849. Recorded Canadian Kootanie by
Dawson, 1886.) Jurassic; Spitzbergen.
- **sagittata**, Heer, 1882, p. 46, pl. xxx, fig. 18.
Atane Beds; Greenland.
- Bajera scanica**, Sternberg, 1825, p. xxviii, pl. xlvii, fig. 2.
Greensand; Hör, Sweden.
- Baieropsis adiantifolia**, Fontaine, 1889, p. 211, pl. xcii, figs. 8, 9;
pl. xciii, figs. 1-3; pl. xciv, figs. 2, 3. (Re-named by Berry,
1910 D, *Acrostichopteris adiantifolia*).
Potomac Formation; Virginia, U.S.A.
- **adiantifolia**, var. **minor**, Fontaine, 1889, p. 212, pl. xciv, fig. 1.
Ibid.
- **denticulata**, Fontaine, 1889, p. 210, pl. xciii, fig. 7. Ibid.
- **denticulata**, var. **angustifolia**, Fontaine, 1889, p. 210, pl. xcii,
fig. 7. Ibid.
- **expansa**, Fontaine, 1889, p. 207, pl. lxxxix, figs. 1, 3; pl. xc, fig. 1;
pl. xci, fig. 2; pl. xcii, fig. 5.
Potomac Formation; Fredericksburg, U.S.A.
- **foliosa**, Fontaine, 1889, p. 209, pl. xciii, figs. 4-6. Ibid.
- **longifolia**, Fontaine, 1889, p. 210, pl. xci, fig. 6. (= *Acrosticho-*
pteris pluripartita (Fontaine), Berry, 1910 D.) Ibid.
- **macrophylla**, Fontaine, 1889, p. 212, pl. xc, fig. 6. Ibid.
- **pluripartita**, Fontaine, 1889, p. 208, pl. lxxxix, fig. 4; pl. xc,
figs. 2-5; pl. xci, figs. 1, 3, 4, 7; pl. xcii, figs. 1, 2, 6. (= *Acro-*
stichopteris pluripartita (Fontaine), Berry, 1910 D.) Ibid.

- Baieropsis pluripartita**, var. *minor*, Fontaine, 1889, p. 208, pl. xci, fig. 5; pl. xcii, figs. 3, 4. (= *Acrostichopteris pluripartita* (Fontaine), Berry, 1910 n.) Potomac Formation; Virginia, U.S.A.
- *sp.*, Dawson, 1892, p. 87. Kootanie Formation; British Columbia.
- Bambusites australis**, Ettingshausen, 1887 A, p. 180 (40), pl. vii, fig. 21; pl. viii, figs. 1-3, 3a. Upper Cretaceous; New Zealand.
- Bambusium latifolium**, Heer, 1881, p. 22, pl. xix, figs. 1-3, 1 c. Cretaceous; Portugal.
- *neocomense*, Heer, 1877, p. 146, pl. lviii, fig. 23. Neocomian; Switzerland.
- Banisteriophyllum cretaceum**, Ettingshausen, 1895, p. 47, pl. iv, fig. 2. Cretaceous; Australia.
- Banksia crenata**, Ettingshausen, 1893, pp. 140, 149; & 1895, p. 31, pl. iii, fig. 11. Cretaceous; Australia.
- *cretacea*, Ettingshausen, 1893, p. 149; & 1895, p. 28, pl. iii, figs. 9, 10. Ibid.
- *longifolia*, Ettingshausen, 1851, p. 730, pl. xxxi, fig. 19. (= *Myrica longifolia*, Unger.) Cenomanian; Niederschoena, Saxony.
- *plagioneura*, Ettingshausen, 1893, p. 141; & 1895, p. 29, pl. iii, fig. 8. Cretaceous; Australia.
- *prototypus*, Ettingshausen, 1851, p. 732; & 1852 A, p. 822, pl. lxiii, figs. 2, 3. Cenomanian; Niederschoena, Saxony.
- *pusilla*, Velenovsky, 1883, p. 7 (32), pl. i (ix), figs. 14-17. Cenomanian; Bohemia.
- *sub-longifolia*, Ettingshausen, 1893, p. 141; & 1895, p. 29, pl. iii, fig. 7. Cretaceous; Australia.
- Banksites pusillus**, Velenovsky, 1882 A, p. 213 [nomen nudum]. Cretaceous; Bohemia.
- *Saportanus*, Velenovsky, 1882 A, p. 213; & 1883, p. 7 (32), pl. i (ix), figs. 18-20. Cenomanian; Bohemia.
- Bauhinia cretacea**, Newberry, 1886 B, p. 77, pl. lvi, fig. 5. Middle Cretaceous; New Jersey, U.S.A.
- ? *gigantea*, Newberry, 1895, p. 93, pl. xx, fig. 1. Amboy Clay; Woodbridge, U.S.A.
- *marylandica*, Berry, 1908 c, p. 219, three text-figs. Magothy Formation; Maryland, U.S.A.
- Belodendron densifolium**, von der Marck, 1864, p. 80, pl. xiii, figs. 8, 9. Upper Senonian; Sendenhorst, Westphalia.
- *gracile*, Debey, 1849, p. 299 [nomen nudum]. Senonian; Aix, Rhenish Prussia.
- *lepidodendroides*, Debey, 1848, p. 121 [nomen nudum]. Ibid.
- *Neesii*, Debey, 1848 A, p. 121 [nomen nudum]. Ibid.
- Benizia calopteris**, Debey & Ettingshausen, 1859 B, p. 216 (36), pl. v figs. 13-17. (= *Asplenium calopteris*, Heer, 1883 A.) Senonian; Aix, Rhenish Prussia.

- Bennettites dacotensis**, McBride, 1893, p. 249, pl. xi, figs. 1, 2.
(= *Cycadeoidea dacotensis* of Ward, 1894).
Dakota Group; S. Dakota, U.S.A.
- **Gibsonianus**, Carruthers, 1870, p. 700, pls. lviii-lx. (= *Bennettites Gibsoni*, Carruthers, 1868.) Lower Greensand; Isle of Wight.
- **Gibsoni**, Carruthers [nomen nudum]. (= *Bennettites Gibsonianus*, Carruthers, 1870.)
- **Maraniana**, Scarabelli MS. in Capellini & Solms-Laubach, 1892, p. 110, pl. iii, fig. 4; pl. ii, fig. 3. Cretaceous (?); Italy.
- **maximus**, Carruthers, 1870, p. 699.
Lower Greensand; Shanklin, Isle of Wight.
- **Schachtii** (Coemans), Carruthers, 1870, p. 699. (= *Cycadeoidea Schachtii*, Capellini & Solms-Laubach, 1892.) Gault; Belgium.
- Benstedtia** sp. (Mackie), Seward, 1896 a, p. 219, pl. xiv, fig. 3. (= *Draena Benstedtii*, Mackie, 1862.) Greensand; Maidstone, Kent.
- Benthamia dubia**, Velenovsky, 1887, p. 11 (72), pl. vii (xxx), figs. 4, 6.
Perucer Beds; Vyserovic, Bohemia.
- Benzoia Masoni** (Lesquereux), Knowlton, 1898, p. 47 (re-naming *Lindera Masoni*, Lesquereux, 1892).
Dakota Group; Kansas, U.S.A.
- **venustum** (Lesquereux), Knowlton, 1898, p. 47 (re-naming *Lindera venusta*, Lesquereux, 1892).
Dakota Group; Kansas, U.S.A.
- Bergeria minuta**, Presl in Sternberg, 1838, p. 184, pl. xlix, figs. 2 a, b, 3.
Cenomanian; Saxony.
- Betula atavina**, Heer, 1883 a, p. 22, pl. lv, figs. 8, 21 b.
Patoot Beds; Greenland.
- **beatriciana**, Lesquereux, 1868, p. 95; & 1874, p. 61, pl. v, fig. 5; pl. xxx, fig. 4.
Dakota Group; Nebraska, U.S.A.
- **Goepperti**, Lesquereux, 1878 a, p. 138, pl. xvii, figs. 21-23.
Laramie Formation; Wyoming, U.S.A.
- **perantiqua**, Dawson, 1883, p. 27, pl. vii, fig. 27.
Upper Cretaceous; Vancouver Island, Canada.
- **Stevensonii**, Lesquereux, 1872, p. 293; & 1878 a, p. 139, pl. xviii, figs. 1-5.
Laramie Formation, Wyoming, U.S.A.
- **tremula**, Heer, 1883 a, p. 21, pl. liii, fig. 1 c; pl. lv, fig. 9.
Patoot Beds; Greenland.
- **vetusta**, Heer, 1883 a, p. 22, pl. lv, fig. 7.
Ibid.
- Betulites denticulata**, Heer in Capellini & Heer, 1867, p. 15, pl. iv, figs. 5, 6.
Dakota Group; Nebraska, U.S.A.
- ? **Hatcheri**, Knowlton in Stanton & Hatcher, 1905, p. 141, pl. xviii, fig. 3.
Judith River Beds; Montana, U.S.A.
- **populifolius**, Lesquereux, 1892, p. 64, pl. vi, figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- **rugosus**, Lesquereux, 1892, p. 65, pl. vi, figs. 3-5.
Ibid.
- **Snowii**, Lesquereux, 1892, p. 64, pl. v, figs. 1-4.
Ibid.
- **Westii**, Lesquereux, 1892, p. 60, pl. iv, figs. 1-22; pl. v, figs. 5-14.
Ibid.

- Betulites Westii*, var. *crassus*, Lesquereux, 1892, p. 63, pl. v, figs. 5-17. Dakota Group; Kansas, U.S.A.
- *Westii*, var. *cuneatus*, Lesquereux, 1892, p. 62, pl. v, fig. 8. Ibid.
- *Westii*, var. *grewiopsideus*, Lesquereux, 1892, p. 63, pl. lxiv, fig. 10. Ibid.
- *Westii*, var. *inæquilateralis*, Lesquereux, 1892, p. 62, pl. v, figs. 10-13. Ibid.
- *Westii*, var. *lanceclatus*, Lesquereux, 1892, p. 62, pl. v, fig. 14. Ibid.
- *Westii*, var. *latifolius*, Lesquereux, 1892, p. 61, pl. iv, figs. 9-11. Ibid.
- *Westii*, var. *multinervis*, Lesquereux, 1892, p. 62, pl. iv, figs. 20-22. Ibid.
- *Westii*, var. *oblongus*, Lesquereux, 1892, p. 61, pl. iv, figs. 17-19. Ibid.
- *Westii*, var. *populoides*, Lesquereux, 1892, p. 63. Ibid.
- *Westii*, var. *obtusus*, Lesquereux, 1892, p. 61, pl. iv, figs. 5-8. Ibid.
- *Westii*, var. *quadratifolius*, Lesquereux, 1892, p. 62, pl. v, fig. 9. Ibid.
- *Westii*, var. *reniformis*, Lesquereux, 1892, p. 62, pl. v, fig. 5. Ibid.
- *Westii*, var. *rhomboidalis*, Lesquereux, 1892, p. 62, pl. v, figs. 6, 7. Ibid.
- *Westii*, var. *rotundatus*, Lesquereux, 1892, p. 61, pl. iv, figs. 12-16. Ibid.
- *Westii*, var. *subintegrifolius*, Lesquereux, 1892, p. 61, pl. iv, figs. 1-4. Ibid.
- *sp.*, Lesquereux, 1892, p. 65, pl. v, fig. 18. Ibid.
- Dakota Group; Dakota, U.S.A.
- Bignonia cordata*, Velenovsky, 1887, p. 9 (70), pl. vi (xxix), fig. 5. (= *Bignoniophyllum cordatum*, Velenovsky, 1889.)
- Cenomanian; Bohemia.
- *pulcherrima*, Bayer, 1899, p. 33, pl. i, figs. 9-14; pl. ii, figs. 4-10; text-figs. 11, 12. Perucur Beds; Bohemia.
- *silesiaca*, Velenovsky, 1887, p. 8 (69), pl. vii (xxx), figs. 1, 2, 5, 11, 15. Cenomanian: Kieselingswalde.
- Bignoniophyllum cordatum*, Velenovsky, 1889, pp. 54, 53. (= *Bignonia cordata*, Velenovsky, 1885.)
- Blechnum priscum*, Ettingshausen, 1887 A, p. 174 (34), pl. vii, figs. 1, 1 a. Upper Cretaceous; New Zealand.
- *zeelandicum*, Crié, 1889, p. 88 (14) [nomen nudum]. Cretaceous; New Zealand.
- Blyttia infracretacea*, Saporta, 1894, p. 122, pl. xxiii, fig. 15; pl. xxiv, fig. 7 b. Urgonian; Portugal.
- Bombacophyllum argillaceum*, Velenovsky, 1889. (= *Bombax argillaceum*, Velenovsky, 1883.)

- Bombax argillaceum**, Velenovsky, 1882 A, p. 213; & 1883, p. 20 (45), pl. ii (x), figs. 17-19; pl. iv (xii), figs. 6-9. (= *Bombacophyllum argillaceum*, Velenovsky, 1889.) Cenomanian; Bohemia.
- **virginiensis**, Fontaine, 1889, p. 310, pl. cli, fig. 4.
Potomac Formation; Virginia, U.S.A.
- Bonaventurea cardinalis**, Debey & Ettingshausen, 1859 B, p. 203, pl. iii, figs. 2-19. Senonian; Aix, Rhenish Prussia.
- Boueina Hochstetteri**, Toulou, 1884, p. 1319, pls. vii-x; & Steinmann, 1901, pp. 1-5, text-figs. 1-7. Upper Neocomian; Servia.
- Bowerbankia attenuata**, Debey, 1849, p. 299 (31) [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **emarginata**, Debey, 1849, p. 299 [nomen nudum]. Ibid.
- **maxima**, Debey, 1849, p. 299 [nomen nudum]. Ibid.
- **repanda**, Debey, 1849, p. 299 [nomen nudum]. Ibid.
- **rotundifolia**, Debey, 1849, p. 299 [nomen nudum]. Ibid.
- Brachyoxylon notabile**, Hollick & Jeffrey, 1909, p. 54, pl. xiii, figs. 2-6; pl. xiv, fig. 2. Raritan Formation; Staten Island, U.S.A.
- Brachyphyllum Brardianum**, Brongniart, 1849 A, p. 110 [nomen nudum].
Lower Cretaceous; France.
- **confusum**, Saporta, 1894, p. 112, pl. xx, fig. 8. Aptian; Portugal.
- **corallinum**, Heer, 1881, p. 21, pl. xii, figs. 1-3, 1 aa.
Cretaceous; Portugal.
- **crassicaule**, Fontaine, 1889, p. 221, pl. c, fig. 4; pl. cix, figs. 1-7; pl. cx, figs. 1-3; pl. cxi, figs. 6, 7; pl. cxii, figs. 6-8; pl. clxviii, fig. 9.
Potomac Formation; Virginia, U.S.A.
- **crassum**, Lesquereux, 1892, p. 32, pl. ii, fig. 5.
Dakota Group; Kansas, U.S.A.
- **macrocarpum**, Newberry MS. in Newberry, 1895 (footnote by Hollick), p. 51; and for internal structure, Jeffrey, 1906 B, p. 384, pl. xxvii. Amboy Clay; New Jersey, U.S.A.
- **obesiforme**, Saporta, 1894, p. 176, pl. xxxi, figs. 12-13; pl. xxxiii, fig. 4 b; pl. xxxiv, fig. 8. Albian, Portugal.
- **obesiforme elongatum**, Saporta, 1894, p. 176, pl. xxxi, fig. 14. Ibid.
- **obesum**, Heer, 1881, p. 20, pl. xvii, figs. 1-4.
Cretaceous; Portugal.
- **Orbignianum**, Brongniart, 1849 B, p. 309; and d'Orbigny, 1852, p. 650, text-fig. 529. Cenomanian; Isle d'Aise.
- **parceramosum**, Fontaine, 1889, p. 223, pl. cx, fig. 4.
Potomac Formation; Virginia, U.S.A.
- **texense**, Fontaine, 1894, p. 269, pl. xxxviii, figs. 3-5; pl. xxxix, figs. 1, 1 a. Trinity Division; Texas, U.S.A.
- **sp. ?**, Fontaine, 1889, p. 224, pl. clxviii, fig. 2.
Dakota Group; Virginia, U.S.A.
- **sp. ?**, Fontaine, 1889, p. 224, pl. cxxxv, fig. 9. Ibid.
- **sp. ?**, Fontaine, 1889, p. 223, pl. cxxxv, fig. 8. Ibid.
- **sp. ?**, Hollick & Jeffrey, 1909, p. 37, pl. ix, figs. 5, 6; pl. vi, fig. 3; pl. xiv, fig. 3. Raritan Formation; New York, U.S.A.

- Brachyphyllum* sp., Jeffrey, 1906 b, p. 389, pl. xviii, figs. 16-17.
- Brasenia antiqua*, Dawson, 1886, p. 15, text-fig. (Re-named by Hollick, 1894 d, *Nelumbo Dawsoni*.)
Belly River Series; Medicine Hat, Canada.
- Braseniopsis venulosa*, Saporta, 1894, p. 192, pl. xxxiv, figs. 1-4.
Albian; Portugal.
- *villarsicoides*, Saporta, 1894, p. 195, pl. xxxv, fig. 9. Ibid.
- Bresciphyllum cretaceum*, Velenovsky, 1889, p. 25, pl. v, figs. 2, 3.
Cenomanian; Bohemia.
- Bromelia? tenuifolia*, Lesquereux, 1892, p. 41, pl. i, fig. 13.
Dakota Group; Kansas, U.S.A.
- Bronnites orientalis*, Unger, 1850, p. 524. Turkey.
- Bryocarpus monostachys*, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- *polystachys*, Debey, 1849, p. 299 [nomen nudum]. Ibid.
- Bumelia marcouana* (Heer), Lesquereux, 1874, p. 90, pl. xxviii, fig. 2.
(= *Leguminosites marcouanus*, Heer in Dana)
Dakota Group; Nebraska, U.S.A.
- ? *rhomboidea*, Lesquereux, 1892, p. 113, pl. li, fig. 10.
Dakota Group; Kansas, U.S.A.
- Butomites cretaceus*, Velenovsky, 1889, p. 25, pl. iii, figs. 10-13, 15.
Cenomanian; Bohemia.
- Caesalpinia Cookiana*, Hollick in Newberry, 1895, p. 94, pl. xlii, figs. 49, 50.
Amboy Clay; New Jersey, U.S.A.
- *raritanensis*, Berry, 1909, p. 257.
Raritan Formation; New Jersey, U.S.A.
- Calamitopsis Konigi*, von der Marck, 1884, p. 81, pl. xiii, fig. 12.
(= *Frenelopsis Konigii*, Hosius & v. d. Marck, 1880.)
Senonian; Drensteinfurth, Westphalia.
- Callistemon cretaceum*, Velenovsky, 1889, p. 22, pl. i, fig. 3.
Cenomanian; Vyserovic, Bohemia.
- Callistemophyllum ambiguum*, Ettingshausen in Reuss, 1854, p. 740 [nomen nudum].
Cenomanian; Moletain, Moravia.
- *Bruderi*, Engelhardt, 1892, p. 115, pl. i, figs. 6, 7.
Cretaceous; Bohemia.
- *Heerii*, Ettingshausen, 1867 a, p. 261, pl. iii, fig. 13.
Cenomanian; Niederschoena, Saxony.
- Callitris? sp.*, Fontaine, 1889, p. 272, pl. clxix, fig. 8.
Potomac Formation; Virginia, U.S.A.
- Calycites alatus*, Hollick, 1905 c, p. 417, pl. lxxii, fig. 8.
Cretaceous Shale; Long Island, U.S.A.
- *diospyriformis*, Newberry, 1895, p. 132, pl. xlii, figs. 39-41.
Amboy Clay; Woodbridge, U.S.A.
- *hexaphylla*, Lesquereux, 1873, p. 402.
Laramie Formation; Wyoming, U.S.A.
- *cbovatus*, Hollick, 1906 a, p. 109, pl. v, fig. 24.
Middle Cretaceous; Martha's Vineyard, U.S.A.

- Calycites parvus**, Newberry, 1895, p. 131, pl. xli, figs. 28-29.
Amboy Clay; Woodbridge, U.S.A.
- **sp.**, Lesquereux, 1892, p. 221, pl. xxii, fig. 8.
Dakota Group; Kansas, U.S.A.
- Campteroneura paradoxa**, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- Camptoneura truncata**, Debey, 1849, p. 299 [nomen nudum]. Ibid.
- Camptopteris biloba**, Presl in Sternberg, 1838, p. 168, in Sternberg,
1825, pl. xlii, fig. 2. Quadersandstein; Hór, Saxony.
- Cannophyllites Hisingeri**, Goeppert, 1848, p. 277 [nomen nudum].
Quadersandstein; Silesia.
- **Nilssoni**, Unger, 1850 A, p. 317. Greensand; Scania, Sweden.
- **septentrionalis**, Nilsson, 1832, p. 346, pl. i, fig. 9. Ibid.
- Capparidoxylon Geinitzi**, Schenk, 1883, p. 12, pl. i, figs. 3-4.
Cretaceous; Cairo, Egypt.
- Cardiopteris tasmanicus**, Johnston, 1896, p. 63, fig. 9.
Cretaceous(?); Tasmania.
- Carex Clarkii**, Berry, 1905 D, p. 347, text-fig. 1.
Middle Cretaceous; Maryland, U.S.A.
- Carlopteris aquensis**, Debey & Ettingshausen, 1859 B, p. 206, pl. iii,
figs. 20-27. Senonian; Aix, Rhenish Prussia.
- **asplenioides**, Debey & Ettingshausen, 1859 B, p. 210, pl. iii,
figs. 29-33; pl. iv, fig. 22. Ibid.
- Carpinites arenaceus**, Goeppert, 1847, p. 306, pl. xxxvii, figs. 5-7.
Cretaceous; Silesia.
- **microphyllus**, Heer, 1883 A, p. 23, pl. lv, fig. 13.
Patoot Beds; Greenland.
- Carpinoxylon compactum**, Vater, 1884, p. 848, pl. xxix, figs. 28, 29.
Lower Senonian (?); Helmstedt.
- Carpites alatus**, Knowlton in Stanton & Hatcher, 1905, p. 150, pl. xvi,
fig. 2. Judith River Beds; Montana, U.S.A.
- **burmanniæformis**, Saporta, 1894, p. 151, pl. xxvii, figs. 9-12.
Urgonian; Portugal.
- **bursæformis**, Lesquereux, 1878 B, p. 306, pl. lx, fig. 30.
Laramie Formation; Wyoming, U.S.A.
- **cocculoides? major**, Lesquereux, 1878 B, p. 307, pl. lx, figs. 38, 39.
Ibid.
- **coniger**, Lesquereux, 1892, p. 221, pl. xxxviii, fig. 17.
Dakota Group; Kansas, U.S.A.
- **cordiformis**, Lesquereux, 1892, p. 220, pl. xxii, fig. 9. Ibid.
- **glumæformis**, Lesquereux, 1878 B, p. 304, pl. xxxv, fig. 4 a; pl. lx,
figs. 14-17. Laramie Formation; Wyoming, U.S.A.
- **granulatus**, Saporta, 1894, p. 209, pl. xxxvi, fig. 17.
Upper Albion; Portugal.
- **judithæ**, Knowlton in Stanton & Hatcher, 1905, p. 149, pl. xvi,
fig. 4. Judith River Beds; Montana, U.S.A.
- **laurineus**, Lesquereux, 1878 B, p. 304, pl. lx, figs. 20, 21.
Laramie Formation; Colorado, U.S.A.

- Carpites ligatus*, Lesquereux, 1878 B, p. 307, pl. lx, figs. 36, 36 a.
Laramie Formation; New Mexico.
- *liriophylli*?, Lesquereux, 1883, p. 77, pl. xi, fig. 5.
Dakota Group; Colorado, U.S.A.
- *minutulus*, Lesquereux, 1878 B, p. 305, pl. lx, fig. 25. Tertiary.
Recorded, Berry, 1905, p. 32, from Cretaceous of Cliffwood, U.S.A.
- *mitratus*, Lesquereux, 1878 B, p. 304, pl. lx, figs. 18, 19.
Laramie Formation; Wyoming, U.S.A.
- *myricarum*, Lesquereux, 1878 B, p. 303, pl. lx, figs. 8-11. Ibid.
- *obovatus*, Lesquereux, 1892, p. 221, pl. lxii, fig. 5.
Dakota Group; Kansas, U.S.A.
- *plicicostatus*, Saporta, 1894, p. 151, pl. xxvi, fig. 8.
Urgonian; Portugal.
- *pruni*, Knowlton in Stanton & Hatcher, 1905, p. 149, pl. xv, figs. 6, 6 a.
Judith River Beds; Montana, U.S.A.
- *rhomboidalis*, Lesquereux, 1878 B, p. 306, pl. lx, figs. 28, 29.
Laramie Formation; Colorado, U.S.A.
- *spiralis*, Lesquereux, 1878 B, p. 306, pl. lx, fig. 27.
Laramie Formation; New Mexico.
- *tileaceus* (Heer), Lesquereux, 1892, p. 221, pl. xxii, figs. 6, 7.
Dakota Group; Kansas, U.S.A.
- *triangulosus*, Lesquereux, 1878 B, p. 302, pl. lx, fig. 4; pl. lxii, figs. 19, 20.
Laramie Formation; Wyoming, U.S.A.
- *utahensis*, Lesquereux, 1878 B, p. 305, pl. lx, fig. 22. Ibid.
- *valvatus*, Lesquereux, 1878 B, p. 307, pl. lx, fig. 37. Ibid.
- *verrucosus*, Lesquereux [*non* (Heer) Schimper], 1878 B, p. 305, pl. lx, fig. 23. Ibid.
- *Viburni*, Lesquereux, 1878 B, p. 305, pl. lx, figs. 26, 26 a. Ibid.
- Carpolithus agglomeratus*, Fontaine, 1889, p. 267, pl. cxxxiv, fig. 5.
Potomac Formation; Virginia, U.S.A.
- *arachioides*, Lesquereux, 1873, p. 403.
Laramie Formation; Wyoming, U.S.A.
- *avellaniformis*, Schlotheim, 1820, p. 421; & 1822, pl. xxi, fig. 6.
Senonian; Aix, Rhenish Prussia.
- *barrensis*, Ward, 1890 B, p. 692, pl. clxix, figs. 14, 15.
Cretaceous; Black Hills, U.S.A.
- *brookensis*, Fontaine, 1889, p. 268, pl. cxxxv, figs. 2, 4; pl. cxxxvi, fig. 6; pl. clxvii, fig. 6. Potomac Formation; Virginia, U.S.A.
- *cliffwoodensis*, Berry, 1905 B, p. 100, pl. xlviii, fig. 6.
Matawan Formation; New Jersey, U.S.A.
- *collophylloides*, Ettingshausen, 1893, p. 151 [*nomen nudum*].
Cretaceous; Australia.
- *complanatus*, Ettingshausen, 1893, p. 151; & 1895, p. 53, pl. iv, fig. 23. Ibid.
- *conjugatus*, Fontaine, 1889, p. 267, pl. cxxxiv, fig. 9.
Potomac Formation; Virginia, U.S.A.
- *cretaceus*, Ettingshausen, 1867 A, p. 262, pl. iii, fig. 2.
Cenomanian; Niederschoena, Saxony.

- Carpolithus curvatus*, Fontaine, 1889, p. 269, pl. cxxxv, fig. 17.
Potomac Formation; Virginia, U.S.A.
- *drupæformis*, Hollick, 1898 D, p. 134, pl. xi, figs. 4, 4a.
Upper Cretaceous (Clay Marl); New Jersey, U.S.A.
- *dubius*, Berry, 1905 B, p. 100, pl. xlviii, fig. 7.
Matawan Formation; New Jersey, U.S.A.
- *euonymoides*, Hollick, 1906 A, p. 110, pl. vii, fig. 2.
Middle Cretaceous; Staten Island, U.S.A.
- *euphorbioides*, Goeppert, 1842 B, p. 157, pl. liv, fig. 19.
Senonian; Aix, Rhenish Prussia.
- *fagiformis*, Ettingshausen, 1893, p. 151; & 1895, p. 53, pl. iv, fig. 22.
Cretaceous; Australia.
- *falcatus*, Lesquereux, 1873, p. 398.
Laramie Formation; Wyoming, U.S.A.
- *fasciculatus*, Fontaine (*non* Lesquereux), 1889, p. 265, pl. cxxxiv, fig. 1.
Potomac Formation; Virginia, U.S.A.
- *floribundus*, Newberry, 1895, p. 133, pl. xlv, figs. 17-21.
Amboy Clay; Woodbridge, U.S.A.
- *foenarius*, Ward, 1899 B, p. 693, pl. clxix, figs. 17, 18.
Lower Cretaceous; Black Hills, U.S.A.
- *geminatus*, Fontaine, 1889, p. 267, pl. cxxxiv, fig. 10.
Potomac Formation; Virginia, U.S.A.
- *Harveyi*, Fontaine, 1894, p. 278, pl. xliii, fig. 3.
Trinity Division; Texas, U.S.A.
- *hirsutus*, Newberry, 1895, p. 134, pl. xlv, figs. 14, 14a.
Amboy Clay; Woodbridge, U.S.A.
- *horridus*, Dawson, 1883, p. 21, pl. i, fig. 3.
Upper Cretaceous; North-West Territory, Canada.
- *hymenospermoides*, Ettingshausen, 1893, p. 151 [*nomen nudum*].
Cretaceous; Australia.
- *juglandiformis*, Berry (*non* Schlotheim), 1905 B, p. 100, pl. xlv, fig. 8.
Matawan Formation; New Jersey, U.S.A.
- *juglandiformis*, Schlotheim, 1822, p. 97, pl. xxii, fig. 5.
Senonian; Aix, Rhenish Prussia.
- *komensis*, Heer, 1882, p. 17, pl. iii, figs. 12, 12b.
Kome Beds; Greenland.
- *latus*, Fontaine, 1889, p. 269, pl. cxxxv, fig. 3.
Potomac Formation; Virginia, U.S.A.
- *longipes*, Heer (*non* Massalongo), 1883 A, p. 45, pl. lix, figs. 8-9.
Patoot Beds; Greenland.
- *mattewanensis*, Berry, 1905 A, p. 48, pl. ii, fig. 7.
Cretaceous; New Jersey, U.S.A.
- (*Zamites*) *meridionalis*, Dawson, 1894, p. 64, pl. x, fig. 42.
Upper Cretaceous; Vancouver Island, Canada.
- *montium-nigrorum*, Ward, 1899 B, p. 692, pl. clxix, fig. 13.
Lower Cretaceous; Black Hills, U.S.A.
- *mucronatus*, Fontaine, 1889, p. 270, pl. cxxxvi, fig. 15.
Potomac Formation; Virginia, U.S.A.

- Carpolithus oblongus*, Goepfert, 1842 B, p. 158, pl. liv, fig. 20.
Senonian; Aix, Rhenish Prussia.
- *obovatus*, Fontaine, 1894, p. 278, pl. xliii, fig. 5.
Trinity Division; Texas, U.S.A.
- *ostryæformis*, Berry, 1905 A, p. 47 (re-naming *Carpolithus virginienensis*, Berry, 1903, *non* Fontaine).
Cretaceous; New Jersey, U.S.A.
- *ovæformis*, Newberry, 1895, p. 134, pl. xlv, figs. 15, 16.
Amboy Clay; Woodbridge, U.S.A.
- *patootensis*, Heer, 1883 A, p. 46, pl. lxiv, fig. 13.
Patoot Beds; Greenland.
- *pruniformis*, Newberry (*non* Heer *nec* Schlotheim), 1895, p. 133, pl. xlv, fig. 42.
New Jersey?, U.S.A.
- *pruniformis*, Schlotheim (*non* Heer), 1822, p. 97, pl. xxi, fig. 4.
Senonian; Aix, Rhenish Prussia.
- *sarthacensis*, Crié, 1877, p. 232 [nomen nudum].
Cretaceous; Mans, France.
- *scrobiculatus*, Heer, 1874 A, p. 120, pl. xxvi, figs. 15, 16.
Atane Beds; Greenland.
- *semisulcatus*, Ettingshausen, 1893, p. 251; & 1895, p. 53, pl. iv, fig. 21.
Cretaceous; Australia.
- *sessilis*, Fontaine, 1889, p. 269, pl. cxxxvi, fig. 9.
Potomac Formation; Virginia, U.S.A.
- *siliculæformis*, Ettingshausen, 1893, p. 151; & 1895, p. 53, pl. iv, fig. 20.
Cretaceous; Australia.
- *Smithiæ*, Mantell, 1844, p. 196; & 1846, p. 53, pl. ii, figs. 3, *a*, *b*, *c*, *d*.
Upper Chalk; Kent.
- *spinosus*, Newberry, 1883, p. 514; & 1898, p. 138, pl. lxxviii, figs. 2, 3.
Upper Cretaceous (?); New Mexico, U.S.A.
- *striatus*, Newberry in Reynolds, 1869, p. 170 [nomen nudum].
Upper Cretaceous (?); Missouri, U.S.A.
- *ternatus*, Fontaine, 1889, p. 265, pl. cxxxiv, figs. 2-4, 6, 8.
Potomac Formation; Virginia, U.S.A.
- *thulensis*, Heer, 1874 A, p. 89, pl. i, fig. 1 *c*.
Kome Beds; Greenland.
- *vaccinioides*, Hollick, 1906 A, p. 110, pl. vii, figs. 19, 19 *a*.
Middle Cretaceous; Staten Island, U.S.A.
- *virginiensis*, Fontaine, 1889, p. 266, pl. cxxxiv, figs. 11-14; pl. cxxxv, figs. 1, 5; pl. clxviii, fig. 7.
Potomac Formation; Virginia, U.S.A.
- *virginiensis*, Fontaine in Berry, 1903 E, p. 100, pl. xlviii, fig. 5.
(Re-named *Carpolithus ostryæformis*, Berry, 1905 A.)
- *vyserovicensis*, Bayer, 1899, p. 47, text-fig. 15.
Perucer Beds; Vyserovic, Bohemia.
- *woodbridgensis*, Newberry, 1895, p. 133, pl. xlv, fig. 22.
Amboy Clay; Woodbridge, U.S.A.
- *sp.*, Dawson, 1893, p. 90, text-fig. 15.
Upper Cretaceous; North-West Territory, Canada.

- Carpolithus* sp., Heer, 1880, p. 8, pl. ii, figs. 10, 11.
Cretaceous; Greenland.
- ? sp., Lesquereux, 1874, p. 114, pl. xxvii, fig. 5; pl. xxx, fig. 11.
Cretaceous; Nebraska, U.S.A.
- sp., Otto, 1854, p. 33, pl. iv, figs. 5, 6. Dippoldswalde, Saxony.
- Cassia angusta*, Heer, 1882, p. 101, pl. xxvii, fig. 6. (= *Palæocassia angustifolia*, Ettingshausen, 1867.) Atane Beds; Greenland.
- *antiquorum*, Heer, 1882, p. 101, pl. xxvii, fig. 5. (= *Leguminosites cassiaeformis*, Heer, 1874 A.) Atane Beds; Greenland.
- *atavia*, Velenovsky, 1887, p. 6, pl. viii, figs. 3-7, 10.
Cenomanian; Kieslingswalde.
- *Etheridgei*, Ettingshausen, 1893, p. 151; & 1895, p. 51, pl. iv, fig. 16. Cretaceous; Australia.
- *Ettingshauseni*, Heer, 1882, p. 100, pl. xxvi, fig. 8.
Atane Beds; Greenland.
- *melanophylla*, Velenovsky, 1887, p. 5, pl. viii, figs. 1, 2, 8, 9, 11, 12. Perucer Beds; Bohemia.
- *phaseolitoides*, Ettingshausen, 1893, p. 151 [nomen nudum].
Cretaceous; Australia.
- *polita*, Lesquereux, 1892, p. 146. Dakota Group; Kansas, U.S.A.
- *præ-mennonina*, Ettingshausen, 1893, p. 151; & 1895, p. 51, pl. iv, fig. 15. Cretaceous; Australia.
- *præ-phaseolitoides*, Ettingshausen, 1895, p. 52, pl. iv, fig. 14. (= *Palæocassia phaseolitoides*, Ettingshausen, 1887 A.) Ibid.
- *problematica*, Lesquereux, 1892, p. 146, pl. xxxviii, fig. 3.
Dakota Group; Kansas, U.S.A.
- sp., Hollick, 1906, p. 84, pl. xxxii, fig. 13.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Castalia* ? *duttoniana*, Knowlton, 1900 A, p. 55, pl. xiii, fig. 7.
Montana Formation; Wyoming, U.S.A.
- *Stantoni*, Knowlton in Stanton & Hatcher, 1905, p. 147, pl. xix, fig. 4. Judith River Beds; Montana, U.S.A.
- Castanea Hausmanni*, Dunker, 1856, p. 181, pl. xxxiv, fig. 1.
Quadersandstein; Blankenburg, Saxony.
- sp., Dawson, 1887, p. 27, pl. i, fig. 8. Souris River, Canada.
- Casuarina Covillei*, Ward, 1895 A, p. 353, pl. iii, fig. 2.
Potomac Formation; Virginia, U.S.A.
- *primæva*, Ettingshausen, 1893, p. 139; & 1895, p. 14, pl. i, figs. 16-20. Cretaceous; Australia.
- Casuarinites cretaceus*, Ettingshausen, 1887 A, p. 181, pl. viii, figs. 6-7. Upper Cretaceous; New Zealand.
- Caudex spinosus*, Lesquereux, 1883, p. 91. (= *Caulinites spinosa*, Lesquereux, 1873.) Dakota Group; Kansas, U.S.A.
- Caulerpa Lehmanni*, Heer, 1877, p. 141, pl. lvii, fig. 11.
Neocomian; Switzerland.
- sp., Lorenz, 1902, p. 23 (56), pl. vii, fig. 3.
Lower Cretaceous; Switzerland.

- Caulerpides bryoides.** (= *Caulerpites bryodes*, Debey & Ettingshausen.)
- Caulerpites articulatus**, Hampe in Stiehler, 1858, p. 56, fig. 3 [nomen nudum]. (= *Delessertites Hampeanus*, Stiehler, 1858.)
- **Brardii**, Sternberg, 1833, p. 23. (= *Fucoides Brardii*, Brongniart, 1828.)
- **Brongniartii**, Debey in Debey & Ettingshausen, 1859 A, p. 203.
Cretaceous; La Rochelle.
- **bryodes**, Debey & Ettingshausen, 1859 A, p. 189, pl. i.
Senonian; Aix, Rhenish Prussia.
- **Diesingii**, Unger, 1850, p. 8; & 1853, p. 77, pl. xxiv, fig. 2.
Cretaceous; Switzerland.
- **Eseri**, Unger, 1850, p. 8; & 1853, p. 77, pl. xxiv, fig. 1.
Cretaceous; Bavaria.
- **incrassatus** (Lesquereux), Knowlton, 1898, p. 61 (re-naming *Delesseria incrassata*, Lesquereux, 1873).
Laramie Formation; New Mexico, U.S.A.
- **lingulatus** (Lesquereux), Knowlton, 1898, p. 61 (re-naming *Delesseria lingulata*, Lesquereux, 1873).
Laramie Formation; New Mexico, U.S.A.
- **montalbanus**, Renger in Fritsch, 1877, p. 72 [nomen nudum].
Cretaceous; Bohemia.
- **Orbignianus** (Brongniart), Sternberg, 1833, p. 23. (= *Fucoides orbignianus*, Brongniart, 1824.)
Lower Cretaceous; Austria and France.
- **tenuis**, Fischer-Ooster, 1858, p. 32, pl. vii, fig. 5.
Upper Cretaceous; Switzerland.
- Caulinites atavinus**, Heer, 1881, p. 22, pl. xix, fig. 4; pl. xvi, fig. 8b.
Cretaceous; Portugal.
- **fecunda**, Lesquereux, 1873, p. 384.
Laramie Formation; Colorado, U.S.A.
- **fimbriatus**, Saporta, 1894, p. 97, pl. xvii, figs. 16-17.
Valanginian; Portugal.
- **inquirendus**, Hollick, 1905 c, p. 406, pl. lxx, fig. 3. (= *Onoclea inquirenda*, Hollick, 1906 A.) Cretaceous Clay; Long Island, U.S.A.
- **sparganicoides**, Lesquereux, 1873, p. 391.
Laramie Formation; Wyoming, U.S.A.
- **spinosa**, Lesquereux, 1873, p. 422. (= *Caudex spinosus*, Lesquereux, 1883.)
Dakota Group; Kansas, U.S.A.
- **stigmarioides**, Ettingshausen, 1867 A, p. 248, pl. ii, fig. 1.
(= *Corticites stigmarioides*, Engelhardt, 1892.)
Cenomanian; Niederschoena, Saxony.
- Caulomorpha Heeri**, Saporta, 1894, p. 217, pl. xxxviii, fig. 9.
Cenomanian; Portugal.
- Caulopteris Brownii**, Renault, 1883, p. 73, pl. viii, fig. 10. (Plate descript. gives name *C. Cottaeana*.) Cretaceous; England.
- **Cottaeana**, Renault, 1883, p. 73 (?), pl. viii, fig. 10. (Text descript. gives name *C. Brownii*.) Cretaceous; France(?).

- Caulopteris cretacea**, Stenzel, 1897, p. 17. (= *Protopteris punctata*, Sternberg, *pro parte*.) Senonian; Westphalia.
- **cyatheoides**, Unger, 1867, p. 649, pl. i, figs. 1-3. Neocomian; Austria.
- **Dicksonioides**, Carruthers, 1865, pl. xiii (misprint for *C. punctata*). Neocomian; Austria.
- **oppoliensis** (Goeppert), Stenzel, 1897, p. 18. (= *Rhizodendron oppoliense*, Goeppert, 1886.) Cretaceous; Oppeln.
- **punctata**, Goeppert, 1836, p. 449. (= *Protopteris punctata*, Presl in Sternberg, 1838.) Cretaceous; Bohemia.
- **Songeri**, Goeppert, 1836, p. 449, pl. xli, fig. 112. (= *Protopteris Songeri*, Presl in Sternberg, 1838.) Cretaceous; Bohemia.
- Ceanothus constrictus**, Hollick, 1906 A, p. 93, pl. xxxiv, figs. 15-17. Middle Cretaceous; Martha's Vineyard, U.S.A.
- **cretaceus**, Dawson, 1883, p. 28, pl. viii, fig. 33. Upper Cretaceous; Vancouver Island, Canada.
- **prodromus**, Heer, 1883 A, p. 41, pl. lxii, fig. 15. Patoot Beds; Greenland.
- **sp. ?**, Hosius & von der Marck, 1880, p. 175, pl. xxxvi, fig. 132. Upper Senonian; Haldem, Westphalia.
- Cedrelospermis venulosus**, Saporta, 1894, p. 98, pl. xvi, fig. 21. Valanginian; Portugal.
- Cedrophloios Bleicheri**, Fliche, 1896, pp. 256-258, pl. xii, fig. 4; pl. xiv, fig. 3. Albian; Vilotte, France.
- Cedroxylon barremianum**, Fliche, 1900, p. 24, pl. ii, fig. 1. Lower Cretaceous; Haute-Marne, France.
- **blevillense**, Lignier, 1907, p. 267, pl. xviii, figs. 15-17; pl. xxi, fig. 66 & diagr.; pl. xxii, fig. 72; pl. xxiii, fig. 87. Albian; Hève, France.
- **Drewi**, Shirley, 1898, p. 13. Upper Cretaceous; Queensland.
- **Gardonienae**, Crié, 1890, p. 236, pl. ii, figs. 1-4. Cenomanian; France.
- **Hoheneggeri**, Felix . . . recorded Liebus, 1901, p. 17, in Perucor Beds of Bohemia.
- **inæquale**, Reiss, 1907, p. 96, fig. 1 of unnumb. pl. Upper Cretaceous; Japan.
- **manehildense**, Fliche, 1896, p. 247, pl. xc, fig. 3. Cenomanian; Sainte-Menehould, France.
- **Matsumuræ**, Stöpes & Fujii, 1910, p. 42, pl. i, fig. 10; pl. iv, figs. 20-23. Upper Cretaceous; Hokkaido, Japan.
- **reticulatum**, Saporta, 1880 (1877), p. 653, pl. iii, fig. 3. Neocomian; France.
- **Ryedalense**, Conwentz, 1892, p. 25, pl. vii, figs. 5-9; pl. viii, fig. 1. Senonian; Sweden.
- **Yendoi**, Stöpes & Fujii, 1910, p. 44, pl. iv, figs. 24-26. Upper Cretaceous; Hokkaido, Japan.
- **sp.**, Schenk, 1869, p. 239, pl. iv, figs. 8-9; pl. vii, fig. 2. Wernsdorfer Beds; Austrian Silesia.

- Cedrus Lennieri*, Saporta, 1880 (1877), p. 652, pl. iv, fig. 1.
Upper Neocomian; Havre, France.
- *lotharingica*, Cornuel, 1882, p. 262, pl. vii, figs. 2-3.
Albian; Houppette, France.
- *oblonga* (Lindley & Hutton), Fliche, 1896, p. 200, pl. viii. (= *Abies oblonga*, Lindley & Hutton.)
Albian; France.
- Celastrophyllum acutidens*, Fontaine, 1889, p. 305, pl. clvi, fig. 8.
Potomac Formation; Virginia, U.S.A.
- *albædomus*, Ward, 1905, p. 489, pl. cviii, fig. 3. Ibid.
- *angustifolium*, Newberry, 1895, p. 100, pl. xiv, figs. 8-17.
Amboy Clay; Woodbridge, U.S.A.
- *arcinerve*, Fontaine, 1889, p. 304, pl. cxlvi, fig. 3.
Potomac Formation; Virginia, U.S.A.
- *australe*, Ettingshausen, 1887 A, p. 188, pl. ix, fig. 14.
Upper Cretaceous; New Zealand.
- *Benedeni*, Saporta & Marion, 1873, p. 67, pl. x, fig. 6; pl. xii, figs. 1-2.
Eocene. (Recorded American Cretaceous. Hollick, 1894 A, p. 58.)
- *Brittonianum*, Hollick in Newberry, 1895, p. 105, pl. xliii, figs. 37, 38, 46, 47.
Amboy Clay; New Jersey (?).
- *brookense*, Fontaine, 1889, p. 305, pl. clviii, fig. 8; pl. clxix, fig. 7.
Potomac Formation; Virginia, U.S.A.
- *crassipes*, Lesquereux, 1892, p. 174, pl. lvii, figs. 6, 7.
Dakota Group; Kansas, U.S.A.
- *crenatum*, Heer, 1883 A, p. 41, pl. lxii, fig. 21.
Patoot Beds; Greenland.
- *cretaceum*, Lesquereux, 1892, p. 173, pl. xxxviii, figs. 12-14.
Dakota Group; Kansas, U.S.A.
- *decurrens*, Lesquereux, 1892, p. 172, pl. xxxvi, fig. 1. Ibid.
- *denticulatum*, Fontaine, 1889, p. 306, pl. clxix, fig. 10; pl. clxxii, fig. 7.
Potomac Formation; Maryland, U.S.A.
- *elegans*, Berry, 1905 E, p. 84, pl. xliii, fig. 6.
Matawan Formation; New Jersey, U.S.A.
- *ensifolium*, Lesquereux, 1874, p. 108, pl. xxi, figs. 2-3.
Dakota Group; Kansas, U.S.A.
- *grandifolium*, Newberry, 1895, p. 104, pl. xix, fig. 8; pl. xxi, figs. 1-4.
Amboy Clay; New Jersey, U.S.A.
- *Hunteri*, Ward, 1895 A, p. 358, pl. iv, fig. 9.
Potomac Formation; Virginia, U.S.A.
- *integrifolium*, Ettingshausen, 1867 A, p. 260, pl. iii, fig. 14.
Cenomanian; Niederschoena, Saxony.
- *lanceolatum*, Ettingshausen, 1867 A, p. 280, pl. iii, fig. 9. Ibid.
- *latifolium*, Fontaine, 1889, p. 306, pl. clxxii, figs. 3, 6; pl. clxxiii, fig. 13.
Potomac Formation; Maryland, U.S.A.
- ? *marylandicum*, Fontaine in Ward, 1905, p. 559, pl. cxvi, fig. 7. Ibid.
- *minus*, Hollick in Newberry, 1895, p. 105, pl. xlii, figs. 51, 52.
Amboy Clay; New Jersey, U.S.A.

- Celastrophyllum myrsinoides**, Lesquereux, 1892, p. 174, pl. lviii, figs. 8, 9. Dakota Group; Kansas, U.S.A.
- **Newberryanum**, Hollick in Newberry, 1895, p. 101, pl. xlix, figs. 1-27. Amboy Clay; Sayreville, U.S.A.
- **obliquum**, Knowlton in Lesquereux, 1892, p. 173, pl. lvii, fig. 5. Dakota Group; Kansas, U.S.A.
- **obovatum**, Fontaine, 1889, p. 307, pl. clxxii, figs. 9, 10. Potomac Formation; Maryland, U.S.A.
- **obtusidens**, Fontaine, 1889, p. 305, pl. clvi, fig. 5. Potomac Formation; Virginia, U.S.A.
- **obtusum**, Heer, 1882, p. 96, pl. xxviii, fig. 15. Atane Beds; Greenland.
- **proteoides**, Fontaine, 1889, p. 304, pl. clxvi, fig. 5. Potomac Formation; Virginia, U.S.A.
- **pulchrum**, Ward, 1899, p. 706, pl. clxxi, figs. 3, 4. Dakota Group; Black Hills, U.S.A.
- **robustum**, Newberry, 1895, p. 103, pl. xlii, figs. 41, 42. Amboy Clay; New Jersey, U.S.A.
- **? saliciforme**, Ward, 1905, p. 494, pl. cviii, fig. 7. Older Potomac Formation; Virginia, U.S.A.
- **serratum**, Saporta & Marion, recorded Heer, 1883 A, p. 41, pl. lxx, fig. 6. Patoot Beds; Greenland.
- **spatulatum**, Newberry, 1895, p. 103, pl. xlii, figs. 43-45. Amboy Clay; New Jersey, U.S.A.
- **tenuinerve**, Fontaine, 1889, p. 306, pl. clxxii, fig. 2. Potomac Formation; Maryland, U.S.A.
- **undulatum**, Newberry, 1895, p. 102, pl. xxxviii, figs. 1-3. Amboy Clay; New Jersey, U.S.A.
- Celastrus arctica**, Heer, 1883 A, p. 40, pl. lxi, figs. 5 d, 5 e. Patoot Beds; Greenland.
- **? sp.**, Hollick, 1893, p. 38, pl. i, fig. 4. Raritan Formation; Staten Island, U.S.A.
- Celtidophyllum cretaceum**, Krasser, 1896, pl. xvi, figs. 8, 9, 10. Cenomanian; Kunstadt, Moravia.
- **cretaceum integrifolia**, Krasser, 1896, p. 116, pl. xvi, figs. 12-14. Ibid.
- **præaustrale**, Krasser, 1896, p. 130, pl. xvi, figs. 8-14. Ibid.
- Celtiophyllum cretaceum**, Krasser, 1889, p. 34 [nomen nudum]. Ibid.
- Celtis ? ovata**, Lesquereux, 1874, p. 66, pl. iv, figs. 2-3. (= *Populites ovata*, Lesquereux, 1868, & *Ampelophyllum ovatum*, Lesquereux, 1876 B.) Dakota Group; Nebraska, U.S.A.
- Cephalotaxites insignis**, Heer, 1883 A, p. 10, pl. liii, fig. 12. Patoot Beds; Greenland.
- Cephalotaxopsis brevifolia**, Fontaine, 1889, p. 238, pl. lv, fig. 3; pl. cvi, fig. 5; pl. cvii, fig. 5. Potomac Formation; Virginia, U.S.A.

- Cephalotaxopsis magnifolia*, Fontaine, 1889, p. 236, pl. civ, figs. 4, 5; pl. cv, figs. 1, 2, 4; pl. evi, figs. 1, 3; pl. evii, figs. 1, 2, 4; pl. eviii, figs. 1, 3, 4. Potomac Formation; Virginia, U.S.A.
- *microphylla*, Fontaine, 1889, p. 238, pl. cviii, fig. 5; pl. cix, fig. 9. Ibid.
- *ramosa*, Fontaine, 1889, p. 237, pl. civ, figs. 2, 3; pl. evi, figs. 2, 4; pl. evii, fig. 3; pl. eviii, fig. 2. Ibid.
- ? *rhytidodes*, Ward, 1905, p. 258, pl. lxviii, fig. 8. Shasta Formation; California, U.S.A.
- *sp.*, Dawson, 1893, p. 89. Kootanie Formation; British Columbia.
- *sp.*, Fontaine in Diller & Stanton, 1894, p. 450. Knoxville Beds; California, U.S.A.
- Cephalotaxospermum carolinianum*, Berry, 1910 A, p. 187. Middle Cretaceous; North Carolina, U.S.A.
- Cerasus meridionalis*, Eichwald, 1865, p. 67, pl. iii, fig. 23. Russia.
- Ceratopetalum primigenium*, Ettingshausen, 1893, p. 138, & p. 150; & 1895, p. 44, pl. iv, fig. 5. Cretaceous; Australia.
- *rivulare*, Ettingshausen, 1887 A, p. 186, pl. ix, figs. 15-16. Upper Cretaceous; New Zealand.
- Ceratophyllum australe*, Ettingshausen, 1893, p. 147; & 1895, p. 14, pl. i, figs. 14-15. Cretaceous; Australia.
- *rivulare*, Crié, 1889, p. 9 [nomen nudum]. Cretaceous; New Zealand.
- Ceratostrobos echinatus*, Velenovsky, 1885, p. 25, pl. xi, figs. 7-15; pl. xiii, fig. 5. Perucer Beds; Bohemia.
- *formosus* (Heer), Schulze, 1888, p. 18. (= *Geinitzia formosa*, Heer.) Senonian; Altenburg.
- *sequoiæphyllum*, Velenovsky, 1885, p. 24, pl. xii, figs. 14-16. Perucer Beds; Bohemia.
- *strictus*, Schulze, 1888, p. 19 [nomen nudum]. (= *Geinitzia cretacea*, Schimper.) Senonian; Altenburg.
- Cercis eocenica*, Lesquereux, 1873, p. 384. Laramie Formation (?); Colorado, U.S.A.
- Cercospora corioccoccum*, Bayer, 1899, pp. 3, 4, text-figs. 1, 1 a, 2. Perucer Beds; Bohemia.
- Cesalpinites marticensis*, Marion, 1890, p. 1054 [nomen nudum]. Turonian; Martiques, France.
- Chamæcyparites Charonis*, Velenovsky, 1889, p. 12, pl. iii, figs. 3-6; pl. ii, fig. 9. Cenomanian; Bohemia.
- Changarniera dubia*, Suportá, 1894, p. 110, pl. xx, figs. 13-14. Lower Cretaceous (?); Portugal.
- Chara acanthica*, Stache, 1880, p. 199 [nomen nudum]. (= *Kosmogyracanthica*, Stache, 1889.) Upper Cretaceous; Corgnale.
- *carinata*, Stache, 1880, p. 198 [nomen nudum]. (= *Kosmogyrarella carinata*, Stache, 1889.) Liburnian (Upper Cretaceous); Cosina.
- *cingulata*, Stache, 1880, p. 198 [nomen nudum]. (= *Kosmogyracringulata*, Stache, 1889.) Ibid.

- Chara cosinensis*, Stache, 1880, p. 198 [nomen nudum]. (= *Nitella cosinensis*, Stache, 1889.) Liburnian (Upper Cretaceous); Cosina.
- *dolium*, Stache, 1880, p. 199 [nomen nudum]. (= *Cristatella dolium*, Stache, 1889.) Upper Cretaceous (?); Corgnale.
- *guttifera*, Stache, 1880, p. 199 [nomen nudum]. (= *Kosmogyyra guttifera*, Stache, 1889.) Ibid.
- *liburnica*, Stache, 1880, p. 198 [nomen nudum]. (= *Lagynophora liburnica*, Stache, 1889.) Upper Cretaceous (?); Cosina.
- *ornata*, Stache, 1880, p. 199 [nomen nudum]. (= *Kosmogyyra ornata*, Stache, 1889.) Ibid.
- *perarmata*, Stache, 1880, p. 199 [nomen nudum]. (= *Kosmogyyra perarmata*, Stache, 1889.) Upper Cretaceous (?); Corgnale.
- *polonica*, Unger, 1860, p. 3, pl. ii, figs. 1-4. Cretaceous (?); Galicia.
- *robusta*, Stache, 1880, p. 179 [nomen nudum]. (= *Nitella robusta*, Stache, 1889.) Upper Cretaceous; Cosina.
- *Stacheana*, Unger, 1860, p. 3, pl. i, figs. 1-4. Ibid.
- *Stantoni*, Knowlton, 1893 a, p. 141, text-figs. 1-3. Bear River Formation; Wyoming, U.S.A.
- *superba*, Stache, 1880, p. 199 [nomen nudum]. (= *Kosmogyyra superba*, Stache, 1889.) Upper Cretaceous (?); Divacea.
- Cheilanthes Mantelli*, Goepfert, 1836, p. 231. (= *Sphenopteris Mantelli*, Brongniart, 1828.) Tilgate Sands; Sussex.
- Cheirolepis Choffati*, Saporta, 1890 a, p. 814; & 1894, p. 90, pl. xvii, figs. 4-9. Valanginian; Portugal.
- Chiropteris elongata*, Rossmässler, MS., in Cotta, 1836, p. 585. Cenomanian; Niederschoena, Saxony.
- *obtusa*, Rossmässler, MS., in Cotta, 1836, p. 585. Ibid.
- *Reichii*, Bronn, 1837, pl. xxviii, fig. 1. (= *Halyserites Reichii*, Sternberg.) Ibid.
- *spatulata*, Newberry, 1891, p. 199, pl. xiv, figs. 1, 2. (= *Sagenopteris*?, Seward, 1910, p. 431.) Kootanie Formation; Montana, U.S.A.
- *Williamsii*, Newberry, 1891, p. 198, pl. xiv, figs. 10, 11. Kootanie Formation; Montana, U.S.A.
- Choffatia Francheti*, Saporta, 1894, p. 150, pl. xxiv, fig. 8; pl. xxvi, figs. 19-22. Urgonian; Portugal.
- Chondrites æqualis*, Sternberg, 1833, p. 26. (= *Fucoides æqualis*, Brongniart, 1824.)
- *aragonensis*, Saporta, 1891 a, p. 312, pl. cclxxv, fig. 1. Cretaceous; Aragon, France.
- *assimilis*, Saporta, 1891 a, p. 312, pl. cclxxv, fig. 2. Cretaceous; France.
- *bignoriensis*, Mantell, 1854, p. 102, text-fig. 9. Chalk; Sussex.
- *Bosqueti*, Miquel, 1853, p. 54, pl. vi, fig. 4. Senonian; Belgium.
- *bulbosus*, Lesquereux, 1873, p. 373; & 1878, p. 42, pl. i, fig. 14. Laramie Formation; New Mexico.

- Chondrites cretaceus**, Presl in Sternberg, 1838, p. 103, pl. xxxiv, fig. 3. Cretaceous (?); Swabian Alps.
- **dichotomus**, Feistmantel, 1874, p. 266. Perucer Beds; Bohemia.
- **difformis** (Brongniart), Sternberg, 1833, p. 26. (= *Fucoides difformis*, Brongniart, 1824.)
- **divaricatus**, Debey & Ettingshausen (*non* Fischer-Ooster), 1859 A, p. 194, pl. ii, fig. 6 b. Senonian; Aix, Rhenish Prussia.
- **elegans**, Debey & Ettingshausen, 1859 A, p. 195, pl. ii, figs. 9-10. Ibid.
- **fastigiatus**, Sternberg, 1833, p. 25. (= section of *Fucoides Targionii*.) Recorded from Lower and Upper Greensand of Sussex by Mantell, etc.
- **filiciformis**, Lesquereux, 1889, p. 32, pl. xvi, fig. 1. Neocomian; Alaska.
- **flexuosus**, Newberry, 1895, p. 34, pl. i, figs. 1-4. Amboy Clay; New Jersey, U.S.A.
- **furcatus**, Sternberg, 1833, p. 27. (= *Fucoides furcatus*, Brongniart, 1824.)
- **furcillatus**, Roemer (*non* Schmalhausen), 1841, p. 1, pl. i, figs. 1, a, b. Quadersandstein; Saxony.
- **furcillatus lator**, von der Marck, 1864, p. 82, pl. xiii, fig. 14. Upper Senonian; Westphalia.
- **fusiformis**, Fischer-Ooster, 1858, p. 53, pl. iv, fig. 3. Lower Cretaceous; Switzerland.
- **intricatus**, Sternberg, 1833, p. 26, as *Chondrides intricatus*, in Schimper, 1874, pl. iii, figs. 4-9. (= *Fucoides intricatus*, Brongniart.) Upper Cretaceous; Westphalia, etc.
- **jugiformis**, Debey & Ettingshausen, 1859 A, p. 193, pl. i, figs. 8-9. Senonian; Aix, Rhenish Prussia.
- **longissimus**, Gümbel, 1861, p. 567. Turonian; Bavaria.
- **Mantelli**, Geinitz, 1850 A, p. 266. (= *Sphaerococcites Mantelli*, Roemer, 1841.) Quadersandstein; Germany.
- **neocomensis**, Heer, 1877, p. 142, pl. lviii, figs. 1-7. Neocomian; Switzerland.
- **polymorphus**, Hosius & von der Marck, 1880, p. 131, pl. xxiv, fig. 3. Upper Senonian; Westphalia.
- **rectangularis**, Gümbel, 1861, p. 563. Neocomian; Bavaria.
- **recurvus**, Sternberg, 1833, p. 26. (= *Fucoides recurvus*, Brongniart, 1824.)
- **Riemsdyki**, Miquel, 1853, p. 55. Senonian; Belgium.
- **rigidus**, Debey & Ettingshausen, 1859 A, p. 197, pl. iii, figs. 8, 15-17. Senonian; Aix, Rhenish Prussia.
- **robionensis**, Saporta & Marion, 1881, p. 94, text-fig. 30. Neocomian; France.
- **serpentinus**, Heer, 1865, p. 190, text-fig. 99; & 1877, p. 142, pl. lvii, figs. 4, 5; pl. lviii, fig. 8. Neocomian; Switzerland.
- **subcurvatus**, Hosius & von der Marck, 1880, p. 131, pl. xxiv, fig. 4. Upper Senonian; Westphalia.

- Chondrites subintricatus**, Debey & Ettingshausen, 1859 A, p. 196, pl. ii, fig. 8. Senonian; Aix, Rhenish Prussia.
- **subsimplex**, Lesquereux, 1878 B, p. 41, pl. i, fig. 13. Laramie Formation; New Mexico.
- **subverticillatus**, Presl in Sternberg, 1838, p. 104, pl. xxviii, fig. 1; pl. lxx, fig. 34. Upper Cretaceous; Westphalia.
- **Targionii** (Br.), Sternberg, 1838, p. 25, pl. ix, fig. 3. (= *Fuccoides Targionii*, Brongniart, 1828.) Greensand; France.
- **Thierensi** (Bosq.), Hosius & von der Marck, 1880, p. 177, pl. xxxvi, fig. 136 a. (= *Delasserites Thierensi*, Miquel.) Lower Senonian; Westphalia.
- **vagus**, Debey & Ettingshausen, 1859 A, p. 196. Senonian; Aix, Rhenish Prussia.
- **sp.**, Otto, 1854, p. 13, pl. ii, fig. 3. Quadersandstein; Saxony.
- Chondrophyllum cretaceum**, Debey, 1848 A, p. 117. Senonian; Aix, Rhenish Prussia.
- **grandidentatum**, Heer, 1869 A, p. 19, pl. xi, fig. 6. Cenomanian; Moletein, Moravia.
- **hederæforme**, Heer, 1871 A, p. 13, pl. iii, figs. 9, 11 b, 2 d. Senonian; Quedlinburg, Saxony.
- **Nordenskiöldi**, Heer, 1871 A, p. 1183; & 1874 A, p. 114, pl. xxxii, figs. 11, 12; pl. xxx, fig. 4 b. (Re-named *Pistia Nordenskiöldi* by Berry, 1910 A.) Atane Beds; Greenland.
- **obovatum**, Newberry, 1895, p. 118, pl. xlii, figs. 26, 27. Amboy Clay; Woodbridge, U.S.A.
- **orbiculatum**, Heer, 1874 A, p. 115, pl. xxxi, fig. 3 c; pl. xxxii, fig. 13. Atane Beds; Greenland.
- **reticulatum**, Hollick in Newberry, 1895, p. 119, pl. xli, figs. 6, 7. Amboy Clay; New Jersey (?), U.S.A.
- **tricuspe**, Schulze, 1888, p. 458. Cretaceous; Quedlinburg, Saxony.
- Chondrophyton dissectum**, Saporta & Marion, 1885, p. 120, text-fig. 126. Turonian; France.
- **laceratum**, Saporta, 1894, p. 219, pl. xxxviii, figs. 4-5. Cenomanian; Portugal.
- **obscuratum**, Saporta, 1894, p. 220, pl. xxxix, fig. 24. Ibid.
- Chonophyllum cretaceum**, Debey, 1848 A, p. 117 [nomen nudum]. Senonian; Aix, Rhenish Prussia.
- **Velenovskyi**, Engelhardt, 1892 A, p. 89, pl. ii, fig. 15. Quadersandstein; Freiburg, Saxony.
- Cinchona rectinervis**, Velenovsky, 1882 A, p. 213 [nomen nudum]. Cretaceous; Bohemia.
- Cinnamomum affine**, Lesquereux, 1868, p. 206. Laramie Formation; Colorado.
- **canadense**, Dawson, 1886, p. 13, pl. iv, fig. 7. Mill Creek Series; Mill Creek, Canada.

- Cinnamomum crassipetiolatum**, Hollick, 1906 A, p. 74, pl. xxx, figs. 3, 4. Middle Cretaceous; Long Island, U.S.A.
- **ellipsoideum**, Saporta & Marion.
(Recorded Heer, 1883 A, from Patoot Beds of Greenland.)
- **ellipticum**, Knowlton in Weed & Knowlton, 1893, p. 54.
Laramie Formation; Montana, U.S.A.
- **Haastii**, Ettingshausen, 1887 A, p. 185, pl. ix, fig. 11.
Upper Cretaceous; New Zealand.
- **Heerii**, Lesquereux, 1859, p. 361; & 1869, p. 431, pl. xxiii, fig. 12.
(= *Daphnogene Heerii*, Lesquereux, 1876 B.)
Dakota Group; Kansas, U.S.A.
- **intermedium**, Newberry (*non* Ettingshausen), 1895, p. 89, pl. xxix, figs. 1-8, 10. Amboy Clay; Woodbridge, U.S.A.
- **intermedium**, Newberry, 1896, pl. xxix, figs. 1-8, 10.
Amboy Clay; New Jersey, U.S.A.
- **Marioni**, Lesquereux, 1892, p. 106, pl. li, figs. 6, 7.
Dakota Group; Kansas, U.S.A.
- **membranaceum** (Lesquereux), Hollick, 1906 A, p. 75, pl. xxix, figs. 5, 6. Middle Cretaceous; Long Island, U.S.A.
- **personatum**, Bayer, 1896, p. 21, text-fig. 6, & p. 35.
Upper Senonian; Kieslingwalde, Bohemia.
- **primigenium**, Ettingshausen, 1895, p. 24, pl. iii, fig. 1.
Cretaceous; Australia.
- **Scheuchzeri**, Heer, recorded by Lesquereux, 1874, p. 23, pl. xxx, figs. 2, 3. Dakota Group; Kansas, U.S.A.
(Originally described from Tertiary.)
- **sezannense**, Watelet, 1866, p. 175, pl. I, fig. 2.
Tertiary; Sezanne. (Recorded Lesquereux, 1891, from Dakota.)
- ? **Stantoni**, Knowlton, 1900 A, p. 60, pl. xv, fig. 1.
Montana Formation; Utah, U.S.A.
- **sp.**, Hollick, 1906 A, p. 75, pl. xxx, fig. 7.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Cissites acerifolius**, Lesquereux, 1892, p. 163, pl. lviii, fig. 1.
Dakota Group; Kansas, U.S.A.
- **acuminatus**, Lesquereux, 1876 A, p. 396; & 1876 B, p. 353, pl. viii, fig. 1. Ibid.
- **acutiloba**, Hollick, 1895, p. 227, pl. cccxxvii, fig. 3. Ibid.
- **affinis**, Lesquereux, 1876 B, p. 352. (= *Platanus affinis*, Lx. 1874.) Ibid.
- **affinis ampla** (Dn.), Knowlton, 1898, p. 70 (re-naming *Platanus affinis ampla*, Dawson). Mill Creek Series; Mill Creek, Canada.
- **alatus**, Lesquereux, 1892, p. 160, pl. xviii, fig. 6.
Dakota Group; Kansas, U.S.A.
- **Brownii**, Lesquereux, 1892, p. 162, pl. xviii, fig. 11. Ibid.
- **crispus**, Velenovsky, 1887, p. 12 (73), pl. iv (xxvii), fig. 6.
Cenomanian; Bohemia.
- **cyclophylla**, Lesquereux, 1876 A, p. 353. (= *Populites cyclophylla*, Hr.)
Dakota Group; Minnesota, U.S.A.

- Cissites dentato-lobatus*, Lesquereux, 1892, p. 164, pl. lxvi, fig. 4.
Dakota Group; Kansas, U.S.A.
- *formosus*, Heer, 1882, p. 85, pl. xxi, figs. 5-8.
Atane Beds; Greenland.
- *formosus magothiensis*, Berry, 1910 n, p. 25.
Magothy Formation; Maryland, U.S.A.
- *Harkerianus*, Lesquereux, 1876 A, p. 397; & 1883 A, p. 67,
pl. iii, figs. 3, 4. (= *Sassafras harkeriana*, Lx., 1873, & *Sassafras*
(*Araliopsis*) *harkerianum*, Lx., 1874.)
Dakota Group; Kansas, U.S.A.
- *Heerii*, Lesquereux, 1876 A, p. 396; & 1876 B, p. 353, pl. vi, fig. 3.
Ibid.
- *ingens*, Lesquereux, 1892, p. 159, pl. xix, figs. 2, 2 a. Ibid.
- *ingens*, var. *parvifolia*, Lesquereux, 1892, p. 160, pl. lvii, figs. 3, 4.
Ibid.
- *insignis*, Heer in Capellini & Heer, 1867, p. 19, pl. ii, figs. 3, 4.
Dakota Group; Nebraska, U.S.A.
- *lobato-crenata* (Lesquereux), Knowlton in Stanton & Knowlton,
1897, p. 145. Wyoming, U.S.A.
- *obtusilobus*, Lesquereux (*non* Saporta), 1892, p. 161, pl. xxxiii,
fig. 5. Dakota Group; Kansas, U.S.A.
- *obtusilobus*, Saporta (*non* Lesquereux), 1894, p. 190, pl. xxxiv,
figs. 12-13. Albian; Portugal.
- *obtusum*, Lesquereux, 1876 B, p. 354. (= *Sassafras obtusum*,
Lesquereux, 1872.) Dakota Group; Kansas, U.S.A.
- *platanoidea*, Hollick, 1895 D, p. 226, pl. cxxxvii, fig. 2. Ibid.
- *populoides*, Lesquereux, 1892, p. 162, pl. xviii, figs. 12-14. Ibid.
- *salisburiaefolius*, Lesquereux, 1883, p. 66. (= *Populus salisburiae-*
folius, Lx., 1868.) Dakota Group; Kansas, U.S.A.
- *sinuosus*, Saporta, 1894, p. 190, pl. xxxiv, fig. 11. Albian; Portugal.
- *vitifolia*, Velenovsky, 1882, p. 214 [nomen nudum]. (= *Cissus*
vitifolia, Velenovsky.) Cenomanian; Bohemia.
- Cissophyllum exulum*, Velenovsky, 1889, p. 24, pl. vi, figs. 4, 5.
(= *Premnophyllum exulum*, Velenovsky.) Ibid.
- *vitifolium*, Velenovsky, 1882 A, p. 214; & 1889, p. 25. (= *Cissus*
vitifolium, Velenovsky.) Ibid.
- Cissus Browniana*, Lesquereux in Winchell, 1885, p. 77; & 1895,
p. 17, pl. A, fig. 8. Dakota Group; Minnesota, U.S.A.
- *lobato-crenatus*, Lesquereux, 1873, p. 396; & 1878 B, p. 240,
pl. xli, figs. 1-3. Laramie Formation; Wyoming.
- *vitifolia*, Velenovsky, 1886, p. 8 (55), pl. ii (xvii), fig. 6; pl. iii
(xviii), fig. 1. Cenomanian; Bohemia.
- Citrophylum aligerum* (Lesquereux), Berry, 1909, p. 258, pl. xviii A,
figs. 1-8. (= *Ficus aligera*, Lesquereux.)
Raritan Formation; New Jersey, U.S.A.
- Cladophlebis acuta*, Fontaine, 1889, p. 74, pl. v, fig. 7; pl. vii, fig. 6;
pl. x, figs. 6, 7; pl. xi, figs. 7, 8; pl. clxvi, fig. 5.
Potomac Formation; Virginia, U.S.A.

- Cladophlebis acuta**, var. **angustifolia**, Fontaine in Ward, 1905, p. 539, pl. cxiv, fig. 5.
Older Potomac Formation; Maryland, U.S.A.
- **alata**, Fontaine, 1889, p. 77, pl. xix, fig. 5.
Potomac Formation; Virginia, U.S.A.
- **angustifolia**, Newberry, 1891, p. 200, pl. xiv, fig. 8.
Kootanie Formation; Montana, U.S.A.
- **argutidens**, Saporta, 1894, p. 74, pl. xviii, figs. 10-12.
Valanginian; Portugal.
- **brevipennis**, Fontaine, 1889, p. 81, pl. xxxvi, fig. 1.
Potomac Formation; Virginia, U.S.A.
- **Browniana** (Dunker), Seward, 1894 A, p. 99. Wealden.
Recorded from American Lower Cretaceous, see Fontaine in Ward, 1905, p. 226, pl. lxxv, fig. 9, & others.
- **columbiana**, Dawson, 1894, p. 55, pl. v, figs. 4, 5.
Upper Cretaceous; Vancouver Island, Canada.
- **confusior**, Saporta, 1894, p. 166, pl. xxv, fig. 6; pl. xxxi, figs. 8-9.
Albian; Portugal.
- **constricta**, Fontaine, 1889, p. 68, pl. ii, fig. 11; pl. iii, fig. 2; pl. vi, figs. 5, 6, 8-14; pl. xxi, figs. 9, 13; pl. clxix, fig. 2.
Potomac Formation; Virginia and Maryland, U.S.A.
- **crenata**, Fontaine, 1889, p. 75, pl. ix, figs. 7-9; pl. x, figs. 1, 2; pl. xiv, figs. 1-3; pl. xix, fig. 1; pl. xx, fig. 6.
Potomac Formation; Virginia, U.S.A.
- **denticulata** (Heer), Fontaine (*non* Brongniart), 1889, p. 71, pl. iv, fig. 2; pl. vii, fig. 7.
Ibid.
- **derelicta**, Saporta, 1894, p. 77, pl. xv, fig. 2; pl. xvi, fig. 9.
Valanginian; Portugal.
- **distans**, Fontaine, 1889, p. 77, pl. xiii, figs. 4, 5.
Potomac Formation; Virginia, U.S.A.
- **Dunkeri** (Schimper), Seward, 1894 A, p. 100. (= *Pecopteris Dunkeri*, Schimper.)
- **falcata**, Fontaine, 1889, p. 72, pl. iv, fig. 8; pl. v, figs. 1-6; pl. vi, fig. 7; pl. vii, figs. 1, 2.
Potomac Formation; Virginia, U.S.A.
- **falcata montanensis**, Fontaine in Ward, 1905, p. 291, pl. lxxi, figs. 14-20. (Re-naming *Thinnfeldia montanensis*, Fontaine.)
Kootanie Formation; Montana, U.S.A.
- **Fisheri**, Knowlton, 1907, p. 109, pl. xi, figs. 2, 2 A. Ibid.
- **fissipennis**, Saporta, 1894, p. 77, pl. xvi, fig. 5.
Valanginian; Portugal.
- **heterophylla**, Fontaine, 1889, p. 493, pl. lxxxiv, fig. 2.
Kootanie Formation; Montana, U.S.A.
- **inæquiloiba**, Fontaine, 1889, p. 80, pl. xxv, fig. 8.
Potomac Formation; Virginia, U.S.A.
- **inclinata**, Fontaine, 1889, p. 76, pl. x, figs. 3, 4; pl. xx, fig. 8.
Ibid.
- **latifolia**, Fontaine, 1889, p. 69, pl. iii, fig. 1; pl. vi, fig. 4. Ibid.
- **Limai**, Saporta, 1894, p. 165, pl. xxix, fig. 8. Albian; Portugal.

- Cladophlebis minutissima**, Saporta, 1894, p. 77, pl. xv, fig. 26;
pl. xvi, fig. 19 c. Valanginian; Portugal.
- **Nathorsti**, Yokoyama, 1894, p. 220, pl. xxviii, figs. 3, 4, 10, 11.
Neocomian; Japan.
- **oblongifolia**, Fontaine, 1889, p. 74, pl. vii, figs. 3-5.
Potomac Formation; Virginia, U.S.A.
- **obtusiloba**, Saporta, 1894, p. 166, pl. xxix, fig. 9.
Albian; Portugal.
- **pachyphylla**, Fontaine, 1889, p. 80, pl. xxv, fig. 9.
Potomac Formation; Virginia, U.S.A.
- **parva**, Fontaine, 1889, p. 73, pl. iv, fig. 7; pl. vi, figs. 1-3. Ibid.
- **petiolata**, Fontaine, 1889, p. 80, pl. xxii, fig. 8. Ibid.
- **rotundata**, Fontaine, 1889, p. 78, pl. xx, figs. 9-10. Ibid.
- **sinulatiloba**, Saporta, 1894, p. 76, pl. xvi, figs. 1-2.
Valanginian; Portugal.
- **skagitensis**, Penhallow, 1907, p. 306, text-fig. 2.
Cretaceous; British Columbia.
- **sphenopteroides**, Fontaine, 1889, p. 79, pl. xxi, fig. 4.
Potomac Formation; Virginia, U.S.A.
- **sulcycadina**, Saporta, 1890 A, p. 814; & 1894, p. 75, pl. xvii,
fig. 18. Valanginian; Portugal.
- **Unger** (Dunker), Ward, 1905, p. 228, pl. lxx, figs. 15-16.
(= *Pecopteris Unger*, Dunker.)
Shasta Formation; California, U.S.A.
- **virginiensis**, Fontaine, 1889, p. 70, pl. iii, figs. 3-8; pl. iv, figs. 1,
3-6. Potomac Formation; Virginia, U.S.A.
- **wyomingensis**, Fontaine in Ward, 1899 B, p. 656, pl. clx, figs. 16,
17. Lower Cretaceous; Black Hills, U.S.A.
- **sp.**, Dawson, 1893, p. 85.
Kootanie Formation; British N.W. Territory, Canada.
- **sp. ?**, Fontaine, 1889, p. 76, pl. x, figs. 5, 8; pl. xx, fig. 7.
Potomac Formation; Virginia, U.S.A.
- Clathraria ? galtiana**, Hosijs & von der Mark, 1880, p. 202, pl. lxii,
fig. 180. Lower Gault; Westphalia.
- **Lyellii**, Mantell, 1827, p. 52, pl. i, figs. 1, 2, 7; pl. ii, figs. 1, 2, 3;
pl. iii, fig. 8; pl. iii, fig. 4.
Wealden; Sussex. (Recorded from Chalk Marl, Isle of
Wight, Mantell, 1847, pp. 295 & 297 text-fig.)
- **Schacht** (Coemans), Schimper, 1870. (= *Cycadeoidea Schacht*,
Capellini & Solms, 1892.) Gault; Belgium.
- Clathropodium foratum**, Saporta, 1875, p. 297, pl. cxxiv, figs. 1, 2.
(= *Cycadeoidea forata*, Saporta in Capellini & Solms, 1892.)
Recorded from Gault; Havre (first described from Jurassic).
- **sp.**, Yokoyama, 1906, p. 37, pl. xii, fig. 2.
Lower Cretaceous; China.
- Clathropteris egyptiaca**, Seward, 1907, p. 253, text-fig. 1.
Jurassic?; Egypt.
(Shown by Newton 1909 to be Senonian.)

- Coccoloba lævigata*, Lesquereux, 1873, p. 387; & 1878 b, p. 208,
pl. xxxv, fig. 7. Wyoming, U.S.A.
- Cocculites imperfectus*, Hollick, 1906 A, p. 63, pl. xii, fig. 14.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- *inquirendus*, Hollick, 1906 A, p. 63, pl. xii, fig. 13.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Cocculus cercifolius*, Saporta, 1890 b, p. 182 [nomen nudum].
Lower Cretaceous; Bagnols, France.
- *cinnamomeus*, Velenovsky, 1887, p. 4 (65), pl. viii (xxxi), figs. 16-21.
Perucer Beds; Lipence, Bohemia.
- *extinctus*, Velenovsky, 1887, p. 3 (64), pl. vi (xxix), figs. 1, 3.
Cenomanian; Silesia.
- *minutus*, Hollick, 1905 c, p. 407, pl. lxx, fig. 6.
Cretaceous Clay; Long Island, U.S.A.
- *princeps*, Saporta, 1890 b, p. 182 [nomen nudum].
Lower Cretaceous; Bagnols, France.
- Coccolopsis ovata*, Fliche, 1896, p. 273, pl. xiii, figs. 3, 4; pl. xvii, fig. 3,
Lower Cenomanian; Sainte-Menehould, France.
- *Zeilleri*, Fliche, 1896, p. 271, pl. xii, figs. 5, 6; pl. xiii, figs. 1, 2.
Lower Cenomanian; Argers, France.
- Codites idicensis*, Squinabol, 1890, p. 185, pl. viii, fig. 3.
Cretaceous (?); Italy.
- *neocomiensis*, Saporta & Marion, 1881, p. 96, text-fig. 31.
Neocomian; France.
- Colutea coronilloides*, Heer, 1882, p. 100. (= *Leguminosites coronil-*
loides, Heer, 1874.) Atane Beds; Greenland.
- *Langeana*, Heer, 1882, p. 100, pl. xl, fig. 7 b. Ibid.
- *obovata*, Berry, 1906 b, p. 175, pl. viii, figs. 5-6.
Magothy Formation; Maryland, U.S.A.
- *primordialis*, Heer, 1882, p. 99, pl. xxvii, figs. 7-11; pl. xliii,
figs. 7-8. Atane Beds; Greenland.
- *protogæa*, Heer, 1883 A, p. 43, pl. lxi, fig. 1 c; pl. lxii, fig. 1 c.
Patoot Beds; Greenland.
- *valde inæqualis*, Heer, 1882, p. 100, pl. xxvii, figs. 12-13.
Atane Beds; Greenland.
- Combretiphyllum acuminatum*, Menzel, 1909, p. 402, pl. ii, fig. 7.
Senonian; Kamerun.
- Compteroneura paradoxa*, Debey, 1850, p. 117 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- *truncata*, Debey, 1850, p. 117 [nomen nudum]. Ibid.
- Comptonia? antiqua*, Nilsson, 1832, p. 346, pl. i, fig. 8. (= *Dryandra*
antiqua, Ettingshausen, 1851, = *Comptonites antiquus*, Unger, 1865.)
Greensand; Sweden.
- *microphylla* (Heer), Berry, 1906 A (re-naming *Rhus microphylla*,
Heer, 1874 A, & others.)
- *tenera*, Hosius & von der Marck, 1885, p. 227, pl. xix, figs. 3-4.
Upper Senonian; Westphalia.

- Comptoniopteris cercalina**, Saporta, 1894, p. 129, pl. xxvi, fig. 24.
Urgonian; Portugal.
- **intermedia**, Marion, 1890, p. 1053. Turonian; Martigues, France.
- **provincialis**, Marion, 1890, p. 1053. Ibid.
- **Saportæ**, Marion, 1890, p. 1053. Ibid.
- **Vasseuri**, Marion, 1890, p. 1053. Ibid.
- Comptonites antiquus** (Nilsson), Hisinger, 1837, p. 111, pl. xxxiv, fig. 7; & Unger, 1865, p. 374, pl. i, fig. 1. (= *Comptonia antiqua*, Wilson, 1832, = *Dryandra antiqua*, Ettingshausen, 1851.)
Greensand; Sweden.
- Confervites? ægagropiloides**, Brongniart, 1828, p. 36, pl. i, figs. 4, 5.
Cretaceous; Denmark.
- **aquensis**, Debey & Ettingshausen, 1859 A, p. 187, pl. i, fig. 4.
Senonian; Aix, Rhenish Prussia.
- **Brardii** (Sternberg), Debey, 1848 A, p. 114.
Cretaceous; La Rochelle, France.
- **cæspitosus**, Debey & Ettingshausen, 1859 A, p. 188, pl. i, figs. 10-12.
Senonian; Aix, Rhenish Prussia.
- **dubius**, Berry, 1903 c, p. 677, text-fig. 9, p. 678.
Matawan Formation; New Jersey, U.S.A.
- **fasciculatâ**, Brongniart, 1828, p. 35, pl. i, figs. 1-3.
Cretaceous; Denmark.
- **orbigniana** (Sternberg), Debey, 1848 A, p. 114.
Cretaceous; La Rochelle, France.
- **ramosus**, Debey & Ettingshausen, 1859 A, p. 188, pl. i, fig. 9.
Senonian; Aix, Rhenish Prussia.
- **Woodwardii**, Mantell, 1844, p. 104, text-fig. 4. Chalk; Norfolk.
- Ccnifero-caulon colymbæforme**, Fliche, 1900, p. 16.
Lower Cretaceous; France.
- Coniopteris nitidula**, Yokoyama, 1906, p. 35, pl. xii, figs. 4, 4 a.
Cretaceous; China.
- Conites familiaris**, Sternberg, 1825, p. xxxix, pl. xlvii, fig. 2.
(= *Zamitostrobis familiaris*, Bronn & Roemer, 1856.)
Pläner; Třebitz, Bohemia.
- **gibbus**, Reuss, 1844, p. 169. (= *Microzamia gibba* (Reuss), Corda, 1846.)
Ibid.
- **insignis**, Bronn & Roemer, 1856, p. 54, pl. xxviii, fig. 13.
Cenomanian; Niederschoena, Saxony.
- **sp.**, Bronn, 1837, p. 577, pl. xxviii, fig. 13. Ibid.
- Conospermities ellipticus**, Fontaine, 1889, p. 279, pl. cxxxviii, fig. 14.
Potomac Formation; Virginia, U.S.A.
- **hakeæfolius**, Ettingshausen, 1867 A, p. 254, pl. iii, figs. 4, 12.
Cenomanian; Niederschoena, Saxony.
- **hakeæfolius**, Velenovsky (*non* Ettingshausen), 1882 A, p. 213 [nomen nudum].
Cretaceous; Bohemia.
- **linearifolius**, Ettingshausen, 1893, pp. 140, 149; & 1895, p. 27, pl. iii, fig. 15.
Cretaceous; Australia.

- Corallinites galaxacera** (*Galaxaura*), Massalongo, 1856, p. 42,
pl. vi, figs. 1-2. Cretaceous; Venice, Italy.
- Cornophyllum vetustum**, Newberry, 1895, p. 119, pl. xix, fig. 10.
Amboy Clay; Woodbridge, U.S.A.
- Cernoxylon myricæforme**, Vater, 1884, p. 846, pl. xxix, figs. 25, 26.
Lower Senonian (?); Helmstedt
- Cornus cretacea**, Caspary, 1888, p. 39. Königsberg, Prussia.
- **cretacea solidior**, Caspary, 1888, p. 40; 1889, p. 29, pl. vi, figs. 4-17. Ibid.
- **Forchhammeri**, Heer, 1882, p. 85, pl. xlv, fig. 13.
Atane Beds; Greenland.
- **Holmiana**, Heer, 1883 A, p. 36, pl. lxii, fig. 12; pl. lxiv, figs. 6, 7.
Patoot Beds; Greenland.
- **obesus**, Dawson, 1894, p. 62, pl. ix, fig. 30.
Upper Cretaceous; Vancouver Island, Canada.
- **platyphylloides**, Lesquereux, 1892, p. 126, pl. lxiv, fig. 15.
Dakota Group; Kansas, U.S.A.
- **præcox**, Lesquereux, 1892, p. 125, pl. xxiii, fig. 5. Ibid.
- **rhamnifolia**, Weber, 1852, p. 192, pl. xxi, fig. 8. Tertiary.
(Recorded Lesquereux, 1872 & 1878 B. American
Laramie & Montana Formations.)
- **Studerii**?, Heer, 1859. Tertiary.
(Recorded Ward, 1872, p. 293, & 1878, p. 244,
pl. xlii, figs. 4, 5, American Cretaceous.)
- **suborbifera**, Lesquereux, 1878 B, p. 243, pl. xlii, fig. 2.
Laramie Formation; Colorado, U.S.A.
- **thulensis**, Heer, 1883 A, p. 37, pl. lxii, figs. 9-11.
Patoot Beds; Greenland.
- Corticites stigmarioides** (Ettingshausen), Engelhardt, 1892, p. 116.
(= *Caulinites stigmarioides*, Ettingshausen) Cretaceous; Bohemia.
- Costarites undulatus**, Debey, 1848 A, p. 115 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- Cratægus aceroides**, Lesquereux, 1892, p. 143, pl. liv, fig. 8; pl. lv,
fig. 1. Dakota Group; Kansas, U.S.A.
- ? **æquidentata**, Lesquereux, 1878 B, p. 297, pl. lviii, figs. 4, 4 a.
Laramie Formation; Wyoming, U.S.A.
- **atavina**, Heer, 1883 A, p. 43, pl. lxiv, fig. 11.
Patoot Beds; Greenland.
- ? **fragarioides**, Heer, 1883 A, p. 43, pl. lxii, fig. 10 b. Ibid.
- **Lacoei**, Lesquereux, 1892, p. 143, pl. lxiv, fig. 14.
Dakota Group; Kansas, U.S.A.
- **lawrenciana**, Lesquereux, 1892, p. 142, pl. xxxviii, fig. 1. Ibid.
- **monmcuthensis**, Berry, 1907 A, p. 675, text-fig. 1.
Magothy Formation; New Jersey, U.S.A.
- **tenuinervis**, Lesquereux, 1892, p. 142, pl. liv, figs. 5-7.
Dakota Group; Kansas, U.S.A.
- Credneria acerifolia**, Richter, 1905, p. 13, pl. ii, fig. 13; pl. iii, fig. 1.
Quadersandstein; Blankenburg, Saxony.

- Credneria acuminata**, Hampe, 1852, p. 7 [nomen nudum]. (In Stiehler, 1858, p. 64, pl. x, figs. 6, 7.)
Quadersandstein; Blankenburg.
- **arcuata**, Velenovsky, 1882 A, p. 14, pl. viii, fig. 9.
Cenomanian; Vyserovic, Bohemia.
- **arcuata**, Richter (*non* Velenovsky), 1905, p. 14, pl. iii, fig. 4; pl. vi, fig. 10.
Quadersandstein; Blankenburg, Saxony.
- **atava**, Richter, 1905, p. 14, pl. iii, fig. 6.
Upper Cretaceous; Neinstedt, Saxony.
- **Beckeriana**, Goeppert, 1845, p. 220 [nomen nudum].
Cretaceous; Silesia.
- **biloba**, Zenker, 1833, p. 19, pl. ii, figs. A & B.
Quadersandstein; Blankenburg, Saxony.
- **bohémica**, Velenovsky, 1882 A, p. 213; & 1882 B, p. 9, pl. iii, fig. 1; pl. iv, figs. 10-11.
Cenomanian; Vyserovic, Bohemia.
- **cuneifolia**, Bronn, 1837, p. 583, pl. xxviii, fig. 11. (= *Eltingshausenia cuneifolia* of Stiehler, 1858.)
Cenomanian; Niederschoena, Saxony.
- **denticulata**, Zenker, 1833, p. 18, pl. ii, fig. E.
Quadersandstein; Blankenburg, Saxony.
- **elongata**, Richter, 1905, p. 14, pl. ii, fig. 11; pl. iv, figs. 4, 7; pl. vi, figs. 6, 7, 8, 9.
Upper Cretaceous; Eselstall, Saxony.
- **Engelhardtii**, Richter, 1905, p. 14, pl. iv, fig. 3.
Upper Cretaceous; Neinstedt, Saxony.
- **expansa**, Brongniart, 1849 A, p. 111 [nomen nudum]. (= *Eltingshausenia expansa* of Stiehler, 1858.)
Cenomanian; Niederschoena, Saxony.
- **Geinitziana**, Unger, 1849, p. 349. (= *Eltingshausenia Geinitziana*, Stiehler, 1858.)
Ibid.
- **glandulosa**, Richter, 1905, p. 14, pl. ii, fig. 12; pl. iv, fig. 1.
Upper Cretaceous; Eselstall, Saxony.
- **grandidentata**, Unger, 1849, p. 348, pl. v, fig. 5. (= *Eltingshausenia grandidentata* of Stiehler, 1858.)
Cenomanian; Niederschoena, Saxony.
- **integerrima**, Zenker, 1833, p. 17, pl. ii, fig. F.
Quadersandstein; Blankenburg, Saxony.
- **lævis**, Velenovsky, 1882 A, p. 213; & 1882 B, p. 13, pl. iii, fig. 4; pl. iv, figs. 2-6. (= *Platanus lævis*, Velenovsky, 1889.)
Cenomanian; Bohemia.
- **Le Conteana**, Lesquereux, 1868, p. 98. (= *Protophyllum Leconteana*, Lx., 1874.)
Dakota Group; Nebraska, U.S.A.
- **macrophylla**, Heer, 1869 A, p. 16, pl. iv.
Upper Cretaceous; Moletain, Moravia.
- **? microphylla**, Lesquereux, 1876 A, p. 397.
Dakota Group; Kansas, U.S.A.
- **oblonga**, Schimper, 1874, p. 60.
Quadersandstein; Blankenburg, Saxony.
- **peltata**, Richter, 1905, p. 14, pl. iv, fig. 5; pl. vi, fig. 11.
Upper Cretaceous; Eselstall, Saxony.

- Credneria posthuma**, Richter, 1905, p. 15, pl. iii, fig. 7.
Upper Cretaceous; Westerhausen, Saxony.
- **Reichi**, Geinitz, 1850 A, p. 274 [nomen nudum]. (= *Ettingshausenia Genitziana*, Stiehler, 1858.) Quadersandstein; Saxony.
- **reticulata**, Eichwald, 1853, p. 230, text-fig. 3. (= *Ettingshausenia reticulata*, Stiehler, 1858.) Neocomian; Kursk, Russia.
- **rhomboidea**, Velenovsky, 1882 A, p. 213; 1882 B, p. 11, pl. iii, figs. 2, 3; pl. iv, fig. 1. Cenomanian; Kuchelbad, Bohemia.
- **Schneideriana**, Goeppert, 1845, p. 220 [nomen nudum].
Tiefenfurth, Silesia.
- **spathulata**, Eichwald, 1853, p. 230, text-figs. 5, 6. (= *Ettingshausenia spathulata*, Stiehler, 1858.) Neocomian; Kursk, Russia.
- **Sternbergii**, Brongniart, 1849 A, p. 160 [nomen nudum]. (= *Ettingshausenia Sternbergi*, Stiehler, 1858.)
Cenomanian; Teschen, Bohemia.
- **subserrata**, Hampe, 1852, p. 7 [nomen nudum]. (In Stiehler, 1858, p. 64, pl. ii, fig. 18.)
Quadersandstein; Blankenburg, Saxony.
- **subtriloba**, Zenker, 1833, p. 20, pl. iii, figs. C, D, G. Ibid.
- **superstes**, Velenovsky, 1882 A, p. 213; & 1882 B, p. 15, pl. iv, figs. 7-9. Cenomanian; Leipa, Bohemia.
- **tenuinervis**, Hosius, 1870 A, p. 95, pl. xii, fig. 2.
Senonian; Legden, Westphalia.
- **tremulæfolia**, Brongniart, 1849 A, p. 111 [nomen nudum].
(= *Ettingshausenia tremulæfolia*, Stiehler, 1858.)
Cenomanian; Niederschoena, Saxony.
- **triacuminata**, Hampe, 1852, p. 7 [nomen nudum]. (In Stiehler, 1858, p. 64, pl. x, figs. 8, 9.)
Quadersandstein; Blankenburg, Saxony.
- **venulosa**, Eichwald, 1853, p. 230, text-fig. 4. (= *Ettingshausenia venulosa*, Stiehler, 1858.) Neocomian; Kursk, Russia.
- **westfalica**, Hosius, 1870 A, p. 95, pl. xii, fig. 1.
Senonian; Legden, Westphalia.
- **Zenkeri**, Richter, 1905, p. 12. Senonian; Quedlinburg, Saxony.
- **Zenkeri acuminata** (Hampe), Richter, 1905, p. 12, pl. ii, fig. 5.
(= *Credneria acuminata*, Hampe.)
Quadersandstein; Blankenburg, Saxony.
- **Zenkeri asymmetrica**, Richter, 1905, p. 12, pl. v, fig. 5. Ibid.
- **Zenkeri denticulata** (Zenker), Richter, 1905, p. 13, pl. ii, figs. 6, 7; pl. vi, figs. 1, 12. (= *Credneria denticulata*, Zenker.)
Ibid.
- **Zenkeri integerrima** (Zenker), Richter, 1905, p. 12, pl. i, fig. 1; pl. iii, figs. 3, 5, 8a, 8b; pl. v, figs. 6, 7, 8. (= *Credneria integerrima*, Zenker, 1833.) Ibid.
- **Zenkeri intermedia**, Richter, 1903, p. 13, pl. vi, figs. 2, 3, 4, 5, 13. Ibid.
- **Zenkeri oblonga** (Schimper), Richter, 1905, p. 13. (= *Credneria oblonga*, Schimper.) Ibid.

- Credneria Zenkeri orbicularis**, Richter, 1905, p. 12, pl. ii, fig. 2;
pl. v, figs. 2, 3. Upper Cretaceous; Saxony.
- **Zenkeri subserata** (Hampe), Richter, 1905, p. 13, pl. ii, fig. 8.
Quadersandstein; Blankenburg, Saxony.
- **Zenkeri subtriloba** (Zenker), Richter, 1905, p. 13, pl. ii, fig. 3;
pl. v, figs. 9–11. Senonian; Quedlinburg.
- **Zenkeri triacuminata** (Hampe), Richter, p. 13, pl. ii, fig. 4;
pl. iv, fig. 2; pl. v, figs. (14), 15, 16. (= *Credneria triacuminata*,
Hampe.) Quadersandstein; Blankenburg, Saxony.
- **Zenkeri triloba**, Richter, 1905, p. 12, pl. v, fig. 12.
Upper Cretaceous; Eselstall, Saxony.
- ? **sp.**, Dunker, 1856, p. 182, pl. xxxv, fig. 1.
Quadersandstein; Blankenburg, Saxony.
- **sp.**, Dunker, 1856, p. 180, pl. xxxiii, fig. 3. Ibid.
- **sp.**, Heer, 1874 A, p. 111, pl. xxxii, figs. 20, 21.
Atane Beds; Greenland.
- **sp.**, Otto, 1854, p. 47, pl. ix, figs. 8, 9. Dippoldswalde, Saxony.
- **sp.**, Richter, 1905, p. 14, pl. iii, fig. 2.
Upper Cretaceous; Salzberg, Saxony.
- **sp.**, Schenk, 1890, p. 485, text-fig. 286.
Quadersandstein; Blankenburg, Saxony.
- Cretovarium japcnicum**, Stopes & Fujii, 1910, p. 70, pl. ix, figs. 58–60.
Upper Cretaceous; Hokkaido, Japan.
- Cristatella doliolium**, Stache, 1889, p. 136, pl. iv, fig. 8. (= *Chara*
dolium, Stache, 1880.) Upper Cretaceous; Cornuaille.
- Crossochorda Marionii**, Dewalque, 1880, p. 45, pl. ii, fig. 1.
Cretaceous; Condroz, Belgium.
- Crotonophyllum cretaceum**, Velenovsky, 1889, p. 20, pl. v, figs. 4–11.
Cenomanian; Vyserovic, Bohemia.
- Cryptomeria primæva**, Corda in Reuss, 1846, p. 89, pl. xlviii, figs. 1–11.
Pläner; Bohemia.
- Cryptomeriopsis antiqua**, Stopes & Fujii, 1910, p. 52, pl. i, fig. 11;
pl. vi, figs. 35–41; text-fig. 18.
Upper Cretaceous; Hokkaido, Japan.
- **mesozoica**, Suzuki, 1910, p. 185, pl. vii, figs. 3–5. Ibid.
- Cryptomerites hungaricus**, Tuzson, 1908, p. 3, pl. ii, figs. 4, 5.
Upper Cretaceous; Hungary.
- Ctenidium dentatum**, Heer, 1881, p. 17, pl. xvi, figs. 12–15.
Cretaceous; Almargem, Portugal.
- **integerrimum**, Heer, 1881, p. 17, pl. xvi, figs. 4–11. Ibid.
- Ctenis imbricata**, Fontaine, 1889, p. 177, pl. cxxxviii, figs. 10–12.
Potomac Formation; Virginia, U.S.A.
- Ctenophyllum latifolium**, Fontaine, 1889, p. 175, pl. lxviii, figs. 2, 3.
Ibid.
- Ctenopteris angustifolia**, Fontaine, 1889, p. 159, pl. lxv, fig. 2;
pl. lxvii, fig. 4. Ibid.
- **columbiensis**, Penhallow, 1902 B, p. 26, text-fig. 1.
Upper Cretaceous; Vancouver Island, Canada.

- Ctenopteris insignis*, Fontaine, 1889, p. 156, pl. lxi, figs. 4, 5; pl. lxii, fig. 1; pl. lxiii, figs. 1, 2. Potomac Formation; Virginia, U.S.A.
- *integrifolia*, Fontaine, 1889, p. 158, pl. lxii, fig. 2; pl. lxx, fig. 3. Ibid.
- *longifolia*, Fontaine, 1889, p. 159, pl. lxvii, fig. 5. Ibid.
- *minor*, Fontaine, 1889, p. 157, pl. lxvii, fig. 3. Ibid.
- *ultima*, Saporta, 1894, p. 84, pl. xv, fig. 27. Valanginian; Portugal.
- *virginiensis*, Fontaine, 1889, p. 157, pl. lxii, fig. 4; pl. lxx, fig. 1; pl. lxvi, fig. 4. Potomac Formation; Virginia, U.S.A.
- Culmites cretaceus*, Ettingshausen (*non* Miquel), 1867 A, p. 247, pl. i, fig. 3. Cenomanian; Niederschöna, Saxony.
- *cretaceus*, Miquel (*non* Ettingshausen), 1853, p. 53. Senonian; Belgium.
- *priscus*, Ettingshausen, 1852 c, p. 16, pl. iii, figs. 4-8. Wernsderfer Beds (see Krasser, 1896, p. 146).
- *sp.*, Roemer, 1870, p. 291, pl. xxviii, fig. 14. Cenomanian; Silesia.
- Cunninghamia elegans*, Corda in Reuss, 1846, p. 93, pl. xlix, figs. 29-31. Quadersandstein; Bohemia.
- *planifolia*, Corda in Reuss, 1846, p. 93, pl. l, figs. 1-3. Quadersandstein; Perutz, Bohemia.
- *stenophylla*, Velenovsky, 1885, p. 15, pl. v, figs. 2, 4, 10, 16. Perucer Beds; Bohemia.
- Cunninghamiostrobus yubariensis*, Stopes & Fujii, 1910, p. 45, pl. v, figs. 27-34, text-figs. 15, 16, 17. Upper Cretaceous; Hokkaido, Japan.
- Cunninghamites borealis*, Heer, 1882, p. 55, pl. xxix, fig. 12. Atane Beds; Greenland.
- *dubius*, Sternberg in Unger, 1867, p. 651, pl. ii, fig. 8. Upper Cretaceous; Austria.
- *elegans* (Corda), Endlicher, 1847 B, p. 305. Middle Cretaceous; Bohemia. (Recorded by Heer, 1883 A, Greenland; Patoot Beds.)
- *elegans linearis*, Hosius & von der Marck, 1880, p. 179, pl. xxxvii, fig. 142. Lower Senonian; Westphalia.
- *oxycedrus*, Presl in Sternberg, 1838, p. 203, pl. xlviii, fig. 3; pl. xlix, fig. 1. Cenomanian; Niederschöna, Saxony.
- *planifolius*, Endlicher, 1847 B, p. 305. (= *Cunninghamia planifolia*, Corda.) Perucer Beds; Bohemia.
- *prisca*, Eichwald, 1861, p. 311. Greensand; Russia.
- *pulchellus*, Knowlton in Stanton & Hatcher, 1905, p. 136, pl. xvi, fig. 1. Judith River Beds; Montana, U.S.A.
- *recurvatus*, Hosius & von der Marck, 1880, p. 179, pl. xxxvii, figs. 143, 144. Lower Senonian; Legden, Westphalia.
- *squamosus*, Heer, 1871 A, p. 9, pl. i, figs. 5-7. Senonian; Quedlinburg, Saxony.

- Cunninghamites squamosus densifolia**, Hosius & von der Marck, 1880, p. 178, pl. xxxvii, figs. 139, 140, 141.
Lower Senonian; Westphalia.
- **squamosus linearis**, Hosius & von der Marck, 1880, p. 179, pl. xxxvii, fig. 142.
Ibid.
- **Sternbergii**, Ettingshausen, 1867 A, p. 246, pl. i, figs. 4-6.
Cenomanian; Niederschoena, Saxony.
- ? **sp.**, Knowlton, 1900 A, p. 29, pl. v, fig. 3.
Montana Formation; Wyoming, U.S.A.
- Cupanites Novæ Zeelandiæ**, Ettingshausen, 1887 A, p. 188, pl. iv, figs. 18-20.
Upper Cretaceous; New Zealand.
- Cupressinea insignis**, Geinitz, 1850 A, p. 270.
Cenomanian; Niederschoena, Saxony.
- Cupressinoxylon ? Bibbinsi**, Knowlton, 1896 A, p. 584, text-figs. 1-4.
Upper Potomac Formation; Maryland, U.S.A.
- **cheyennense**, Penhallow, 1900, p. 76.
Comanche Cretaceous; Kansas, U.S.A.
- **columbianum**, Knowlton, 1889 A, p. 49, pl. iv, fig. 5; pl. v, figs. 3, 4; pl. vi, figs. 1-5; pl. vii, fig. 1.
Potomac Formation; Virginia, U.S.A.
- **comanchense**, Penhallow, 1900, p. 77.
Comanche Cretaceous (?); Kansas, U.S.A.
- **elongatum**, Knowlton, 1889 B, p. 7, pl. iii, figs. 1-4.
Laramie Formation; Montana, U.S.A.
- **Glasgowi**, Knowlton, 1889 B, p. 6, pl. ii, figs. 1-5.
Cretaceous (?); Iowa, U.S.A.
- **infracretaceum**, Fliche, 1896, p. 248, pl. xvii, figs. 1, 2.
Albian; Montblainville, France.
- **Kiprivanowi**, Mercklin in Krendowskij, 1880, p. 266, pl. i, figs. 4, 5.
Cretaceous; Isjum, Russia.
- **Lennieri**, Lignier, 1907, p. 275, pl. xix, figs. 25-28; pl. xxi, diag. 8; pl. xxii, fig. 73; pl. xxiii, fig. 88.
Lower Cretaceous; Hève, France.
- **McGeel**, Knowlton, 1889 A, p. 46, pl. ii, fig. 5; pl. iii, figs. 1-5.
Potomac Formation; Washington, U.S.A.
- **pulchellum**, Knowlton, 1889 A, p. 45, pl. ii, figs. 1-4.
Potomac Formation; Virginia, U.S.A.
- **sequoianum** (Mercklin), Vater emend. 1884, p. 813.
Lower Senonian; Harzburg.
- **Severzovi**, Mercklin, 1855, p. 59, pl. xiii, figs. 7-9.
Greensand; Russia.
- **turoniense**, Hosius & von der Marck, 1880, p. 199, pl. xlii, fig. 175 a-c.
Turonian; Westphalia.
- **ucranicum**, Goeppert, 1850 A, p. 201, pl. xxvi, figs. 1-4.
Cretaceous; Charcow, Ucranica.
- **vectense**, Barber, 1898, p. 337, pl. xxiii; pl. xxiv, figs. 1-15.
Lower Greensand; Isle of Wight.
- **Wardi**, Knowlton, 1889 A, p. 48, pl. iv, figs. 1-4; pl. v, figs. 1, 2.
Potomac Formation; Columbia, U.S.A.

- Cupressites Cookii**, Newberry, 1870, p. 9 [nomen nudum].
 Amboy Clay; New Jersey, U.S.A.
- **obtusifolius**, Eichwald, 1865, p. 44, pl. iii, figs. 6-7. Russia.
- Cupressoxylon cheyennense**, Penhallow, 1907 A, p. 238.
 Comanche Cretaceous; Kansas, U.S.A.
- **comanchense**, Penhallow, 1907 A, p. 239. Ibid.
- **Dawsoni**, Penhallow, 1907 A, p. 240.
 (Recorded in Cretaceous of Medicine Hat, Canada.)
- **macrocarpoides**, Penhallow, 1907 A, p. 238.
 Cretaceous; Medicine Hat, Canada.
- Cussonia ? lacerata**, Saporta, 1894, p. 189, pl. xxxv, fig. 3 B.
 Albian; Portugal.
- **partita**, Velenovsky, 1882 B, p. 20, pl. vii, fig. 1. (= *Cussoniophyllum partita*, Velenovsky, 1889.)
 Cenomanian; Hodkovic, Bohemia.
- Cussoniophyllum partitum**, Velenovsky, 1889, p. 22, pl. v, fig. 1.
 (= *Cussonia partita*, Velenovsky, 1882.) Cenomanian; Bohemia.
- Cyathea angusta**, Heer, 1883 A, p. 1, pl. i, figs. 4-5.
 Patoot Beds; Greenland.
- **fertilis**, Heer, 1882, p. 21, pl. xxxi, figs. 3-8.
 Atane Beds; Greenland.
- **Hammeri**, Heer, 1882, p. 22, pl. xxxi, figs. 1-2; pl. xxxv, figs. 4, 4b.
 Ibid.
- Cyatheetes ? nebraskana** (Heer), Knowlton, 1893, p. 81 (re-naming
Pecopteris nebraskana, Heer). Dakota Group; Kansas, U.S.A.
- Cycadeites ? Wohlfahrti** (Otto), Geinitz, 1875 B, p. 232, pl. lxi,
 figs. 3 a, b. (= *Arundites Wohlfahrti*, Otto, 1852).
 Quadersandstein; Dippoldiswalde.
- Cycadeocarpus (Dioonites) columbianus**, Dawson, 1873, p. 69,
 pl. i. Lower Cretaceous; British Columbia.
- Cycadeoidea Argonnensis**, Fliche, 1896, p. 153, pl. i, fig. 5.
 Albian; Clermont, France.
- **arida-gamantiensis**, Leriche, 1909, p. 501, text-fig. 3.
 Senonian; Templeux-la-Fosse, France.
- **aspera**, Ward, 1899 A, p. 213; & 1899 B, p. 624, pl. cxvii.
 Lower Cretaceous; Black Hills, U.S.A.
- **Bibbinsi**, Ward, 1897 C, p. 15; & 1905, p. 456, pl. lxxxiii, fig. 3;
 pl. lxxxiv, fig. 3; pl. lxxxv; pl. lxxxvii, figs. I, 2, 4, 6, 7; II, 1, 2, 4,
 8, 9, 10, 11; III, 5, 7; IV, 2, 3, 6, 8, 11; V, 12, 15; pl. lxxxix,
 figs. I, 5; II, 2, 5, 6; pls. ci-civ.
- **cicatricula**, Ward, 1899 A, p. 203; & 1899 B, p. 609, pls. lxxxiii-
 lxxxiv. Lower Cretaceous; Black Hills, U.S.A.
- **Clarkiana**, Ward, 1905, p. 472, pl. lxxxix, figs. I, 2, 4; pl. cvi.
 Potomac Formation; Maryland, U.S.A.
- **Colei**, Ward, 1899 A, p. 211; & 1899 B, p. 619, pls. ex-cxii.
 Lower Cretaceous; Black Hills, U.S.A.

- Cycadeoidea Colleti**, Fliehe, 1896, p. 151, pl. ii; pl. iii, figs. 1, 2.
Albian; Islettes, France.
- **colossalis**, Ward, 1899 A, p. 197; & 1899 B, p. 603, pls. lxxvii-lxxxii.
(See also Wieland, 1906, p. 98.)
Lower Cretaceous; Black Hills, U.S.A.
- **dakotensis** (McBride), Ward, 1894 B, p. 86. (= *Bennettites dakotensis*, McBride, 1893.) Illust. Wieland, 1906, pl. v, phot. 2, pls. xxxii, xxxiv, xxxv, xxxvi, xxxvii, xxxviii, xxxix, xl, xli, xlii, etc.
Lower Cretaceous; S. Dakota, U.S.A.
- **etrusca**, Capellini & Solms Laubach, 1892, p. 110, pl. i, fig. 2; pl. iv, fig. 1; pl. v, figs. 4, 7, 8. Lower Cretaceous (?); Italy.
- **excelsa**, Ward, 1899 A, p. 225; & 1899 B, p. 637, pls. cliii-clv.
Lower Cretaceous; Black Hills, U.S.A.
- **Ferretiana**, Capellini & Solms Laubach, 1892, pp. 84, 115.
Cretaceous (?); Italy.
- **Fisheræ**, Ward, 1905, p. 470, pl. lxxxvii, fig. III, 9; pl. cv.
Potomac Formation; Maryland, U.S.A.
- **Fontaineana**, Ward, 1897 c, p. 13; & 1905, p. 439, pl. lxxxvi; pl. lxxxvii, figs. I, 1; III, 2, 6, 8; IV, 1, 7, 9; V, 1, 10, 11, 13, 14, 16, 18, 21; pl. lxxxix, figs. I, 1; III, 1, 3, 6, 7, 11; pl. xcvi; pl. xevii; pl. xcviii.
Ibid.
- **forata** (Saporta), Capellini & Solms Laubach, 1892, p. 96.
(= *Clathropodium foratum*, Saporta, 1875.)
Gault; Havre, France.
- **formosa**, Ward, 1899 A, p. 222; & 1899 B, p. 634, pls. cxliv-cxlv.
See also Wieland, 1906, pl. xiv, phot. 2.
Lower Cretaceous; Black Hills, U.S.A.
- **furcata**, Ward, 1899 A, p. 210; & 1899 B, p. 618, pls. cvi-cix.
Ibid.
- **Gibsoni** (Carruthers), Ward. (= *B. Gibsonianus*, Carruthers, 1870.)
Lower Greensand; Isle of Wight.
- **Goucheriana**, Ward, 1897 c, p. 14; & 1905, p. 451, pl. lxxxvii, figs. 1, 3; pl. lxxxix, figs. 1, 3; pl. xcix. See also Wieland, 1906, p. 32, fig. 7.
Potomac Formation; Maryland, U.S.A.
- **Guillieri**, Crié, 1884, p. 512.
Mans, France.
- **helichorea**, Ward, 1900 A, p. 337, pl. iv, upper figure. (= *C. dakotensis*, Ward.) See also Wieland, 1906, p. 162.
Lower Cretaceous; Wyoming, U.S.A.
- **inclusa** (Carruthers), Schimper, 1874, p. 556. (= *Mantellia inclusa*, Carruthers, 1870.)
Potton Sands; Cambridgeshire.
- **ingens**, Ward, 1899 A, p. 221; & 1899 B, p. 632, pls. cxxxiii-cxliii.
See also illust. Wieland, 1906, pls. i, ii, iii, iv, iv A, etc.
Lower Cretaceous; S. Dakota, U.S.A.
- **insolita**, Ward, 1899 A, p. 214; & 1899 B, p. 625, pls. cxviii & cxix.
Lower Cretaceous; Black Hills, U.S.A.
- **Jenneyana**, Ward, 1894 B, p. 87; & 1899 B, p. 627, pls. cxxi-cxxiii.
See also illust. Wieland, 1906, pl. xiv, phot. 1, etc.
Ibid.

- Cycadeoidea Marshiana**, Ward, 1899 A, p. 208; & 1899 B, p. 616, pls. ci-cv. See also illust. Wieland, 1906, pl. v, phot. 1 & 3, pls. vii, viii, ix, phot. 1, xii, xiii, xxxiii, etc.
- Lower Cretaceous; Black Hills, U.S.A.
- **marylandica** (Font.), Capellini & Solms Laubach, 1892, p. 85. (= *Tysonia marylandica*, Fontaine, 1889.)
- Potomac Formation; Maryland, U.S.A.
- **Masseiana**, Capellini & Solms Laubach, 1892, p. 111, pl. i, fig. 1. (= *Raumeria Masseiana*, Capellini, 1890.) Cenomanian (?); Italy.
- **McBridei**, Ward, 1899 A, p. 205; & 1899 B, p. 612, pls. xci-c. See also Wieland, 1906. Lower Cretaceous; Black Hills; U.S.A.
- **McGeeana**, Ward, 1897 c, p. 12; & 1905, p. 434, pl. lxxxvii, figs. III, 3, 10, IV, 15, V, 8, 9, 19, 20; pl. lxxxix, figs. II, 1, 8, III, 4, 9, 10; pl. xciv. Potomac Formation; Maryland, U.S.A.
- **minima**, Ward, 1900 A, p. 341, pl. ii, lower fig.
- Lakota Formation; S. Dakota, U.S.A.
- **minnekahtensis**, Ward, 1899 A, p. 200; & 1899 B, p. 606, pls. lxxvi-lxxix. See also Wieland, 1906, pl. xliii, photos. 4, 5, etc.
- Lower Cretaceous; Black Hills, U.S.A.
- **mirabilis** (Lesquereux), Ward, 1894 B, p. 86. (= *Zamiostrobus*.)
- Pre-Laramie (?); Colorado, U.S.A.
- **munita**, Cragin, 1889, p. 65. Trinity Division; Kansas, U.S.A.
- **nana**, Ward, 1899 A, p. 227; & 1899 B, p. 639, pls. clvi & clvii.
- Lower Cretaceous; Black Hills, U.S.A.
- **Niedzwiedzki**, Raciborski, 1892, p. 355; & 1893, p. 301, pls. vii, viii.
- Lower Cretaceous (?); Carpathians.
- **occidentalis**, Ward, 1899 A, p. 215; & 1899 B, p. 626, pl. cxx.
- Lower Cretaceous; Black Hills, U.S.A.
- **Paynei**, Ward, 1899 A, p. 212; & 1899 B, p. 620, pls. cxiii-cxv. See also Wieland, 1906, pl. xliii, phot. 7, etc. Ibid.
- **protea**, Ward, 1900 A, p. 343, pl. iv, lower right fig. Ibid.
- **pulcherrima**, Ward, 1899 A, p. 201; & 1899 B, p. 608, pls. lxxx-lxxxii. See also Wieland, 1906, p. 171, fig. 91. Ibid.
- **reticulata**, Ward, 1900 A, p. 340, pl. iv, lower left fig. Ibid.
- **rhombica**, Ward, 1900 A, p. 336, pl. ii, upper fig. Ibid.
- **semiglobosa**, Fliche, 1896, p. 155, pl. iv, figs. 1, 2.
- Albian; Clermont, France.
- **Stantoni**, Ward, 1905, p. 276, pl. lxx.
- Shasta Formation; California, U.S.A.
- **Stillwalli**, Ward, 1899 A, p. 223; & 1899 B, p. 635, pls. cxlvii-clii. See also Wieland, 1906. Lower Cretaceous; Black Hills, U.S.A.
- **superba**, Ward, 1900 A, p. 334, pl. iii, lower fig. See also illust. Wieland, 1906, pl. ix, phot. 2; pls. x, xi, etc.
- Lowest Cretaceous (?); S. Dakota, U.S.A.
- **turrita**, Ward, 1899 A, p. 203; & 1899 B, p. 610, pls. lxxxv-xc; & Wieland, 1906, pl. xxviii, phot. 6, pl. xxx, phot. 2, etc.
- Lower Cretaceous; Black Hills, U.S.A.

- Cycadeoidea Tysoniana**, Ward, 1897 c, p. 11; & 1905, p. 432,
pl. lxxxvii, figs. 1, 5; pl. xciii.
Potomac Formation; Maryland, U.S.A.
- **Uhleri**, Ward, 1897 c, p. 14; & 1905, p. 454, pl. lxxxvii, figs. IV, 10;
pl. c. Ibid.
- **utopiensis**, Ward, 1900 a, p. 338, pl. iii, fig. 727.
Lakota Formation; Wyoming, U.S.A.
- **Wellsii**, Ward, 1899 a, p. 199; & 1899 b, p. 605, pls. lxxiii-lxxv.
See also Wieland, 1906, pl. xliii, phot. 6.
Lower Cretaceous; Black Hills, U.S.A.
- **Wielandi**, Ward, 1899, p. 621, pl. cxvi. (*C. Paynei*, ex parte.) See
also Wieland, 1906, p. 110, etc., pls. xv, xvi, xvii, phot. 4, xxi,
xxii, etc. Ibid.
- **Yatesii**, Carruthers, 1867, p. 199, pl. ix. (= *Yatesia Morrisii*, Car-
ruthers, 1870.) Lower Greensand; Potton.
- **sp.**, Fliche, 1896, p. 157, pl. v, fig. 1. Albain (?); Clermont, France.
- **sp.**, Wieland, 1900, p. 219, pl. i, fig. 12, text-fig. 13, p. 220.
Lower Cretaceous; S. Dakota, U.S.A.
- Cycadeospermum acutum**, Fontaine, 1889, p. 270, pl. cxxxv, fig. 12.
Potomac Formation; Virginia, U.S.A.
- **angustum**, Fontaine, 1889, p. 271, pl. cxxxv, fig. 20. Ibid.
- **californicum**, Fontaine in Ward, 1905, p. 257, pl. lxviii, fig. 4.
Shasta Formation; California, U.S.A.
- **columnare**, Lesquereux, 1892, p. 31, pl. xlv, figs. 7, 8.
Dakota Group; Kansas, U.S.A.
- **ellipticum**, Fontaine, 1889, p. 271, pl. cxxxv, fig. 19.
Potomac Formation; Virginia, U.S.A.
- **lineatum**, Lesquereux, 1892, p. 30, pl. i, fig. 14.
Dakota Group; Kansas, U.S.A.
- **montanense**, Fontaine in Ward, 1905, p. 310, pl. lxxiii, fig. 7.
Kootanie Formation; Montana, U.S.A.
- **obovatum**, Fontaine, 1889, p. 270, pl. cxxxv, fig. 13.
Potomac Formation; Virginia, U.S.A.
- **rotundatum**, Fontaine, 1889, p. 271, pl. cxxxvi, fig. 12. Ibid.
- **Schmidtianum**, Geinitz, 1879, p. 114, pl. iv, figs. 2 a, b, c.
Senonian; Silesia.
- **spatulatum**, Fontaine, 1889, p. 271, pl. cxxxv, figs. 11, 21.
Potomac Formation; Virginia, U.S.A.
- **turonicum**, Engelhardt, 1892, p. 94, pl. i, figs. 4, 5.
Turonian; Bohemia.
- Cycadinocarpus circularis**, Newberry; Ward in Smith, 1894,
p. 348; & Newberry, 1895, p. 46, pl. xlv, figs. 1-4.
Amboy Clay; Alabama, U.S.A.
- Cycadites affinis**, Eichwald, 1861, p. 311. Greensand; Russia.
- **cenomanensis**, Crie, 1879, p. 14 [nomen nudum].
Cenomanian; France.
- **contiguus**, Eichwald, 1865, p. 41, pl. iii, fig. 2.
Neocomian; Kursk, Russia.

- Cycadites Dicksoni**, Heer, 1871, p. 1182; & 1874 A, p. 99, pl. xxviii, fig. 7; pl. xxvii, fig. 9 c. (= *Pseudocycas Dicksoni* (Heer), Nathorst, 1907, p. 8) (= *Cycas Dicksoni*, Heer, 1882.) Atane Beds; Greenland.
- **giganteus**, Hisinger, 1837, p. 109, pl. xxxiii, fig. 5.
Greensand; Scania, Sweden.
- **linearis**, Sternberg, 1825, p. xxxiii, pl. 1, fig. 3.
Greensand; Hör, Sweden.
- **Nilssoni**, Sternberg, 1825, p. xxxii, pl. xlvii, fig. 1. (= *Dewalquea Nilssoni*, Nathorst, 1881 & 1894.) Ibid.
- **Nilssoniana (Nilssonianus)**, Brongniart, 1828, p. 98.
Greensand; Sweden.
- **pungens**, Lesquereux, 1892, p. 30, pl. ii, fig. 6.
Dakota Group; Kansas, U.S.A.
- **pygmæus**, Saporta, 1894, p. 173, pl. xxxi, fig. 10; pl. xxxii, fig. 5.
Albian; Portugal.
- **Sarthacensis**, Crié, 1879?, p. 22.
Cretaceous; Sainte-Croix, France.
- **Saxbyanus**, R. Brown, (1851) 1855, p. 130.
Cretaceous (?); Isle of Wight.
- **Schachti**, Coemans, 1867, p. 7, pl. iii, figs. 1-3.
Cretaceous; La Louvière, Belgium.
- **tenuisectus**, Saporta, 1894, p. 171, pl. xxxii, figs. 1-4, 6.
Albian; Portugal.
- **Unjiga**, Dawson, 1883, p. 20, pl. i, figs. 2, 2 a.
Upper Cretaceous; North-West Territory, Canada.
- **zamiaeifolius**, Sternberg, 1825, p. xxxiii, pl. xliii, fig. 3. (= *Zamites Schlotheimii*, Presl.)
Greensand; Hör, Sweden.
- Cycadopsis aquisgranensis**, Debey, 1848 B, p. 140. (= *Pinites aquisgranensis*, Goeppert, 1842 B.)
Senonian; Aix, Rhenish Prussia.
- **araucarina**, Debey, 1848, p. 141. Ibid.
- **cryptomerioides**, Miquel, 1833, p. 42, pl. iii, figs. 1-6.
Senonian; Limburg.
- **Foersteri**, Debey, 1848, p. 142. Ibid.
- **Monheimi**, Debey, 1848, p. 141. Ibid.
- **Pitzl**, Debey, 1848, p. 141. Ibid.
- **thujoides**, Debey, 1848, p. 142. Ibid.
- Cycadopteris Dunkeri**, Schenk, 1871 A, p. 6, pl. ii, figs. 1-2.
Senonian; Austrian Silesia.
- Cycadoxylum westfalicum**, Hosius & von der Mark, 1880, p. 193, pl. xli, figs. 164, 165. Lower Senonian; Haltern, Westphalia.
- Cycas Dicksoni**, Heer, 1882, p. 42, pl. xiv, fig. 10; pl. xvi, fig. 7.
(= *Cycadites Dicksoni*, Heer, 1874 A; *Pseudocycas Dicksoni*, Nathorst, 1907.) Atane Beds; Greenland.
- **Steenstrupi**, Heer, 1882, p. 40, pl. v, figs. 1 a, 1 b. (= *Pseudocycas Steenstrupi* (Heer), Nathorst, 1907.) Atane Beds; Greenland.
- **sp.**, Velenovsky, 1887 A, p. 642. Cenomanian; Bohemia.

- Cyclopitys Delgadoi**, Saporta, 1890 A, p. 814; & 1894, p. 91, pl. xviii, figs. 1-4. Valanginian; Portugal.
- Cyclopteris Klipsteini**, Dunker, 1846, p. 11, pl. ix, fig. 7. Neocomian; Duingen.
- **Moquensis**, Newberry in Ives, 1861, p. 129, pl. iii, figs. 1, 2. Base of Cretaceous (?); Arizona, U.S.A.
- **squamata**, Ettingshausen, 1852 c, p. 13, pl. iv, fig. 1. Wernsdorfer Beds (see Krasser, 1896, p. 146).
- Cylindrites arteriæformis**, Goepfert, 1842 A, p. 117, pl. i. Quadersandstein; Silesia.
- **conicus**, Hosius & von der Marck, 1880, p. 191, pl. xl, fig. 160. Lower Senonian; Belgium.
- ? **cretaceus**, Miquel, 1853, p. 55. Senonian; Belgium.
- **Daedaleus**, Goepfert, 1842 A, p. 117, pl. xlix, figs. 1, 2. Quadersandstein; Silesia.
- **latifrons**, Saporta, 1880 (1877), p. 642, pl. i, fig. 1. Lower Cretaceous; Villequier, France.
- **spongioides**, Goepfert, 1842 A, p. 115, pl. xlv, figs. 1-5; pl. xlviii, figs. 1, 2. (= *Typha gigantea*, Unger, 1870 A.) Quadersandstein; Silesia.
- **spongioides** (Goepfert), emend. Richter, 1909, p. 8, pls. xi, xii, xiii. Lower Cretaceous; Quedlinburg.
- Cyparissidium cretaceum**, Schenk, 1876, p. 167, pl. xxix, figs. 10-11. Upper Cretaceous; Brandenburg, Tyrol.
- **gracile**, Heer, 1874 A, p. 74, pl. xvii, figs. 5, b, c; pl. xviii, fig. 6 b; pl. xix; pl. xx, figs. 1, d, e; pl. xxi, figs. 9 b, 10 d. Kome Beds; Greenland.
- ? **japonicum**, Yokoyama, 1894, p. 229, pl. xx, figs. 3 a, 6, 6 a, 13; pl. xxiv, fig. 4. Neocomian (?); Japan.
- **minimum**, Velenovsky, 1885, p. 19, pl. ix, figs. 6-7; pl. x, fig. 4. Perucer Beds; Landsberg, Bohemia.
- **mucronatum**, Heer, 1883 A, p. 12, pl. xlviii, figs. 6 c, d, 16-17. Patoot Beds; Greenland.
- **pulchellum**, Velenovsky, 1885, p. 18, pl. v, figs. 3, 5, 6, 9. Cenomanian (?); Knezioka, Bohemia.
- **Suessii**, Schenk, 1876, p. 167, pl. xxviii, fig. 13 (*Widdringtonites*). Upper Cretaceous; St. Wolfgang, Austria.
- Cyperacites ambiguus**, Ettingshausen, 1893, pp. 139, 147; & 1895, p. 13, pl. i, fig. 3. Cretaceous; Australia.
- **arcticus**, Heer, 1874 A, p. 86, pl. xii, fig. 4 b. Kome Beds; Greenland.
- **hyperboreus**, Heer, 1874 A, p. 86, pl. xxiv, fig. 4. Ibid.
- **sp.**, Knowlton, 1900 A, p. 32, pl. v, fig. 8. Montana Formation; Wyoming, U.S.A.
- **sp.** (Dn.) Knowlton, 1898, p. 83 (re-naming *Cyperites sp.*, Dawson). Kootanie Formation; British Columbia.
- Cyperites sp.**, Dawson, 1893, p. 91, text-fig. 16. Ibid.

- Cyperites?* sp., Hollick, 1894 A, p. 63, pl. clxxx, fig. 3.
Middle Cretaceous; New York, U.S.A.
- Cytisus cretaceus*, Dunker, 1856, p. 182, pl. xxxiv, fig. 3.
Quadersandstein; Blankenburg, Saxony.
- Czekanowskia capillaris*, Newberry, 1895, p. 61, pl. ix, figs. 14-16.
Amboy Clay; New Jersey, U.S.A.
- (*Sclerophyllina*) *dichotoma*, Heer, 1868, p. 82, pl. xlv, fig. 6;
& 1882, p. 14. Kome Beds; Greenland.
- *nervosa*, Heer, 1881, p. 18, pl. xvii, figs. 5-7 a, 8-11.
Cretaceous; Almagem, Portugal.
- Dacrydium cupressinum*, Ettingshausen, 1887 A, p. 178, pl. vii,
figs. 17-18, 18 a. Upper Cretaceous; New Zealand.
- Dacrydites incertus*, Marik, 1901, p. 10, pl. i, fig. 20.
Cenomanian; Bohemia.
- Dacrydium densifolium*, Velenovsky, 1885, p. 12, pl. xii, figs. 1-4.
Ibid.
- Dactyolepis cryptomerioides*, Hollick & Jeffrey, 1909, p. 52, pl. x,
figs. 12, 13. Raritan Formation; New York, U.S.A.
- Dadoxylon ægyptiacum*, Unger, 1859, p. 228, pl. i, figs. 3-5.
(= *Araucarites Ægyptiacus*, Goeppert, and *Araucarioxylon Ægyptiacum*, Kraus in Schenk, 1883.) Cenomanian (?); Cairo, Egypt.
- *Dantzii*, Potonié, 1902 A, p. 229, pl. ii, figs. 1-8.
Upper Cretaceous (?); Dutch East Africa.
- Dalbergia apiculata*, Newberry, 1895, p. 90, pl. xliii, figs. 17-19.
Amboy Clay; Woodbridge, U.S.A.
- *hyperborea*, Heer, 1882, p. 102, pl. xxvi, fig. 4 a.
Atane Beds; Greenland.
- *irregularis*, Hollick, 1906 A, p. 85, pl. xxxii, fig. 11.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- *minor*, Hollick, 1906 A, p. 85, pl. xxxii, fig. 12. Ibid.
- *Rinkiana*, Heer, 1882, p. 102, pl. xxvi, figs. 1-3.
Atane Beds; Greenland.
- Dalbergiophyllum nelsonicum*, Ettingshausen, 1887 A, p. 189,
pl. vi, fig. 3. Upper Cretaceous; New Zealand.
- *phaseolitoides*, Ettingshausen, 1887 A, p. 189, pl. ix, fig. 17. Ibid.
- *rivulare*, Ettingshausen, 1887 A, p. 188, pl. vi, fig. 4. Ibid.
- Dammara acicularis*, Knowlton in Stanton & Hatcher, 1905, p. 134,
pl. xv, figs. 2-5. Judith River Beds; Montana, U.S.A.
- *borealis*, Heer, 1882, p. 54, pl. xxxviii, fig. 5.
Atane Beds; Greenland.
- ? *cliffwoodensis*, Hollick, 1898 A, p. 128, pl. xi, figs. 5-8.
Upper Cretaceous (Clay Marl); New Jersey, U.S.A.
- *macrosperma*, Heer, 1883 A, p. 17, pl. liii, fig. 11.
Patoot Beds; Greenland.
- *Mantelli*, Ettingshausen, 1887 A, p. 176, pl. vii, fig. 20.
Upper Cretaceous; New Zealand.

- Dammara microlepis**, Heer, 1882, p. 55, pl. xl, fig. 5.
Atane Beds; Greenland.
- **minor**, Hollick, 1906 A, p. 40, pl. ii, figs. 35-37. (= *Protodammara speciosa*, Hollick & Jeffrey, 1909.)
Raritan Formation; Staten Island, U.S.A.
- **northportensis**, Hollick, 1905 c, p. 405, pl. lxx, figs. 1-2.
Cretaceous Clay; Long Island, U.S.A.
- Dammarites albens**, Presl in Sternberg, 1838, p. 203, pl. lii, figs. 11, 12.
Upper Cretaceous; Balkans.
- **Bayeri**, Zeiller, 1905, p. 338, pl. vii, figs. 8-11.
Upper Cretaceous; Bulgaria.
- **caudatus**, Lesquereux, 1892, p. 32, pl. i, figs. 9, 10.
Dakota Group; Kansas, U.S.A.
- **crassipes**, Goeppert, 1842 A, p. 122, pl. liii, fig. 3.
Senonian; Silesia.
- **dubius**, Dawson, 1894, p. 56, pl. vi, fig. 8.
Upper Cretaceous; Vancouver Island, Canada.
- **emarginatus**, Lesquereux, 1892, p. 33, pl. i, fig. 11.
Dakota Group; Kansas, U.S.A.
- Dammaphyllum striatum**, Velenovsky, 1889, pp. 47 & 53.
(= *Podozamites striatus*, Velenovsky, 1885.)
Cenomanian; Bohemia.
- Danæites firmus**, Heer, 1868, p. 81, pl. xlv; & 1874 A, p. 56, pl. ix, fig. 1 a; pl. xii, figs. 1, 2.
Kome Beds; Greenland.
- **Schlotheimi**, Debey & Ettingshausen, 1859 B, p. 202, pl. iii, fig. 1.
Senonian; Aix, Rhenish Prussia.
- Daphnites Goepperti**, Ettingshausen, 1867 A, p. 253, pl. ii, fig. 8.
Cenomanian; Niederschoena, Saxony.
- Daphnogene cretacea**, Lesquereux, 1876 B, p. 343 (substituted for *Cinnamomum Scheuchzeri*, Heer, Lesquereux, 1874, p. 83, pl. xxx, figs. 2, 3).
Dakota Group; Kansas, U.S.A.
- **excellens**, Eichwald, 1865, p. 64, pl. iii, fig. 17.
Cretaceous; Russia.
- **Heerii**, Lesquereux, 1876 B, p. 343. (= *Cinnamomum Heerii*, Lesquereux, 1859.)
Dakota Group; Kansas, U.S.A.
- **primigenia**, Ettingshausen, 1867 A, p. 252, pl. i, fig. 13; pl. iii, fig. 15.
Cenomanian; Niederschoena, Saxony.
- Daphnophyllum angustifolium**, Lesquereux, 1892, p. 98, pl. xxxvi, fig. 8.
Dakota Group; Kansas, U.S.A.
- **crassinervium**, Heer, 1869 A, p. 18, pl. vii, fig. 2; pl. xi, fig. 5.
Upper Cretaceous; Moleteine, Moravia.
- **dakotense**, Lesquereux, 1892, p. 99, pl. li, figs. 1-4; pl. lii, fig. 1.
Dakota Group; Kansas, U.S.A.
- **ellipticum**, Heer, 1869 A, p. 18, pl. vii, fig. 3.
Upper Cretaceous; Moleteine, Moravia.
- **Fraasii**, Heer, 1869 A, p. 17, pl. vi, figs. 1, 2.
Ibid.
- Davallites Richardsoni**, Dawson, 1883, p. 25, pl. v, figs. 18, 18 a, 18 b.
Upper Cretaceous; Protection Island, Canada.

- Debeya affinis**, Ettingshausen, 1895, p. 46, pl. iii, figs. 21, 22.
Cretaceous; Australia.
- **australiensis**, Ettingshausen, 1893, pp. 137, 150; & 1895, p. 45, pl. iii, figs. 19, 20. Cretaceous; Australia.
- **serrata**, Miquel, 1853, p. 38, pl. i, fig. 1. (= *Phyllites Geinitzensis*, Goeppert, 1865.) Senonian; Limburg.
- **sp.**, Roemer, 1870, p. 354, pl. xxxix, fig. 10. Upper Cretaceous; Upper Silesia.
- Delesseria Friedaui**, Unger, 1850, p. 29; & 1853, p. 80, pl. xxvi, fig. 2. Upper Cretaceous; Styria.
- **incrassata**, Lesquereux, 1873, p. 374. (= *Caulerpites incrassatus*, Knowlton, 1898.) Laramie Formation; New Mexico, U.S.A.
- **lingulata**, Lesquereux, 1873, p. 374. (= *Caulerpites lingulatus*, Knowlton, 1898.) Ibid.
- **Reichii**, Engelhardt, 1892 A, p. 80. (= *Halysites Reichii*, Sternberg.) Cenomanian; Niederschoena, Saxony.
- Delessertites Hampeanus**, Stiehler, 1858, p. 56, pl. xi, fig. 12. Quadersandstein; Blankenburg, Saxony.
- **Thierensi**, Miquel, 1853, p. 54, pl. i, fig. 4. (= *Phyllites Thierensi*, Bosquet MS., Debey, 1851.) Senonian; Limburg.
- Delgadopsis rhizostigma**, Saporta, 1894, p. 141, pl. xxiii, figs. 3, b-c; pl. xxv, figs. 1-4, 9-18; pl. xxvi, fig. 2. Urgonian; Cercal, Portugal.
- Dermatophyllites acutus**, Heer, 1882, p. 80, pl. xlii, fig. 7. Atane Beds; Greenland.
- **borealis**, Heer, 1874 A, p. 112, pl. xxxii, figs. 8, 8b. Ibid.
- Dewalquea aquisgranensis**, Saporta & Marion, 1873, p. 61, pl. viii, figs. 5-7. Senonian; Aix, Rhenish Prussia.
- **coriacea**, Velenovsky, 1889, p. 23, pl. iv, figs. 1-6. (= *Aralia coriacea*, Velenovsky.) Cenomanian; Bohemia.
- **dakotensis**, Lesquereux, 1892, p. 211, pl. lix, figs. 5, 6. Dakota Group; Kansas, U.S.A.
- **gelindenensis**, Saporta & Marion, recorded Hosius & v. d. Marck, 1880, p. 174, pl. xxxiv, fig. 124. Upper Senonian; Westphalia.
- **groenlandica**, Heer, 1882, p. 87, pl. xxix, figs. 18-19; pl. xlii, figs. 5-6; pl. xliiv, fig. 11. Atane Beds; Greenland.
- **haldemiana angustifolia**, Hosius & von der Marck, 1880, p. 173, pl. xxxiii, figs. 116, 117; pl. xxxiv, figs. 118-122. Upper Senonian; Haldem, Westphalia.
- **haldemiana latifolia**, Hosius & von der Marck, 1880, p. 173, pl. xxxiv, fig. 115; pl. xxxv, fig. 114. Ibid.
- **insignis**, Hosius & von der Marck, 1880, p. 172, pl. xxxii, figs. 111-113; pl. xxxiii, fig. 109; pl. xxxiv, fig. 110; pl. xxxv, fig. 123. Ibid.
- **Nilssoni** (Nilsson), Nathorst, 1881 A, p. 83; & 1894, p. 196, text-fig. (= *Cycadites Nilssoni*, Sternberg, 1825.) Greensand; Scania, Sweden.

- Dewalquea pentaphylla**, Velenovsky, 1886, p. 14, pl. viii, figs. 11, 12.
Cenomanian; Bohemia.
- **primordialis**, Lesquereux in Winchell, 1885, p. 77; & Lesquereux, 1895, p. 18, pl. A, fig. 10. Dakota Group; Minnesota, U.S.A.
- **Smithi**, Berry, 1910 E, p. 36, text-fig. 1.
Tuscaloosa Formation; Alabama, U.S.A.
- **trifoliata**, Newberry, 1895, p. 129, pl. xxii, figs. 4-7.
Amboy Clay; Woodbridge, U.S.A.
- Diceras cenomanicus**, Velenovsky, 1889, p. 14, pl. ii, figs. 5-7.
Cenomanian; Vyserovic, Bohemia.
- Dicksonia borealis**, Heer, 1882, p. 23, pl. xlv, fig. 2.
Atane Beds; Greenland.
- **conferta**, Heer, 1882, p. 23, pl. xxxv, figs. 5-7. Ibid.
- **groenlandica**, Heer, 1882, p. 23, pl. xxxv, figs. 8-9. Ibid.
- **microphylla**, Heer, 1878, p. 27, pl. viii, figs. 1-4.
Lower Cretaceous (?); Atyrkan, Siberia.
- **montanensis**, Fontaine in Ward, 1905, p. 286, pl. lxxi, figs. 1-4.
Kootanie Formation; Montana, U.S.A.
- **munda**, Dawson, 1886, p. 11, pl. iii, figs. 5, 5 A.
Mill Creek Series; Mill Creek, Canada.
- **pachyphylla**, Fontaine in Ward, 1905, p. 224, pl. lxx, fig. 1.
Shasta Formation; California, U.S.A.
- **ptericides**, Ettingshausen, 1887 A, p. 175, pl. vii, figs. 4-6.
Upper Cretaceous; New Zealand.
- **punctata** (Sternberg), Heer, 1882, p. 24, pl. xlvii. (= *Protopteris punctata*, Sternberg.)
Atane Beds; Greenland.
- **tosana**, Yokoyama, 1894, p. 213, pl. xxv, figs. 13, 13 a.
Neocomian; Ryoseki, Japan.
- Dicksoniopteris Naumanni**, Nathorst, 1890, p. 11, pl. v, fig. 4.
Recorded Yokoyama, 1894, p. 214, pl. xxv, fig. 4. Ibid.
- Dicotylophyllum cerciforme**, Saporta, 1894, p. 147, pl. xxvi, fig. 14.
Urgonian; Cercal, Portugal.
- **corrugatum**, Saporta, 1894, p. 148, pl. xxvi, fig. 16. Ibid.
- **hederaceum**, Saporta, 1894, p. 148, pl. xxvi, fig. 15. Ibid.
- **lacerum**, Saporta, 1894, p. 149, pl. xxvi, fig. 3. Ibid.
- Dicropteris longifolia**, Pomel, 1849, p. 339. (= *Baiera longifolia* (Pomel), Heer, 1876.)
- Dictyophyllum Dicksoni**, Heer, 1871, p. 1181; 1874 A, p. 55, pl. iii, figs. 9, 9 b, c, d.
Kome Beds; Greenland.
- **sp.**, Heer, 1878, p. 29, pl. viii, figs. 16 b, 16 c, 18.
Lower Cretaceous; Siberia.
- Dictyopteris anomala**, Saporta, 1894, p. 81, pl. xv, fig. 28.
Valanginian; Portugal.
- **infracretacea** [see *infracretacica*], Saporta, 1894, p. 99. Ibid.
- **infracretacica** (error for *infracretacea*), Saporta, 1894, p. 81, pl. xvi, fig. 20.
- **tenella**, Saporta, 1894, p. 82, pl. xv, fig. 25. Ibid.

- Didymosorus comptoniaefolius**, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **comptonifolius**, Debey & Ettingshausen, 1859 B, p. 186, pl. i, figs. 1-5. (= *Zonopteris comptoniaefolia*, Debey, 1848. = *Gleichenia comptoniaefolia*, Heer, 1874 A.) Ibid.
- **gleichenioides**, Debey & Ettingshausen, 1859 B, p. 190, pl. i, figs. 10, f. Ibid.
- **varians**, Debey, 1849, p. 299; & Debey & Ettingshausen, 1859 B, p. 190, pl. i, figs. 7-9. Ibid.
- Diemenia lancifolia**, Ettingshausen, 1893, p. 149; & 1895, p. 24, pl. iii, fig. 4. Cretaceous; Australia.
- Dillenia** sp., Otto, 1854, p. 47, pl. ix, figs. 5-7. Dippoldiswalde, Saxony.
- Dicönites borealis**, Dawson, 1883, p. 24, pl. iii, fig. 37. Lower (?) Cretaceous; North-West Territory, Canada.
- **Buchianus** (Ett.), Bornemann, 1856, p. 57 (no descript.). (= *Pterophyllum Buchianum*, Ettingshausen, 1852. Recorded from American Potomac, Fontaine, 1889.) Wealden; Silesia.
- **Buchianus abietinus** (Goeppert), Ward, 1905, p. 250, pl. lxvii, figs. 1-3 (re-naming *Pterophyllum abietinum*, Goeppert). Shasta Formation; California, U.S.A.
- **Buchianus**, var. **angustifolius**, Fontaine, 1889, p. 185, pl. lxvii, fig. 6; pl. lxviii, fig. 4; pl. lxxi, fig. 2. Potomac Formation; Virginia, U.S.A.
- **Buchianus**, var. **obtusifolius**, Fontaine, 1889, p. 184, pl. clxviii, fig. 3. Ibid.
- **Buchianus rarinervis**, Fontaine, 1894, p. 264, pl. xxvii, figs. 3, 4. Trinity Division; Texas, U.S.A.
- **columbianus**, Dawson, 1893, p. 91. (= *Cycadeospermum (Dioönites) columbianus*, Dn. 1873.) Lower Cretaceous; Queen Charlotte Island, Canada.
- **cretosus** (Reiche), Schimper, 1872, p. 211. (= *Pterophyllum cretosum*.) Cenomanian; Niederschoena, Saxony.
- **Dunkerianus** (Goeppert), Miquel. Recorded Fontaine, 1894, p. 265, pl. xxxvi, fig. 12; pl. xxxvii, fig. 1. American Trinity Division.
- Dioscorea? cretacea**, Lesquereux, 1874, p. 56, pl. xxviii, fig. 10. Dakota Group; Kansas, U.S.A.
- Dicspyrophyllum provectum**, Velenovsky, 1889, pp. 50, 53. (= *Diospyros provecta*, Velenovsky, 1886.) Cenomanian; Bohemia.
- Diospyros ambigua**, Lesquereux, 1892, p. 110. (= *D. anceps*, Lx., 1874, & *Quercus anceps*, Lx., 1868.) Dakota Group; Nebraska, U.S.A.
- **amboyensensis**, Berry, 1909, p. 262 (re-naming *Phyllites ellipticus*). Raritan Formation; New Jersey, U.S.A.
- **anceps**, Lesquereux, 1874, p. 89, pl. vi, fig. 6. (= *Quercus anceps*, Lesquereux, 1868, p. 96.) Dakota Group; Nebraska, U.S.A.

- Diospyros apiculata**, Lesquereux, 1892, p. 110, pl. xiv, fig. 3.
Dakota Group; Kansas, U.S.A.
- **?celastroides**, Lesquereux, 1892, p. 113, pl. xx, fig. 7.
Dakota Group; Kansas, U.S.A.
- **cretacea**, Ettingshausen, 1893, pp. 138, 150; & 1895, p. 39, pl. iii, figs. 17, 18, 24.
Cretaceous; Australia.
- **eminens**, Dawson, 1894, p. 62, pl. x, fig. 40.
Upper Cretaceous; Vancouver Island, Canada.
- **judithæ**, Knowlton in Stanton & Hatcher, 1905, p. 146, pl. xviii, figs. 4, 5; pl. xix, fig. 3. Judith River Beds; Montana, U.S.A.
- **nitida**, Dawson, 1883, p. 22, pl. iii, fig. 10.
Cretaceous; Peace River, North-West Territory, Canada.
- **primæva**, Heer in Capellini & Heer, 1867, p. 19, pl. i, figs. 6, 7.
Dakota Group; Nebraska, U.S.A.
- **prodrcmus**, Heer, 1874 a, p. 113, pl. xxxii, figs. 3-7; pl. xxviii, fig. 6 c.
Atane Beds; Greenland.
- **provecta**, Velenovsky, 1886, p. 49, pl. xxiii, figs. 1-5, 10.
(= *Diospyrophyllum provectum*, Velenovsky, 1889.)
Cenomanian; Bohemia.
- **pseudoanceps**, Lesquereux, 1892, p. 111, pl. xxii, fig. 1; & 1895, p. 17, pl. B, fig. 6. Dakota Group; Minnesota, U.S.A.
- **rotundifolia**, Lesquereux, 1874, p. 89, pl. xxx, fig. 1.
Dakota Group; Kansas, U.S.A.
- **Schweinfurthi**, Heer, 1876 b, p. 6, figs. 1-10 (fruits).
Upper Cretaceous; Egypt.
- **Steenstrupi**, Heer, 1883 a, p. 32, pl. lxiv, fig. 1.
Patoot Beds; Greenland.
- **vancouverensis**, Dawson, 1883, p. 28, pl. viii, fig. 32.
Upper Cretaceous; Vancouver Island.
- **Wodani**, Unger, 1850 a, p. 435. Recorded American Laramie Formation, Lesquereux, 1878, p. 233, pl. lix, fig. 13.
- **sp.**, Dawson, 1894, p. 62, pl. x, fig. 41.
Cretaceous; British Columbia, Canada.
- Diphyllites membranaceus**, Heer, 1883 a, p. 45, pl. lx, fig. 4 a.
Patoot Beds; Greenland.
- Diplopora Mühlbergii**, Lorenz, 1902, p. 52, text-figs. 3-6.
Lower Cretaceous; Switzerland.
- Dipteriphyllum cretaceum** (Velenovsky), Krasser, 1896, p. 123, pl. xv, fig. 7. (= *Platyserium cretaceum*, Velenovsky).
Perucer Beds; Vyserovic, Bohemia.
- Discophorites angustilobus**, Heer, 1877, p. 145, pl. lviii, figs. 18, 19.
Neocomian; Switzerland.
- **Fischeri**, Heer, 1877, p. 145, pl. lviii, figs. 16, 17. Ibid.
- **Schneiderianus**, Geinitz, 1879, p. 113, pl. iv, fig. 1.
Neocomian(?); Caucasia.
- Dombeyopsis obtusa**, Lesquereux, 1873, p. 375; & 1878, p. 255, pl. xlvii, figs. 4, 5.
Laramie Formation; Colorado, U.S.A.

- Dombeyopsis obtusiloba**, Lesquereux, 1868, p. 100. (= *Menispermites obtusiloba*, var.?, Lesquereux, 1874.) Dakota Group; Nebraska, U.S.A.
 — **platanoides**, Lesquereux, 1878 B, p. 254, pl. xlvii, figs. 1-2.
 Laramie Formation; Montana, U.S.A.
 — **trivialis**, Lesquereux, 1873, p. 380; & 1878 B, p. 255, pl. xlviii, fig. 3.
 Laramie Formation; Colorado, U.S.A.
Dombeyoxylon ægyptiacum, Schenk, 1883, p. 13.
 Cretaceous; Cairo, Egypt.
Dorstenia ? sp., Penhallow, 1907, p. 310, text-fig. 5.
 British Columbia, Canada.
Dracæna australis, Morris MS., Goeppert in Bronn, 1849, p. 37 [nomen nudum].
 Cretaceous (?); Europe.
 — **Benstedti**, König MS. in Mantell, 1851, p. 49; & Mackie, 1862 B, pl. xxii. (= *Benstedtia* sp., Seward, 1896 B.)
 Lower Greensand; Maidstone.
Dracænites Jourdei, Marion, 1890, p. 1054 [nomen nudum].
 Turonian; Martignes, France.
Dryandra antiqua, Ettingshausen, 1851, p. 739 (re-naming *Comptonites antiqua*, Nilsson).
 Cretaceous; Sweden.
 — **cretacea**, Velenovsky, 1882 B, p. 12 [nomen nudum]; & 1883, p. 1, pl. ix, figs. 1-5.
 Cretaceous; Bohemia.
 — **Huttoniana**, Crié, 1889, p. 79 (5) [nomen nudum].
 Cretaceous; New Zealand.
 — **pteroides**, Ettingshausen, 1851, p. 737, pl. xxxii, fig. 9.
 Cretaceous; near Trieste.
Dryandroides coriacea, Velenovsky, 1882 A, p. 213 [nomen nudum].
 Cretaceous; Bohemia.
 — **geinoglypha**, Bayer, 1896, p. 18, text-figs. 11, 12, & p. 35.
 Upper Senonian; Kieslingswalde, Bohemia.
 — **haldemiana**, Hosius & von der Marck, 1880, p. 168, pl. xxxi, figs. 91-100; pl. xxxii, figs. 101-104.
 Upper Senonian; Haldem, Westphalia.
 — **hieraciifolia**, Debey (in litt.), see Hosius & v. d. Marck, 1880, p. 166. (= *Quercus hieraciifolia*.)
 Ibid.
 — **latifolius**, Ettingshausen, 1867 A, p. 257, pl. iii, fig. 10.
 Cenomanian; Niederschoena, Saxony.
 — (**Myrica**) **macrophylla**, Hosius & v. d. Marck, 1880, p. 169, pl. xxxii, fig. 105.
 Upper Senonian; Haldem, Westphalia.
 — **minor**, Feistmantel, 1874, p. 275.
 Perucer Beds; Bohemia.
 — **pakawauica**, Ettingshausen, 1887 A, p. 186, pl. ix, fig. 13.
 Upper Cretaceous; New Zealand.
 — **quercinea**, Velenovsky, 1882 A, p. 213; & 1883, p. 33, pl. x, figs. 8 a-15.
 Cenomanian; Bohemia.
 — **serratus**, Velenovsky, 1882 A, p. 213 [nomen nudum].
 Cretaceous; Bohemia.
 — **Zenkeri**, Ettingshausen, 1867 A, p. 257, pl. iii, figs. 1, 3, 11. (= *Salix fragiliformis*, Zenker, 1833.)
 Cenomanian; Niederschoena, Saxony.

- Drynaria astrostigmosa*, Bayer, 1899, p. 9, pl. i, figs. 5, 6, text-figs. 4, 4 a. Perucur Beds; Bohemia.
- *dura* (Velenovsky), Bayer, 1899, p. 15, text-figs. 6, 6 a. (= *Lambertia dura*, Velenovsky, 1883.) Ibid.
- *fascia*, Bayer, 1899, p. 10, text-figs. 5, 5 a. Perucur Beds; Vyserovic, Bohemia.
- *tumulosa*, Bayer, 1899, p. 19, pl. i, figs. 1, 1 a, 2 a, 3, 4. Ibid.
- Dryophyllum Alberti-Magni*, Debey, 1881, p. 89, fig. 6 on plate. Senonian; Aix, Rhenish Prussia.
- *anceps*, Lesquereux, MS. Nebraska, U.S.A.
- *aquamarum*, Ward, 1885, p. 551, pl. xxxvii, figs. 3-5. Laramie Formation; Wyoming, U.S.A.
- *aquisgranense*, Debey, 1881, p. 88, fig. 1 on plate. Senonian; Aix, Rhenish Prussia.
- *basidentatum*, Ward, 1885, p. 551, pl. xxxvii, fig. 11. Laramie Formation; Wyoming, U.S.A.
- *Beuthianum*, Debey, 1881, p. 96, fig. 22 on plate. Senonian; Aix, Rhenish Prussia.
- *bruneri*, Ward, 1887, p. 27, pl. x, figs. 5, 6. Montana Formation; Wyoming, U.S.A.
- *camptoneurum*, Debey, 1881, p. 96. Senonian; Aix, Rhenish Prussia.
- *crenatum*, Lesquereux, 1876 A, p. 371; & 1878 B, p. 162, pl. lxii, figs. 10, 11. Montana Formation; Wyoming, U.S.A.
- *Crepini*, Debey, 1881, p. 94, fig. 18 on plate. Senonian; Aix, Rhenish Prussia.
- *cretaceum*, Debey, 1881, p. 88, figs. 2-5 on plate. Ibid.
- *Dethimusianum*, Debey, 1881, p. 95, fig. 20 on plate. Ibid.
- *elongatum*, Dawson, 1894, p. 58, pl. vii, fig. 20. Upper Cretaceous; Vancouver Island, Canada.
- *Eodrys*, Debey, 1881, p. 94, fig. 19 on plate. Senonian; Aix, Rhenish Prussia.
- *exiguum*, Debey, 1881, p. 95, fig. 21 on plate. Ibid.
- *falcatum*, Ward, 1885, p. 551, pl. xxxvii, fig. 10. Laramie Formation (?); Wyoming, U.S.A.
- *gracile*, Debey, 1881, p. 90, figs. 10-11 on plate. Senonian; Aix, Rhenish Prussia.
- *Heeri*, Debey, 1881, p. 89, figs. 7, 8 on plate. Ibid.
- (*Quercus*) *Holmesii*, Lesquereux, 1883, p. 38, pl. iv, fig. 8. (= *Quercus Holmesii*, Lesquereux, 1892.) Dakota Group (?); Colorado, U.S.A.
- (*Quercus*) *latifolium*, Lesquereux, 1876 A, p. 393; & 1876 B, p. 340, pl. vi, fig. 1. Dakota Group; Kansas, U.S.A.
- *Lerschianum*, Debey, 1881, p. 93, figs. 15-16 on plate. Senonian; Aix, Rhenish Prussia.
- *Lesquereuxianum*, Debey, 1881, p. 93, fig. 17 on plate. Ibid.
- *Lesquereuxii*, Ettingshausen, 1893, p. 148; & 1895, p. 16, pl. i, fig. 30. Cretaceous; Australia.

- Dryophyllum Neillianum**, Dawson, 1894, p. 58, pl. vii, fig. 19.
Upper Cretaceous; Vancouver Island, Canada.
- **Nelsnicum**, Ettingshausen, 1887 A, p. 182, pl. viii, figs. 11, 11 a.
Upper Cretaceous; New Zealand.
- **occidentale**, Dawson, 1894, p. 58, pl. vii, figs. 17, 18. (Re-named by Knowlton, 1898, *Quercus* (?) *Dryophyllum* *occidentalis*.)
Upper Cretaceous; Vancouver Island, Canada.
- **primordiale**, Lesquereux, 1883, p. 37. (= *Quercus primordialis*, 1868.)
Dakota Group; Kansas, U.S.A.
- **regaliaquense**, Debey, 1881, p. 92, figs. 12-14 on plate.
Senonian; Aix, Rhenish Prussia.
- (*Quercus*) **salicifolium**, Lesquereux, 1876 A, p. 399; & 1876 B, p. 340, pl. viii, fig. 2.
Dakota Group (?); Colorado, U.S.A.
- **subfalcatum**, Lesquereux, 1876 c, p. 379.
Laramie Formation; Wyoming, U.S.A.
- **tenuifolium**, Debey, 1881, p. 90, fig. 9 on plate.
Senonian; Aix, Rhenish Prussia.
- **westphaliense**, Saporta, 1867, p. 35.
Quadersandstein; Haldem, Westphalia.
- **sp.**, Dawson, 1894, p. 58, pl. xii, fig. 59.
Upper Cretaceous; Vancouver Island, Canada.
- (*Dryophanes*) **sp.**, Debey, 1881, p. 97, figs. 23-26 on plate.
Quadersandstein; Aix, Rhenish Prussia.
- Dryopteris angustipinnata** (Font.), Knowlton, 1898, p. 91 (re-naming *Aspidium angustipinnatum*, Fontaine, 1889).
Potomac Formation; Virginia, U.S.A.
- **angustipinnata montanense** (Font.), Knowlton, 1898, p. 91 (re-naming *Aspidium angustipinnata montanense*, Fontaine, 1893).
Kootanie Formation; Montana, U.S.A.
- **duttoniana**, Knowlton, 1900 A, p. 4 [nomen nudum].
Montana Formation; Wyoming, U.S.A.
- **Kennerleyi** (Newberry), Knowlton, 1898, p. 92 (re-naming *Aspidium Kennerlyi*, Newberry, 1863).
Upper Cretaceous; Vancouver Island.
- ? **kootaniensis**, Knowlton, 1907, p. 111, pl. xi, figs. 4, 4 A.
Kootanie Formation; Montana, U.S.A.
- **Oerstedii** (Heer), Knowlton, 1898, p. 92 (re-naming *Aspidium Oerstedii*, Heer, 1882).
Atane Beds; Greenland.
- **spp.**, Knowlton, 1898, pp. 91-93, re-naming various species of *Aspidium*.
- Echinostrobus minor**, Velenovsky, 1889, p. 10, pl. i, figs. 11, 12, 15.
Cenomanian; Vyserovic, Bohemia.
- **squammosus**, Velenovsky, 1885, p. 16, pl. vi, figs. 3, 6, 7, 8.
Ibid.
- Elæodendron marylandicum**, Berry, 1910 II, p. 24, pl. viii, fig. 1.
Magothy Formation; Maryland, U.S.A.

- Elæodendron priscum**, Ettingshausen, 1893, p. 150; & 1895, p. 48,
pl. iv, fig. 6. Cretaceous; Australia.
- **speciosum**, Lesquereux, 1892, p. 175, pl. xxxvi, figs. 2, 3.
Dakota Group; Kansas, U.S.A.
- **strictum**, Hollick, 1906 A, p. 89, pl. xxxii, fig. 6.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- **sp.**, Hollick, 1906 A, p. 89, pl. xxxiii, fig. 7.
Cretaceous; Long Island, U.S.A.
- Elate oblonga**, Unger, 1845, p. 199. (= *Abies oblonga*, Lindley &
Hutton, 1835.)
- Elatoxylon Withamii**?, Hartig, 1848, p. 168.
Quadersandstein; Quedlinburg, Saxony.
- Eleoxylon cretaceum**, Brongniart, 1849 A, pp. 76, 111. (= *Pinus*
cretacea, Corda.) Cretaceous; Bohemia.
- Embothrites**? **daphneoides**, Lesquereux, 1883, p. 51. (= *Embo-*
thrium? *daphneoides*, Lesquereux, 1874.)
- Embothrium**? **daphneoides**, Lesquereux, 1874, p. 87, pl. xxx, fig. 10.
Dakota Group; Kansas, U.S.A.
- Encephalartopsis nervosa**, Fontaine, 1889, p. 174, pl. lxx, fig. 4;
pl. lxxi, figs. 3, 4; pl. lxxii, figs. 3, 4.
Potomac Formation; Virginia, U.S.A.
- Encephalartos cretaceus**, Lesquereux, 1892, p. 29, pl. i, fig. 12.
Dakota Group; Kansas, U.S.A.
- Eolirion lusitanicum**, Saporta, 1894, p. 180, pl. xxxiv, figs. 2-3.
Albian; Portugal.
- ? **nervosum**, Hosius & von der Marck, 1880, p. 143, pl. xxvi,
fig. 24. Upper Senonian; Haldem, Westphalia.
- **primigenium**, Schenk, 1871 A, p. 20, pl. vii, fig. 4.
Urgonian; Grod'schitz, Austrian Silesia.
(Recorded Heer, 1874 A, pl. xxiv, figs. 1-3, Greenland.)
- **subfalcatum**, Hosius & von der Marck, 1880, p. 142, pl. xxvi,
fig. 23. Upper Senonian; Westphalia.
- **sp.**?, Hosius & von der Marck, 1880, p. 182, pl. xxxvii, fig. 150.
Lower Senonian; Westphalia.
- Ephedrites baccatus**, Marik, 1901, p. 14, pl. ii, fig. 3.
Cenomanian; Bohemia.
- **i vernonensis**, Fontaine in Ward, 1905, p. 495, pl. cvii, fig. 8.
Lower Potomac Formation; Virginia, U.S.A.
- Equisetites annularioides**, Heer, 1874 A, p. 61, pl. xiii, fig. 9.
Kome Beds; Greenland.
- **Burchardti**, Dunker, 1846, p. 2, pl. v, fig. 7. Wealden.
(Recorded from Urgonian of Portugal, etc.)
- **grönlandicus**, Heer, 1874 A, p. 61, pl. xiii, fig. 10.
Kome Beds; Greenland.
- **inæqualis**, Eichwald, 1861, p. 310. Greensand; Russia.
- **notabilis**, Eichwald, 1865, p. 34, pl. iv, fig. 7. Neocomian; Russia.
- **peruanus**, Neumann, 1907, p. 78, pl. ii, figs. 1, 2.
Neocomian; Peru.

- Equisetites?* sp., Jasche, 1858, p. 93, pl. iv, fig. 3.
Quadersandstein; Prussian Saxony.
- Equisetum amissum*, Heer, 1874 A, p. 60, pl. xiii, figs. 2-8; pl. xxii, figs. 11 b, c.
Kome Beds & Atane Beds; Greenland.
- *arenarium*, Hampe, 1852, p. 7 [nomen nudum].
Quadersandstein; Blankenburg, Saxony.
- *Burchardti*, Dunker. (= *Equisetites Burchardti*, Dunker, 1846.)
- *Heerii*, Schenk, 1876, p. 165, pl. xxix, fig. 1.
Upper Cretaceous; Brandenburg, Tyrol.
- *Lyelli*, Mantell, 1833, p. 245, text-figs. 1-3.
Wealden; Tilgate Forest.
(Recorded from American Potomac, Fontaine, 1889.)
- *marylandicum*, Fontaine, 1889, p. 65, pl. ii, fig. 10.
Potomac Formation; Maryland, U.S.A.
- *maximum*, Hampe, 1852, p. 7 [nomen nudum].
Quadersandstein; Blankenburg, Saxony.
- *montanense*, Fontaine in Weed & Pirsson, 1898, p. 481.
Judith River Formation; Montana, U.S.A.
- *nodosum*, Lesquereux, 1883, p. 25.
Dakota Group; Kansas, U.S.A.
- *robustum*, Newberry, 1863, p. 513; & 1898, p. 15, pl. xvi, figs. 1, 2.
Tertiary & Cretaceous; Washington, U.S.A.
- *texense*, Fontaine, 1894, p. 263, pl. xxxvi, fig. 1.
Trinity Division; Texas, U.S.A.
- *virginicum*, Fontaine, 1889, p. 63, pl. i, figs. 1-6, 8; pl. ii, figs. 1-3, 6, 7, 9.
Potomac Formation; Virginia, U.S.A.
- *Zeilleri*, Richter, 1905, p. 7, pl. i, figs. 2, 12.
Quadersandstein; Quedlinburg, Saxony.
- *sp.?*, Fontaine, 1889, p. 65, pl. ii, fig. 8.
Potomac Formation; Virginia, U.S.A.
- *sp.*, Heer, 1874 A, p. 124, pl. xxxviii, fig. 8. Cretaceous; Spitzbergen.
- Eremophyllum fimbriatum*, Lesquereux, 1874, p. 107, pl. viii, fig. 1.
(= *Ficus? fimbriata*, Lesquereux, 1868.)
Dakota Group; Nebraska, U.S.A.
- Etheridgea subglobosa*, Ettingshausen, 1893, pp. 141-150; & 1895, p. 46, pl. iv, fig. 3.
Cretaceous; Australia.
- Ettingshausenia cuneifolia*, Stiehler, 1858, p. 67. (= *Credneria cuneifolia*, Bronn.)
Cenomanian; Niederschoena, Saxony.
- *cuneiformis*, Krasser, 1889, p. 34; & 1896, p. 116 [nomen nudum].
Cenomanian; Kunstadt, Moravia.
- *expansa*, Stiehler, 1858, p. 67. (= *Credneria expansa*, Brongniart, 1849 A.)
Cenomanian; Niederschoena, Saxony.
- *Geinitziana*, Stiehler, 1858, p. 67. (= *Credneria Reichi*, Geinitz, and *C. Geinitziana*, Unger.)
Ibid.
- *grandidentata*, Stiehler, 1858, p. 67. (= *Credneria grandidentata*, Unger, 1849.)
Ibid.
- *irregularis*, Krasser, 1889, p. 34; & 1896, p. 116 [nomen nudum].
Cenomanian; Kunstadt, Moravia.

- Ettingshausenia mcraivica**, Krasser, 1896, p. 116 [nomen nudum].
Cretaceous; Moravia.
- **pseudo-Guillelmæ**, Krasser, 1889, p. 34; & 1896, p. 116 [nomen nudum].
Cenomanian; Kunstadt, Moravia.
- **reticulata**, Stiehler, 1858, p. 67. (= *Credneria reticulata*, Eichwald, 1853.)
Neocomian; Kursk, Russia.
- **spathulata**, Stiehler, 1858, p. 67. (= *Credneria spathulata*, Eichwald, 1853.)
Ibid.
- **Sternbergi**, Stiehler, 1858, p. 67. (= *Credneria Sternbergii*, Brongniart, 1849 A.)
Cenomanian; Teschen, Bohemia.
- **tremulæfolia**, Stiehler, 1858, p. 67. (= *Credneria tremulæfolia*, Brongniart, 1849 A.)
Cenomanian; Niederschoena, Saxony.
- **venulosa**, Stiehler, 1858, p. 67. (= *Credneria venulosa*, Eichwald, 1853.)
Neocomian; Kursk, Russia.
- **sp.**, Meek & Hayden, 1859 c, p. 222, text-fig. 3.
Lower Cretaceous; Kansas, U.S.A.
- Eucalyptophyllum oblongifolium**, Fontaine, 1889, p. 325, pl. clxii, fig. 4.
Potomac Formation; Virginia, U.S.A.
- Eucalyptus angusta**, Velenovsky, 1887, p. 64, pl. xxvi, figs. 2-12.
(= *E. angustus*, Velenovsky, 1889, p. 21, pl. vi, fig. 1.)
Cenomanian; Vyserovic, Bohemia.
- ? **angustifolia**, Newberry, 1895, p. 111, pl. xxxii, figs. 1, 6, 7.
Amboy Clay; South Amboy, New Jersey, U.S.A.
- ? **attenuata**, Newberry, 1895, p. 111, pl. xvi, figs. 2, 3, 5. (= *Ficus daphnogenoides*, Berry.)
Ibid.
- **baldemiana**, Debey in Hosius & v. d. Marck, 1880, p. 174. (Misprint (?) for *E. haldemiana*.)
- **borealis**, Heer, 1882, p. 94, pl. xl, figs. 3, 4; pl. xli, fig. 14.
Atane Beds; Greenland.
- **Chcffati**, Saporta, 1894, p. 207, pl. xxxvii, fig. 1.
Upper Albian; Portugal.
- **cretacea**, Ettingshausen, 1893, p. 150; & 1895, p. 48, pl. iv, figs. 7, 8.
Cretaceous; Australia.
- **dakotensis**, Lesquereux, 1892, p. 137, pl. xxxvii, figs. 14-19.
Dakota Group; Kansas, U.S.A.
- **Davidsoni**, Ettingshausen, 1893, p. 150; & 1895, p. 49, pl. iv, fig. 10.
Cretaceous; Australia.
- ? **dubia**, Berry (*non* Ettingshausen, 1887), 1905 n, p. 87, pl. lii, fig. 1.
(= *Eucalyptus Wardiana*, Berry, 1905 A.)
Matawan Formation; New Jersey, U.S.A.
- **Geinitzi**, Heer, 1882, p. 93, pl. xix, fig. 1 c; pl. xiv, figs. 4-9; pl. xli, figs. 12, 13. (= *Myrtophyllum Warderi*, Lesquereux, 1892, = *M. Geinitzi*, Heer.)
Atane Beds; Greenland.
- **Gouldii**, Ward, 1897 A, p. 576, text-figs. 1, 2, p. 577.
Dakota Group; Kansas, U.S.A.
- **haldemiana**, Debey in Hosius & von der Marck, 1880, p. 174, pl. xxxv, figs. 125-128.
Upper Senonian; Westphalia.
- **inæquilatera**, von der Marck, 1864, p. 77, pl. xiii, fig. 1. Ibid.

- Eucalyptus latifolia**, Hollick, 1906 A, p. 97, pl. xxxvi, figs. 1-5.
Middle Cretaceous; Long Island, U.S.A.
- **linearifolia**, Berry, 1907, p. 203 (re-naming *Eucalyptus nervosa*, Newberry).
Middle Cretaceous; N. Carolina, U.S.A.
- ? **nervosa**, Newberry MS. in Hollick, 1894 A, p. 56, pl. clxxiv, fig. 10; & in Newberry, 1895, p. 112, pl. xxxii, figs. 3, 4, 5, 8.
(= *Eucalyptus linearifolia*, Berry, 1907.)
Amboy Clay; Long Island, U.S.A.
- **oxleyana**, Ettingshausen, 1893, p. 150; & 1895, p. 49, pl. iv, fig. 9.
Cretaceous; Australia.
- ? **parvifolia**, Newberry, 1895, p. 112, pl. xxxii, figs. 9, 10.
Amboy Clay; South Amboy, New Jersey, U.S.A.
- **proto-Geinitzii**, Saporta, 1894, p. 206, pl. xxxvi, fig. 16; pl. xxxvii, fig. 11.
Upper Albian; Portugal.
- **rosieriana**, Ward, 1905, p. 530, pl. cxiii, figs. 9, 10.
Older Potomac Formation; Maryland, U.S.A.
- **Schübleri** (Heer), Hollick, 1906 A, p. 96, pl. xxxvi, fig. 6.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- **scoliophylla**, Ettingshausen, 1893, p. 150; & 1895, p. 49, pl. iv, figs. 12, 13.
Cretaceous; Australia.
- **Wardiana**, Berry, 1905 A, p. 47. (= *Eucalyptus dubia*, Berry, 1905 B, non Ettingshausen.)
Middle Cretaceous; New Jersey, U.S.A.
- **warraghiana**, Ettingshausen, 1893, p. 150; & 1895, p. 50, pl. iv, fig. 11.
Cretaceous; Australia.
- **sp.**, Krasser, 1896, p. 116, pl. xii, fig. 3.
Cenomanian; Kunstadt, Moravia.
- Eugenia primæva**, Lesquereux, 1892, p. 137, pl. liii, figs. 5-9.
Dakota Group; Kansas, U.S.A.
- Eugeinitzia proxima**, Hollick & Jeffery, 1909, p. 43, pl. x, fig. 10; pl. xxv, figs. 1-3.
Raritan Formation; Staten Island, U.S.A.
- Euonyminium Auerbachi**, Mercklin, 1855, p. 23, pl. i, fig. 3; pl. iii.
Greensand; Russia.
- Euphorbiophyllum antiquum**, Saporta & Marion, 1885, p. 117, text-fig. 125 c, e.
Turonian; France.
- **primordiale**, Saporta, 1894, p. 218, pl. xxxix, fig. 23.
Cenomanian; Portugal.
- Eurysaxis squamosa** (Heer), Schulze, 1888, p. 18. (= *Cunninghamites squamosus*, Heer.)
Senonian; Altenburg.
- Fagophyllum nervosum**, Dawson, 1894, p. 58, pl. vii, fig. 16.
Upper Cretaceous; Vancouver Island, Canada.
- **retosum**, Dawson, 1894, p. 57, pl. vii, fig. 15.
Ibid.
- Fagoxylon hokkaidense**, Stopes & Fujii, 1910, pp. 64-66, pl. vii, figs. 50-53.
Upper Cretaceous; Hokkaido, Japan.
- Fagus cretacea**, Newberry, 1870, p. 23; & 1878, pl. ii, fig. 3; & 1898, p. 68, pl. i, fig. 3.
Dakota Group; Kansas, U.S.A.
- **leptoneura**, Ettingshausen, 1893, p. 134; 1895, p. 20, pl. ii, fig. 9.
Cretaceous; Australia.

- Fagus Nelsonica**, Ettingshausen, 1887 A, p. 183, pl. ix, figs. 9, 9a.
Upper Cretaceous; New Zealand.
- **orticulata**[-um], Lesquereux, 1892, p. 51, pl. xlvii, fig. 6.
Dakota Group; Kansas, U.S.A.
- **polyclada**, Lesquereux, 1868, p. 95; & 1874, p. 67, pl. v, fig. 6.
Dakota Group; Nebraska, U.S.A.
- **præ-ninnisiana**, Ettingshausen, 1893, p. 135; & 1895, p. 21,
pl. ii, figs. 1-5. Cretaceous; Australia.
- **præ-ulmifolia**, Ettingshausen, 1893, p. 137; & 1895, p. 21, pl. ii,
figs. 6-8. Ibid.
- **prisca**, Ettingshausen, 1867 A, p. 249, pl. ii, figs. 3, 3b.
Cenomanian; Niederschoena, Saxony.
- **producta**, Ettingshausen, 1887 A, p. 183, pl. ix, fig. 1.
Upper Cretaceous; New Zealand.
- **proto-nucifera**, Dawson, 1883, p. 21, pl. ii, figs. 6 & 6a. [Isolated
leaf, and also fruit given the same name.]
Cretaceous; Peace River, North-West Territory, Canada.
- Fasciculites ambiguus**, Eichwald, 1865, p. 71, pl. v, fig. 7.
Neocomian; Moscow, Russia.
- **groenlandicus**, Heer, 1868, p. 85, pl. xlv, fig. 23.
Kome Beds; Greenland.
- **ovata**, Stenzel, 1872, p. 72 [nomen nudum].
- **varians**, Unger, 1850 A, p. 339. (= *Palmacites varians*, Corda, 1846.)
Cretaceous; Bohemia.
- Fasciostelopteris Tansleii**, Stopes & Fujii, 1910, pp. 10-15, pl. i,
fig. 7; pl. ii, figs. 2, 3; text-fig. 4.
Upper Cretaceous; Hokkaido, Japan.
- Fegonium dryandraforme**, Vater, 1884, p. 838, pl. xxviii, figs. 7-10.
Lower Senonian (?); Brunswick.
- **Schenki**, Vater, 1884, p. 839, pl. xxviii, figs. 11-14. Ibid.
- Feildeniopsis crassinervis**, Fontaine, 1889, p. 205, pl. lxxxv, fig. 5.
Potomac Formation; Virginia, U.S.A.
- Feistmantelia oblonga**, Ward, 1899 B, p. 693, pl. clxix, fig. 19.
Lower Cretaceous; Black Hills, U.S.A.
- **virginica**, Fontaine in Ward, 1905, p. 484, pl. cvii, fig. 3.
Older Potomac Formation; Virginia, U.S.A.
- Ficophyllum crassinerve**, Fontaine, 1889, p. 291, pl. cxlv, fig. 3;
pl. cxlv, fig. 3; pl. cxlvi, fig. 1; pl. cxlvii, fig. 4; pl. cxlviii, figs. 1,
2, 4; pl. clvii, fig. 4; pl. clxxiii, fig. 10.
Potomac Formation; Virginia, U.S.A.
- **eucalyptoides**, Fontaine, 1889, p. 294, pl. cxlv, figs. 1-2. Ibid.
- **serratum**, Fontaine, 1889, p. 294, pl. cxlv, fig. 2; pl. cxlix, fig. 9.
Ibid.
- **tenuinerve**, Fontaine, 1889, p. 292, pl. cxl, fig. 3; pl. cxli, fig. 2;
pl. cxlv, figs. 1, 4; pl. cxlvii, fig. 2; pl. cxlix, figs. 1, 3, 5; pl. clvi,
fig. 1. Ibid.
- Ficoxylon cretaceum**, Schenk, 1883, p. 14, pl. v, figs. 17-19.
Cretaceous; Cairo, Egypt.

- Ficus aligera**, Lesquereux, 1892, p. 84, pl. x, figs. 3-6. (Re-named *Citrophylllum aligerum*, Berry, 1909.)
Dakota Group; Kansas, U.S.A.
- **angulata**, Hosius & von der Mark, 1880, p. 166, pl. xxxi, fig. 89.
Upper Senonian; Haldem, Westphalia.
- ? **angustata**, Lesquereux, 1883, p. 47.
Dakota Group; Kansas, U.S.A.
- **angustifolia**, Hosius, 1870 A, p. 99, pl. xv, figs. 21-22.
Senonian; Legden, Westphalia.
- **arenacea brevipetiolata**, Lesquereux, 1878 B, p. 195, pl. xxix, figs. 2, 5.
Laramie Formation; Wyoming, U.S.A.
- ? **arctica**, Heer, 1883 A, p. 26, pl. lxx, fig. 4.
Patoot Beds; Patoot, Greenland.
- **asarifolia**, Ettingshausen, 1867 B, p. 156, pl. xxv, figs. 2, 3, 6.
Tertiary; Friesen.
(Recorded by Lesquereux from Montana Formation, U.S.A.)
- **asarifolia minor**, Lesquereux, 1876 C, p. 367.
Montana Formation; Wyoming, U.S.A.
- **atavina**, Heer, 1882, p. 69, pl. xi, figs. 5 b, 7 b, 8 b; pl. xvii, fig. 8 b;
pl. xix, fig. i b; pl. xx, figs. 1, 2. (= *Ficus protogaea*, Heer, 1874 A.)
Atane Beds; Greenland.
- **austiniana**, Lesquereux in Winchell, 1885, p. 76; & Lesquereux, 1895, p. 14, pl. A, fig. 5.
Dakota Group; Minnesota, U.S.A.
- **Beckwithii**, Lesquereux, 1883, p. 46, pl. xvi, fig. 5; pl. xvii, figs. 3, 4.
Dakota Group; Colorado, U.S.A.
- **Berthoudi**, Lesquereux, 1892, p. 78, pl. xii, fig. 3 (pre-occupied name, see *F. Lesquereuxii*, Kn.).
Dakota Group; Kansas, U.S.A.
- **bumelioides**, Ettingshausen, 1867 A, p. 251, pl. ii, fig. 6.
Cenomanian; Niederschoena, Saxony.
- **Cecropiæ-lobus**, Bayer, 1893, pp. 15, 41, fig. 12; & Bayer in Fritsch, 1893, p. 130, text-figs. 1 & 4.
Senonian; Priesen, Bohemia.
- **contorta**, Dawson, 1894, p. 60, pl. ix, fig. 31.
Upper Cretaceous; Vancouver Island, Canada.
- ? **corylifolius**, Lesquereux, 1873, p. 394 [nomen nudum].
Laramie Formation (?); Wyoming, U.S.A.
- **crassinervis**, Hosius, 1870 A, p. 99, pl. xvi, figs. 25-26.
Senonian; Legden, Westphalia.
- **crassipes**, Heer, 1882, p. 70, pl. xvii, fig. 9 a; pl. xxiv, figs. 1, 2; pl. xlii, fig. 2 c; pl. xliii, figs. 4-6; pl. xlvi, fig. 15. (= *Proteoides crassipes*, Heer, 1874 A.)
Atane Beds; Greenland.
- **cretacea**, Hosius, 1870 A, p. 99, pl. xv, figs. 19-20.
Senonian; Legden, Westphalia.
- ? **cuneatus**, Newberry, 1863, p. 524.
Cretaceous (?); Washington State, U.S.A.
- **daphnogenoides** (Heer), Berry, 1905 B, p. 329, pl. xxi. (= *Proteoides daphnogenoides*, Heer in Capellini & Heer, 1867.)
Middle Cretaceous; Maryland, U.S.A.

- Ficus deflexa**, Lesquereux, 1892, p. 80, pl. iii, fig. 13; pl. xvi, fig. 3.
Dakota Group; Kansas, U.S.A.
- **densinervis**, Hosius & von der Marek, 1880, p. 135, pl. xxv, figs. 10, 11, 12.
Upper Senonian; Westphalia.
- **dentata**, Hosius, 1870 A, p. 100, pl. xvi, fig. 27.
Senonian; Legden, Westphalia.
- **distorta**, Lesquereux, 1876 A, p. 393; & 1876 B, p. 342, pl. v, fig. 5.
Dakota Group; Kansas, U.S.A.
- **elongata**, Hosius (*non* Velenovsky), 1870 A, p. 98, pl. xiv, figs. 15-16.
Senonian; Legden, Westphalia.
- **elongata**, Velenovsky (*non* Hosius), 1882 A, p. 213; & 1883, p. 40, pl. xii, fig. 4.
Cenomanian; Bohemia.
- ? **fimbriatus**, Lesquereux, 1868, p. 96. (= *Eremophyllum fimbriatum*, Lesquereux, 1874.)
Dakota Group; Nebraska, U.S.A.
- **fracta**, Velenovsky, 1887, p. 71, pl. xxxi, fig. 15.
Cenomanian; Kieslingswalde.
- **Fredericksburgensis**, Fontaine, 1889, p. 295, pl. cxlviii, figs. 3, 5.
Potomac Formation; Virginia, U.S.A.
- **Geinitzii**, Ettingshausen, 1867 A, p. 250, pl. ii, figs. 7, 9-11.
Cenomanian; Niederschoena, Saxony.
- **Glascoena**, Lesquereux, 1883, p. 48.
Dakota Group; Kansas, U.S.A.
- **gracilis**, Hosius, 1870 A, p. 99, pl. xv, figs. 23-24.
Senonian; Legden, Westphalia.
- ? **halliana**, Lesquereux, 1874, p. 68, pl. xxviii, figs. 3, 9.
Dakota Group; Minnesota, U.S.A.
- **Haydenii**, Lesquereux, 1873, p. 394; & 1878 B, p. 197, pl. xxx, fig. 1.
Laramie Formation; Wyoming, U.S.A.
- **Hellandiana**, Heer, 1882, p. 70, pl. xxxvii, fig. 8.
Atane Beds; Greenland.
- **hesperia**, Knowlton, 1900 A, p. 45, pl. ix, fig. 5.
Montana Formation; Wyoming, U.S.A.
- **inæqualis**, Lesquereux, 1892, p. 82, pl. xlix, figs. 6-8; pl. i, fig. 3.
Dakota Group; Kansas, U.S.A.
- **incompleta**, Knowlton, 1900 A, p. 46, pl. ix, fig. 2.
Montana Formation; Wyoming, U.S.A.
- **ipswichiana**, Ettingshausen, 1895, p. 22, pl. ii, fig. 12.
Cretaceous; Australia.
- **Ipswichii**, Ettingshausen, 1893, p. 137.
Ibid.
- **irregularis**, Lesquereux, 1876 C, p. 368; & 1878 B, p. 196, pl. xxxiv, figs. 4-7; pl. lxiii, fig. 9. (= *Ulmus? irregularis*, Lx., 1873.)
Montana Formation; Wyoming, U.S.A.
- **Krausiana**, Heer, 1869 A, p. 15, pl. v, figs. 3-6.
Upper Cretaceous; Moletein, Moravia.
- **lanceolato-acuminata**, Ettingshausen, 1872, p. 182, pl. vi, figs. 3, 4. (Recorded Lesquereux, 1892, American Dakota.)
Tertiary; Sagor, Croatia.

- Ficus laurifolia**, Hosius & von der Marck, 1880, p. 136, pl. xxv, fig. 13.
Upper Senonian; Westphalia.
- **laurophylla**, Lesquereux, 1876 B, p. 342, pl. v, fig. 7 (substituted for *Laurophyllum reticulatum*).
Dakota Group; Kansas, U.S.A.
- **laurophyllidia**, Dawson, 1894, p. 60, pl. x, fig. 37.
Upper Cretaceous; Vancouver Island, Canada.
- **Lesquereuxii**, Knowlton, 1898, p. 102 (re-naming *Ficus Berthoudi*, Lx., 1892).
Dakota Group; Kansas, U.S.A.
- **longifolia**, Hosius, 1870 A, p. 99, pl. xv, figs. 17-18.
Senonian; Legden, Westphalia.
- **macrophylla**, Lesquereux, 1892, p. 76, pl. xi, fig. 1.
Dakota Group; Kansas, U.S.A.
- **magnoliæfolia**, Lesquereux, 1883, p. 47, pl. xvii, figs. 5-6. Ibid.
- **maxima**, Dawson, 1883, p. 21, pl. ii, fig. 5.
Upper Cretaceous; North-West Territory, Canada.
- **melanophylla**, Lesquereux, 1892, p. 83, pl. l, fig. 2.
Dakota Group; Kansas, U.S.A.
- **missouriensis**, Knowlton, 1900 A, p. 12, pl. i, fig. 5.
Montana Formation; Coal Banks, U.S.A.
- **Mohlana**, Heer, 1869 A, p. 15, pl. v, fig. 2.
Upper Cretaceous; Moletsein, Moravia.
- **montana**, Knowlton, 1900 A, p. 48, pl. xi, figs. 2, 3; pl. xii, fig. 1.
Montana Formation; Wyoming, U.S.A.
- **Mudgei**, Lesquereux, 1892, p. 83, pl. xii, fig. 4.
Dakota Group; Kansas, U.S.A.
- **myricoides**, Hollick in Newberry, 1895, p. 71, pl. xxxii, fig. 18; pl. xli, figs. 8, 9.
(?) Amboy Clay; New Jersey, U.S.A.
- **neurocarpa**, Hollick, 1903, p. 103, text-fig. B, 1, p. 104.
Dakota Group; Kansas, U.S.A.
- **oblanceolata**, Lesquereux, 1873, p. 387 (re-named *Pistacia oblanceolata*, Knowlton, 1898).
Laramie Formation; Wyoming, U.S.A.
- **ovata**, Newberry, 1895, p. 70, pl. xxiv, figs. 1-3 (re-named *Ficus ovatifolia*, Berry, 1909).
Amboy Clay; New Jersey, U.S.A.
- **ovatifolia**, Berry, 1909, p. 253 (re-naming *Ficus ovata*, Newberry, 1895).
Raritan Formation; New Jersey, U.S.A.
- **pedunculata**, Feistmantel, 1874, p. 276.
Perucer Beds; Niederschoena, Saxony; and Bohemia.
- **Peruni**, Velenovsky, 1883, p. 41, pl. xii, figs. 1-3.
Cenomanian; Bohemia.
- **planicostata**, Lesquereux, 1873, p. 393; & Knowlton, 1900 A, p. 52, pl. x, fig. 4.
Montana Formation; Wyoming, U.S.A.
- **planicostata latifolia**, Lesquereux, 1873, p. 393; & 1878 B, p. 202, pl. xxxi, fig. 9.
Laramie Formation; Wyoming, U.S.A.
- **populcides**, Knowlton, 1900 A, p. 44, pl. viii, fig. 3.
Montana Formation; Wyoming, U.S.A.
- **præcursor**, Lesquereux, 1892, p. 81, pl. xlix, fig. 5.
Dakota Group; Kansas, U.S.A.

- Ficus primordialis*, Heer in Capellini & Heer, 1867, p. 16, pl. iii, fig. 1.
Dakota Group; Nebraska, U.S.A.
- *problematica*, Knowlton, 1900 A, p. 46, pl. ix, fig. 3.
Montana Formation; Wyoming, U.S.A.
- *proteoides*, Lesquereux, 1892, p. 77, pl. xii, fig. 2. (= *Ficus daphnogenoides* (Heer), Berry, 1905 B.)
Dakota Group; Kansas, U.S.A.
- *protogæa*, Ettingshausen (*non* Heer), 1867 A, p. 249, pl. ii, fig. 5.
Cenomanian; Niederschoena, Saxony.
- *protogæa*, Heer (*non* Ettingshausen), 1871, p. 1183; & 1874 A, p. 108, pl. xxx, figs. 1-8; pl. xxix, fig. 2 b. (= *Ficus atavina*, Heer, 1882.)
Atane Beds; Greenland.
- *reticulata* (Lesquereux), Hollick in Newberry, 1898, p. 88, pl. xxii, figs. 2, 3 (re-naming *Laurophyllum reticulatum*, Lx., 1873).
Dakota Group; Kansas, U.S.A.
- *Reuschii*, Hosius, 1870 A, p. 98, pl. xiv, figs. 13-14.
Senonian; Legden, Westphalia.
- *Reussii*, Ettingshausen in Reuss, 1854, p. 740 [nomen nudum].
Cenomanian; Moletuin, Moravia.
- *ramnoides*, Knowlton, 1900 A, p. 47, pl. x, figs. 1-3; pl. xi, fig. 1.
Montana Formation; Wyoming, U.S.A.
- ? *rhomboides*, Lesquereux, 1868, p. 96.
Dakota Group; Nebraska, U.S.A.
- *rotundata*, Dawson, 1894, p. 60, pl. ix, figs. 32, 33 a.
Upper Cretaceous; Vancouver Island, Canada.
- *sapindifolia*, Hollick, 1905 c, p. 411, pl. lxxviii, fig. 5.
Cretaceous Clay; Long Island, U.S.A.
- *similis*, Ettingshausen, 1887 A, p. 185, pl. ix, figs. 10, 10 a.
Upper Cretaceous; New Zealand.
- *sinuosa*, Ward, 1887, p. 41, pl. xxii, fig. 2.
Laramie Formation; Wyoming, U.S.A.
- *speciosissima*, Ward, 1887, p. 39, pl. xxi, fig. 3.
Montana Formation; Wyoming, U.S.A.
- *squarrosa*, Knowlton, 1900 A, p. 45, pl. viii, fig. 2. Ibid.
- *Stephensoni*, Berry, 1910 A, p. 194, pl. xxiii, figs. 2, 3.
Middle Cretaceous; North Carolina, U.S.A.
- *Sternbergii*, Lesquereux, 1873, p. 423; & 1892, p. 82, pl. 1, fig. 1.
Dakota Group; Kansas, U.S.A.
- *stylosa*, Velenovsky, 1882 A, p. 213; & 1883, p. 39, pl. xii, fig. 5.
Cenomanian; Bohemia.
- *subgoepperti*, Shirley, 1902, p. 7, pl. i.
Upper Cretaceous; Queensland, Australia.
- *suspecta*, Velenovsky, 1887, p. 71, pl. xxviii, figs. 6, 9.
Perucer Beds; Vyserovic, Bohemia.
- *tenuifolia*, Hosius, 1870 A, p. 100, pl. xvi, fig. 28.
Senonian; Legden, Westphalia.
- *trinervis*, Knowlton, 1900 A, p. 42.
Montana Formation; Wyoming, U.S.A.

- Ficus? undulata*, Lesquereux, 1892, p. 84, pl. xii, fig. 5.
Dakota Group; Kansas, U.S.A.
- *virginiensis*, Fontaine, 1889, p. 295, pl. cxliii, figs. 1, 3; pl. cxliv, fig. 1.
Potomac Formation; Virginia, U.S.A.
- *Wardii*, Knowlton, 1900 A, p. 48, pl. ix, fig. 1.
Montana Formation; Wyoming, U.S.A.
- *Wellingtoniæ*, Dawson, 1894, p. 60, pl. ix, figs. 33, 34.
Upper Cretaceous; Vancouver Island, Canada.
- *Willisiana*, Hollick, 1894 A, p. 52, pl. clxxvi, figs. 2, 5.
Cretaceous; Long Island, U.S.A.
- *Woolsoni*, Newberry MS., in Hollick, 1893, p. 33, pl. ii, figs. 1, 2 c; & Newberry, 1895, p. 70, pl. xx, fig. 3; pl. xxiii, figs. 1-6.
Amboy Clay; New York, U.S.A.
- *sp.*, Hollick, 1903, p. 104, text-fig. B. 2, 3.
Dakota Group; Kansas, U.S.A.
- *sp.*, Lesquereux, 1892, p. 85, pl. x, figs. 7, 8. Ibid.
- Filicites sp.*, Nilsson, 1824, p. 146, pl. ii, figs. 1-3. Greensand; Sweden.
- Fittonia squamata*, Carruthers, 1870, p. 690, pl. lvi. (= *Clathraria anomala*, in part, Mantell, 1847.)
Upper Cretaceous (?); Isle of Wight.
- Flabellaria chamæropifolia*, Goeppert, 1836, p. 426; & 1842 A, p. 120, pl. lii, figs. 1-3. Senonian; Silesia.
- *coryphæfolia*, Goeppert (probably error for *F. chamæropifolia*), 1836, p. 87.
- *longirachis*, Unger, 1850 A, p. 332; & 1853, p. 91, pl. xxxi, fig. 1; pl. xxxii, fig. 1. Cretaceous; Austria.
- *magothiensis*, Berry, 1905 c, p. 32, text-figs. 1, 2.
Magothy Formation; Maryland, U.S.A.
- ? *minima*, Lesquereux, 1874, p. 56, pl. xxx, fig. 12.
Dakota Group; Kansas, U.S.A.
- *sub-longirachis*, Ettingshausen, 1887 A, p. 181, pl. viii, figs. 4, 4 a.
Upper Cretaceous; New Zealand.
- Folium arcuatum*, Velenovsky, 1882 A, p. 214 [nomen nudum].
Cretaceous; Bohemia.
- *daphnophyllum*, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- *lanceolatum*, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- *lingua*, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- *productum*, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- *ranunculus*, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- Fontainea grandifolia*, Newberry, 1895, p. 96, pl. xlv, figs. 1-4.
Amboy Clay; Woodbridge, U.S.A.
- Fraxinus præcox*, Heer, 1883 A, p. 33, pl. lxiv, fig. 2.
Patoot Beds; Greenland.
- Frenelites Reichii*, Ettingshausen, 1867 A, p. 246, pl. i, figs. 10 a-10 c.
(= *Widdringtonites Reichii*, Heer, 1882.)
Cenomanian; Niederschoena, Saxony.

- Frenelopsis bohémica*, Velenovsky, 1888 A, p. 590, pl. (unnumb.), figs. 1-3, 10; & 1889, p. 13, pl. ii, figs. 3, 4.
Cenomanian; Vyserovic, Bohemia.
- *gracilis*, Newberry, 1895, p. 59, pl. xii, figs. 1-3 a.
Amboy Clay; New Jersey, U.S.A.
- *Hoheneggeri* (Ett.), Schenk, 1871 A, p. 13, pl. iv, figs. 5-7; pl. v, figs. 1, 2; pl. vi, figs. 1-6; pl. vii, fig. 1. (Recorded Heer from Greenland, 1874.)
Wernsdorfer Beds; Carpathians.
- *Königii*, Hosius & von der Marck, 1880, p. 132, pl. xxxvii, fig. 148. (= *Calamitopsis Königii*, v. d. Marck, 1864.)
Upper Senonian; Westphalia.
- *leptoclada*, Saporta, 1894, p. 100, p. 113, pl. xix, fig. 18; pl. xxi, figs. 9-11.
Neocomian (? Aptian); Portugal.
- *occidentalis*, Heer, 1881, p. 21, pl. xii, figs. 3 b, 4-7.
Cretaceous; Portugal.
- *parceramosa*, Fontaine, 1889, p. 218, pl. cxi, figs. 1-5; pl. cxii, figs. 1-5; pl. cxviii, fig. 1. Potomac Formation; Virginia, U.S.A.
- *ramosissima*, Fontaine, 1889, p. 215, pls. xcv-xcix; pl. c, figs. 1-3; pl. ci, fig. 1.
Ibid.
- *varians*, Fontaine, 1894, p. 273, pl. xl, figs. 1, 2; pl. xli, figs. 1-3 a.
Trinity Division; Texas, U.S.A.
- Fritschia nobilis*, Velenovsky, 1885, p. 8, pl. iii, figs. 1-3, 6, 11.
Pläner; Bohemia.
- Fucoides æqualis*, Brongniart, 1824, pp. 307, 310, pl. xix, fig. 7; & 1828, p. 58, pl. v, fig. 4. (= *Chondrites æqualis*, Sternberg, 1833.)
Greensand; Vernasque.
- *Brardii*, Brongniart, 1828, p. 77, pl. ii, figs. 8-10 (synonym for *Carpolithes hemlocinus*?, Schlotheim). (= *Caulerpites Brardii*, Sternberg, 1833.)
Cretaceous; France.
- *Brongniarti*, Mantell (*non* Harlan), 1829, p. 204. Chalk; Sussex.
- *caniculatus*, d'Archiac, 1837, p. 159. Lower Greensand; France.
- *cauliformis*, Fritsch, 1883, p. 136, text-fig. 131, p. 137.
Lower Cretaceous; Chorousek, Bohemia.
- ? *columnaris*, Fritsch, 1883, p. 135, text-fig. 130, p. 137. Ibid.
- *cylindricus*, Sternberg, 1825, p. vii, pl. lxviii, fig. 1.
Quadersandstein; Bohemia.
- *dichotomus*, Reich, in litt. (= *Haliserites Reichii*, Sternberg, 1833.)
- ? *dichotomus*, Fritsch (*non* Morris *nee* Reich), 1889, p. 119, text-fig. 167.
Turonian (?); Koschlitz, Bohemia.
- *difformis*, Brongniart, 1824, pp. 307, 310, pl. xix, fig. 6; & 1828, p. 57, pl. v, fig. 5. (= *Chondrites difformis*, Sternberg, 1833.)
Cretaceous (?); Bayonne.
- *friburgensis*, Heer, 1877, p. 143, pl. lvii, fig. 3.
Neocomian; Switzerland.
- *furcatus*, Brongniart, 1824, p. 309, pl. xix, fig. 3; & 1828, p. 62, pl. v, fig. 1. (= *Chondrites furcatus*, Sternberg, 1833.)
Cretaceous; Vernasque.

- Fucoides funiformis**, Fritsch, 1883, p. 135, text-fig. 129 on p. 136.
Lower Cretaceous; Bohemia.
- **intricatus**, Brongniart, 1824, pp. 307, 311, pl. xix, fig. 8; & 1828,
p. 59, pl. v, figs. 6, 7, 8, (= *Chondrites intricatus*, Sternberg, 1833.)
Greensand; France.
- **latifrons**, Heer, 1877, p. 143, pl. lvii, figs. 1-2.
Upper Cretaceous; Switzerland.
- **Lyngbianus**, Brongniart, 1828, p. 82, pl. ii, figs. 20-21. (= *Sargassites Lyngbianus*, Sternberg, 1833.) Cretaceous; Denmark.
- **Orbignianus**, Brongniart, 1824, p. 308, pl. xix, fig. 1; & 1828,
p. 78. (= *Caulerpites Orbignianus*, Sternberg, 1833.)
Cretaceous; France.
- **recurvus**, Brongniart, 1824, p. 309, pl. ii, fig. 67, pl. xix, fig. 4;
& 1828, p. 62, pl. v, fig. 2. (= *Chondrites recurvus*, Sternberg, 1833.)
Greensand; Vernasque.
- ? **strangulatus**, Fritsch, 1883, p. 136, text-fig. 132, p. 137.
Lower Cretaceous; Bohemia.
- **strictus**, Brongniart, 1824, p. 308, pl. xix, fig. 2. (= *Rhomelites strictus*, Sternberg.)
Greensand; Isle d'Aix.
- **Targionii**, Brongniart, 1828, p. 56, pl. iv, figs. 2 & 6. Ibid.
- **tuberculosus**, Brongniart, 1828, p. 54, pl. vii, fig. 5. (= *Laminarites tuberculosus*, Sternberg.) Ibid.
- **sp.**, Jasche, 1853, p. 96, pl. iv, fig. 1. Cretaceous; Saxony.
- Fucus lignitum**, Lesquereux, 1876 c, p. 364; & 1878 b, p. 42, pl. lxi,
figs. 24, 24 a. Montana Formation; Wyoming, U.S.A.
- Galla quercina**, Lesquereux, 1892, p. 58, pl. vii, fig. 2.
Dakota Group; Kansas, U.S.A.
- Geinitzia biformis**, Knowlton, 1900 a, p. 28 (re-naming *Sequoia biformis*, Lesquereux). Montana Formation; Wyoming, U.S.A.
- **cretacea**, Endlicher, 1847 b, p. 281; & Unger, 1853, p. 93,
pl. xxxiv, fig. 6. Cretaceous; Austria.
- **formosa**, Heer, 1871 a, p. 6, pl. i; pl. ii, figs. 9, 10; & 1871 b,
p. 395. Senonian; Quedlinburg.
- **hyperborea**, Heer, 1883 a, p. 16, pl. li, fig. 13.
Patoot Beds; Greenland.
- **Jenneyi**, Fontaine in Ward, 1899 b, p. 676, pl. clxvi, figs. 5-11;
pl. clxvii. Dakota Group; Black Hills, U.S.A.
- **longifolia** (Lesquereux), Knowlton, 1900 a, p. 28 (re-naming
Sequoia longifolia, Lesquereux).
Montana Formation; Wyoming, U.S.A.
- **macrobracteata**, Richter, 1899 b, p. 44. Senonian; Quedlinburg.
- **microcarpa**, Richter, 1905, p. 15, pl. i, figs. 6, 7. Ibid.
- **prisca**, Eichwald, 1865, p. 48, pl. iii, figs. 19 a, b, c (re-naming
Cunninghamites prisca, 1861). Neocomian; Moscow.

- Geinitzia Reichenbachii** (Geinitz), Hollick & Jeffrey, 1909, p. 38, pl. v, figs. 7-10; pl. viii, figs. 3, 4; pl. xvi, figs. 2-4; pl. xvii, figs. 1-4; pl. xviii, figs. 1-4. (= *Araucarites Reichenbachii*, Geinitz, 1842; = *Sequoia Reichenbachii*, Heer, 1868.)
 Raritan Formation; Staten Island, U.S.A.
- **sp.**, Hollick & Jeffrey, 1909, p. 42, pl. viii, figs. 5, 6; pl. xviii, figs. 5, 6; pl. xix, figs. 1, 2. Ibid.
- **sp.**, Newberry, 1873, p. 10. (= *Geinitzia formosa*, Heer.)
 Cretaceous; New Jersey, U.S.A.
- Gelidinium trajectomosanum**, Debey & Ettingshausen, 1859 A, p. 199, pl. iii, fig. 6 h. Senonian; Maestricht.
- Gecnomites tenuirachis**, Lesquereux, 1878 B, p. 117, pl. xi, fig. 1. Laramie Formation; New Mexico, U.S.A.
- **Ungerii**, Lesquereux, 1878 B, p. 118, pl. xi, fig. 2. Ibid.
- Ginkgo? acetaria**, Ward, 1905, p. 551, pl. cviii, fig. 12. Older Potomac Formation; Maryland, U.S.A.
- **arctica**, Heer, 1874 c, p. 261, pl. 807, fig. 7. (= *Baiera arctica*, Heer.) Kome Beds; Greenland.
- **Baynesiana** (Dawson), Knowlton, 1898, p. 110 (re-naming *Salisburyia Baynesiana*, Dn., 1883).
 Upper Cretaceous; Vancouver Island.
- **Jaccardi**, Heer, 1877, p. 146, pl. lviii, fig. 20. Lower Aptian; Switzerland.
- **laramiensis**, Ward, 1885, p. 496, fig. 7. Montana Formation; Wyoming, U.S.A.
- **lepida**, Heer, 1876, p. 62, pl. xii; pl. vii, fig. 7. Kootanie Formation; British Columbia.
- **multinervis**, Heer, 1882, p. 46, pl. v, fig. 1 d; pl. viii, figs. 2 b, 3, 4; pl. ix, fig. 3 b. Atane Beds; Greenland.
- **nana** (Dawson), Knowlton, 1898, p. 111 (re-naming *Salisburyia nana*, Dn., 1886). Kootanie Formation; British Columbia.
- **primordialis**, Heer, 1874 c, p. 261, pl. 807, figs. 8, 9, 10. Atane Beds; Greenland.
- **pusilla** (Dawson), Knowlton, 1898, p. 111 (re-naming *Salisburyia pusilla*, Dn., 1894.) [Knowlton's name invalidated by Heer's Jurassic species, Heer, 1876 c, p. 61.]
 Upper Cretaceous; Vancouver Island.
- **sibirica**, Heer, 1876 c, p. 61, pl. vii, fig. 6; pl. ix, fig. 5 f; pl. xi. Jurassic, recorded from Kootanie Formation, British Columbia.
- **tenuestriata**, Heer, 1882, p. 14, pl. ii, fig. 12 a. Kome Beds; Greenland.
- **sp.** (nuts of), Dawson, 1886, p. 9, pl. ii, fig. 4. Kootanie Formation; British Columbia, Canada.
- Ginkgoeladus Novæ Zeelandiæ**, Ettingshausen, 1887 A, p. 179, pl. vii, fig. 19. Upper Cretaceous; New Zealand.
- Gleditsiophyllum triacanthoides**, Berry, 1910 A, p. 197. Middle Cretaceous; North Carolina, U.S.A.

- Gleichenia acutiloba*, Heer, 1871 A, p. 5, pl. i, figs. 2, 2*b*.
Senonian; Quedlinburg.
- *acutipennis*, Heer, 1874 A, p. 53, pl. x, figs. 12, 13.
Kome Beds; Greenland.
- *comptoniæfolia*, Heer, 1874 A, p. 49, pl. xi, figs. 1, 2. (= *Didymosorus comptoniiifolius*, Debey & Ettingshausen, 1859 B.) Ibid.
- *crenata*, Velenovsky, 1888 B, p. 9, pl. iii, figs. 15-17. (= *Gleichenites crenata*, Engelhardt, 1892 A.)
Perucer Beds; Vyserovic, Bohemia.
- *delawarensis*, Berry, 1907 A, p. 670, text-figs. 3, 3*a*.
Magothy Formation; Delaware, U.S.A.
- *delicatula*, Heer, 1874 A, p. 54, pl. ix, figs. 11 *e, f*; pl. x, figs. 16, 17.
Kome Beds; Greenland.
- *Drechsleriana*, Goeppert, 1865 B, p. 399 [nomen nudum].
Cretaceous; Austria.
- *Giesekiana*, Heer, 1868, p. 78, pl. xliii, figs. 1 *a*, 2 *a*, 3 *a*; pl. xlv, figs. 2, 3.
Kome Beds; Greenland.
- ? *Gilbert-Thompsoni*, Fontaine in Ward, 1905, p. 232, pl. lxvi, fig. 11.
Shasta Formation; California, U.S.A.
- *gracilis*, Heer, 1871, p. 1181; & 1874 A, p. 52, pl. x, figs. 1-11.
Kome Beds & Atane Beds; Greenland.
- *cf. gracilis*, Heer in Zeiller, 1905, p. 330, pl. vii, figs. 5, 5*a*.
Senonian; Bulgaria.
- *Kurriana*, Heer, 1869 A, p. 6, pl. ii, figs. 1-4. (= *Mertensia Kurriana*, Engelhardt, 1892 A.) Cenomanian; Moletain, Moravia.
- *longipennis*, Heer, 1871, p. 1181; & 1874 A, p. 46, pl. vi, figs. 4, 5, 6; pl. viii, figs. 1, 2, 3.
Kome Beds; Greenland.
- *micromera*, Heer, 1874 A, p. 55, pl. x, figs. 14, 15. Ibid.
- *multinervosa*, Velenovsky, 1888 B, p. 8, pl. iii, figs. 1, 2.
Perucer Beds; Bohemia.
- *Nauckhoffii*, Heer, 1874 A, p. 90, pl. xxv, fig. 4.
Atane Beds; Greenland.
- *nervosa*, Heer, 1874 A, p. 53, pl. xi, figs. 3-6.
Kome Beds; Greenland.
- *Nordenskiöldi*, Heer, 1874 A, p. 50, pl. ix, figs. 6-12. Ibid.
- (*Mertensia*) *obscura*, Ettingshausen, 1887 A, p. 175, pl. vii, figs. 7, 7*a*.
Upper Cretaceous; New Zealand.
- *obtusata*, Heer, 1882, p. 37, pl. xxx, figs. 7-16.
Atane Beds; Patoot, Greenland.
- *optabilis*, Heer, 1880 B, p. 5, pl. i, fig. 13. Kome Beds; Greenland.
- *protogæa*, Debey & Ettingshausen, 1859 B, p. 191, pl. i, figs. 11-12, *g, h*.
Senonian; Aix, Rhenish Prussia.
- *rhombifolia*, Hollick, 1902, p. 147, pl. iii, fig. 3.
Laramie Formation; Colorado, U.S.A.
- *rigida*, Heer, 1868, p. 80, pl. xlv, figs. 1-1*b*.
Kome Beds; Kome, Greenland.
- *Rinkiana*, Heer, 1868, p. 80, pl. xliii, fig. 6. Ibid.
- *rotula*, Heer, 1874 A, p. 48, pl. viii, figs. 4-5; pl. ix, figs. 1-4. Ibid.

- Gleichenia Saundersii**, Berry, 1903 c, p. 679, text figs. 1-3, p. 678.
Matawan Formation; New Jersey, U.S.A.
- **thulensis**, Heer, 1874 A, p. 48, pl. x, fig. 18; pl. v, fig. 9 b.
Kome Beds; Pattofsk, Greenland.
- **Vahlia**, Heer, 1883 A, p. 7, pl. xlix, figs. 8 a, 9.
Patoot Beds; Patoot, Greenland.
- **vidovlensis**, Marik, 1901, p. 1, pl. i, figs. 1, 2.
Cenomanian; Bohemia.
- **votrubensis**, Bayer, 1899, p. 22, pl. ii, fig. 1, text-figs. 7, 7 a.
Perucer Beds; Bohemia.
- **Zippei** (Corda), Heer, 1868, p. 79, pl. xliii, fig. 4. (= *Pecopteris Zippei*, Corda; = *Gleichenites Zippei*, Seward, 1910; = *Mertensia Zippei*, Engelhardt, 1892 A.)
Kome Beds; Greenland.
- **Zippei** (Corda), Zeiller, 1905, p. 328. (*Pecopteris*).
Upper Cretaceous; Bulgaria.
- ? sp., Bayer in Fritsch & Bayer, 1901, p. 85, text-fig. 33. Bohemia.
- Gleichenites coriaceus**, Marik, 1901, p. 7, pl. i, fig. 16.
Cenomanian; Bohemia.
- **crenata**, Engelhardt, 1892 A, p. 82. (= *Gleichenia crenata*, Velenovsky, 1888 B.)
Cretaceous; Bohemia.
- **Zippei** (Corda), Seward, 1910, p. 354 (re-naming *Gleichenia Zippei*, Heer, 1868).
Kome Beds; Greenland.
- Glossozamites acuminatus**, Yokoyama, 1906, p. 38, pl. xii, figs. 5 b, 7.
Cretaceous; China.
- **brevior**, Saporta, 1890 A, p. 814; & 1894, p. 88, pl. xvi, fig. 32.
Valanginian; Portugal.
- **dilaceratus**[-um], Saporta, 1890 A, p. 814.
Ibid.
- **distans**, Fontaine, 1889, p. 176, pl. lxxviii, fig. 5.
Potomac Formation; Virginia, U.S.A.
- **Fontaineanus**, Ward, 1899 B, p. 667, pl. clxii, figs. 16-18.
Lower Cretaceous; Black Hills, U.S.A.
- **Hoheneggeri** (Schenk), Yokoyama, 1906, p. 36, pl. xii, figs. 1, 1 a; pl. xii, figs. 5 a, 6. (= *Podozamites Hoheneggeri*, Schenk.)
Cretaceous; China.
- **Klipsteini** (Dunker), Fontaine in Diller & Stanton, 1894, p. 450. (= *Cyclopteris Klipsteini*, Dunker, 1846.)
Horsetown Beds; California, U.S.A.
- **laceratus**, Saporta, 1894, p. 89, pl. xvi, figs. 26-27.
Valanginian; Portugal.
- **modestior**, Saporta, 1894, p. 88, pl. xvi, fig. 30.
Ibid.
- **parvifolius**, Yokoyama, 1894, p. 226, pl. xxi, figs. 5, 5 a.
Neocomian; Yuasa, Japan.
- **Schenkii**, Heer, 1874 A, p. 99, pl. xvi, figs. 5-8.
Kome Beds; Greenland.
- Glyptostrobus australis**, Ettingshausen, 1893, p. 147; & 1895, p. 12, pl. i, figs. 11-13.
Cretaceous; Australia.
- **brockense** (Fontaine), Ward, 1895 A, p. 359 (re-naming *Taxodium brockense*, Fontaine, 1889).
Potomac Formation; Virginia, U.S.A.

- Glyptostrobus brookense angustifolium** (Fontaine), Knowlton, 1898, p. 112 (re-naming *Taxodium brookense angustifolium*, Fontaine, 1889). Potomac Formation; White House Bluff, U.S.A.
- **cenomanensis**, Crié, 1884, p. 512 [nomen nudum].
Cenomanian; France.
- **europæus cretaceus**, Velenovsky, 1885, p. 26, pl. vi, fig. 2;
pl. vii, figs. 2, 3, 9, 10. Perucur Beds; Bohemia.
- **expansus**, Ward, 1905, p. 538. (= *Taxodium expansum*, Fontaine, 1889.)
- **fastigiatus** (Fontaine), Ward, 1895 A, p. 380 (re-naming *Taxodium fastigiatum*, Fontaine, 1889). Potomac Formation; Virginia, U.S.A.
- **gracillimus**, Lesquereux, 1868, p. 92; & 1874, p. 52, pl. i, figs. 3, 8, & 11-11 f. Dakota Group; Nebraska, U.S.A.
- **grönlandicus**, Heer, 1874 A, p. 76, pl. xvii, fig. 9; pl. xx, figs. 9, 10, 11; pl. xxii, fig. 12. Kome Beds; Greenland.
- **intermedius**, Heer, 1883 A, p. 13, pl. lii, figs. 6-7.
Patoot Beds; Greenland.
- **sp.**, Dawson, 1883, p. 25.
Upper Cretaceous; Vancouver Island, Canada.
- **? sp.**, Knowlton, 1900 A, p. 31, pl. v, fig. 4.
Montana Formation; Wyoming, U.S.A.
- Granularia sp.**, T. Lorenz, 1902, p. 56, pl. vii, fig. 2.
Lower Cretaceous; Switzerland.
- Grevillea constans**, Velenovsky, 1882 A, p. 213; & 1883, p. 28, pl. ix, figs. 6-10. Cenomanian; Bohemia.
- **Dvraki**, Bayer, 1899, p. 27, pl. i, fig. 15, text-figs. 9, 9 a. Ibid.
- **Oxleyana**, Ettingshausen, 1893, pp. 140, 149; & 1895, p. 27, pl. iii, fig. 14. Cretaceous; Australia.
- **palmata**, Debey MS. in Saporta & Marion, 1873, p. 61, pl. viii, figs. 5-7. (= *Devalquea aquisgranensis*, Saporta & Marion.)
Senonian; Aix, Rhenish Prussia.
- **Reussii**, Ettingshausen, 1851, p. 721 (re-naming *Salicites angustus*, Reuss). Cretaceous; Bohemia.
- **tenera**, Velenovsky, 1887, p. 72, pl. xxx, figs. 9, 14, 16.
Perucur Beds; Bohemia.
- Grewiopsis æquidentata**, Lesquereux, 1892, p. 180, pl. lviii, fig. 4.
[Same as *G. Mudgei*.] Dakota Group; Kansas, U.S.A.
- **Cleburni**, Lesquereux, 1876 A, p. 381; & 1878 B, p. 259, pl. lxii, fig. 12. Laramie Formation; Wyoming,
& Point of Rocks, Montana, U.S.A.
- **flabellata** (Lesquereux), Knowlton, 1898, p. 114 (re-naming *Populites flabellata*, Lx., 1868). Dakota Group; Kansas, U.S.A.
- **Haydenii**, Lesquereux, 1874, p. 97, pl. iii, figs. 2, 4; pl. xxiv, fig. 3.
[Same as *G. flabellata*.] Ibid.
- **Mudgei**, Lesquereux, 1892, p. 181, pl. lxvi, fig. 3. [This is founded on the same specimen as *G. æquidentata*.] Ibid.
- **pakawanica**, Ettingshausen, 1887 A, p. 187, pl. ix, fig. 21.
Upper Cretaceous; New Zealand.

- Grewiopsis paliurifolia**, Ward, 1887, p. 92, pl. lvi, fig. 3.
Laramie Formation; Wyoming, U.S.A.
- **viburnifolia**, Ward, 1887, p. 89, pl. xl, fig. 2; & recorded Hollick,
1894 A, p. 59, pl. clxxiv, fig. 8. Cretaceous; Long Island, U.S.A.
- Guatteria cretacea**, Hollick, 1906 A, p. 73, pl. xxi, figs. 1-4.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Gyminda primordialis**, Hollick, 1906 A, p. 88, pl. xxxiii, fig. 5. Ibid.
- Gymnogramme bohemia**, Bayer, 1899, p. 45, pl. ii, fig. 2, text-figs.
14, 14 a. Perucur Beds; Bohemia.
- **Haydenii**, Lesquereux, 1872, p. 295; & 1878, p. 59, pl. v, figs. 1-3.
Laramie Formation; Wyoming, U.S.A.
- Gyrochorte porrecta**, De Stefani, 1881, p. 281.
Upper Cretaceous; Italy.
- Gyrolithes Davreuxi**, Saporta, 1884, p. 27, pl. v, figs. 1-4'; pl. vi,
figs. 1, 2. Greensand; France.
- Gyrophyllites angustifolius**, Heer, 1865, p. 190, text-fig. 100 b.
Cretaceous; Switzerland.
- **Doblhoffii**, Lorenz, 1901, p. 579, pl. iv, fig. 11. Flysch; Austria.
- **Kastneri**, Lorenz, 1901, p. 579, pl. iv, figs. 5, 7-8. Ibid.
- **Kwassizensis**, Glocker, 1841, p. 322, text-fig.
Quadersandstein; Austria.
- **obtusifolius**, Heer, 1865, p. 190, text-fig. 101; & 1877, p. 144,
pl. lviii, fig. 13. Cretaceous; Switzerland.
- **Oosteri**, Heer, 1865, p. 190, text-fig. 100 a; & 1877, p. 144, pl. lviii,
fig. 14. Neocomian; Switzerland.
- **pentamerus**, Heer, 1865, p. 190, text-fig. 102; & 1877, p. 144,
pl. lviii, fig. 15. Ibid.
- **Petteri**, Lorenz, 1901, p. 579, pl. iv, figs. 9-10. Flysch; Austria.
- **quassazensis**, Glocker in Goepfert, 1865 c, p. 12. (= *Gyrophyllites*
Kwassizensis, Glocker, 1841.)
- **vestanensis**, Squinabol, 1890, p. 189, pl. xi, fig. 2.
Cretaceous (?); Italy.
- Haastia speciosa**, Ettingshausen, 1887 A, p. 180, pl. viii, fig. 5.
Upper Cretaceous; New Zealand.
- Halimeda Fuggeri**, Lorenz, 1897, p. 177, pl. i. Flysch; Salzburg.
- Haliserites contortuplicatus**, von der Mark, 1864, p. 81, pl. xiii,
fig. 13. Upper Cretaceous; Westphalia.
- **gracilis**, Debey & Ettingshausen, 1859 A, p. 189, pl. i, figs. 1-2.
Senonian; Aix, Rhenish Prussia.
- **Reichii**, Sternberg, 1833, p. 34, pl. xxiv, fig. 7. (= *Fucoides*
dichotomus, Reich in litt.) Greensand; Freiberg.
- **Schlotheimi**, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **trifidus**, Debey, 1848 A, p. 114 [nomen nudum]. Ibid.
- Halocharis longifolia**, Miquel, 1853, p. 49, pl. v, figs. 4-8.
Senonian; Limburg, Belgium.

- Halymenites cylindricus**, Sternberg, 1825, pl. xlviii, fig. 1.
(= *Münsteria cylindrica*, Otto.) Quadersandstein; Bohemia.
- **major**, Lesquereux, 1873, p. 373; & 1878 B, p. 38, pl. i, figs. 7, 8.
Montana Formation (& above); U.S.A.
- **Rioana**, Zigno, 1864, p. 525 [nomen nudum]. Cretaceous (?); Italy.
- **striatus**, Lesquereux, 1873, p. 373; & 1878 B, p. 37, pl. i, fig. 6.
Laramie Formation; New Mexico, U.S.A.
- **verticillatus**, Münster (*in litt.*), Sternberg, 1838, p. 104.
Cretaceous; Westphalia.
- Halyserites gracilis**, Debey & Ettingshausen, 1859 A, p. 189, pl. i, figs. 1, 2. Senonian; Aix, Rhenish Prussia.
- **Reichii** (Sternberg), Unger, 1850 A, p. 10. (= *Halyserites Reichii*, Sternberg, = *Delesseria Reichii*, Engelhardt, 1892 A.)
Senonian; Aix, Rhenish Prussia.
- **Schlotheimi**, Debey, 1848 (?); & 1850, p. 116. Ibid.
- **trifidus**, Debey, 1848 A, p. 114. Ibid.
- Hamamelidoxylon Renaulti**, Lignier, 1907, p. 301, pl. xix, fig. 44; pl. xx, figs. 45-52; pl. xxi, fig. 68; pl. xxiii, figs. 85, 93; text-figs. 3-5, pp. 294-296. Cenomanian; France.
- Hamamelites ? cordatus**, Lesquereux, 1883, p. 71, pl. iv, fig. 3.
Dakota Group; Kansas, U.S.A.
- **kansaseana**, Lesquereux, 1876, p. 355, pl. vii, fig. 4. (= *H. kansaseanus*, 1883, = *Alnus kanseana*, 1874, re-named by Knowlton, 1898, *Quercus kanseana*.) Ibid.
- **quadrangularis**, Lesquereux, 1876 B, p. 355. (= *Alnites quadrangularis*, Lx., 1874, & *Populites quadrangularis*, Lx., 1868.) Ibid.
- **quercifolius**, Lesquereux, 1883, p. 71. Ibid.
- **tenuinervis**, Lesquereux, 1883, p. 70. Ibid.
- Hausmannia ? arctica** (Heer), Richter, 1906, p. 20. (= *Jeanpaulia arctica* & *Jeanpaulia borealis*, Heer, 1874 A.) Urgonian; Greenland.
- **? Brongniarti** (Debey & Ettingshausen), Richter, 1906, p. 19, pl. vii, fig. 23. (= *Asplenium Brongniarti*, Debey & Ettingshausen, 1859 B.) Senonian; Quedlinburg, Saxony.
- **? californica**, Fontaine in Ward, 1905, p. 238, pl. lxx, fig. 47.
Shasta Formation; California, U.S.A.
- **? cordata** (Heer), Richter, 1906, p. 24, pl. vii, fig. 19. (= *Protorhipis cordata*, Heer, 1882.) Kome Beds; Greenland.
- **cretacea** (Velenovsky), Richter, 1906, p. 20. (= *Platyseriphyllum cretaceum*, Velenovsky, 1889.) Cenomanian; Bohemia.
- **dichotoma**, Dunker, 1846, p. 12, pl. v, fig. 1; pl. vi, fig. 12.
Recorded from Senonian, Quedlinburg, by Richter, 1906.
- **dichotoma euryphylla**, Richter, 1906, p. 19, pl. iii, fig. 11; pl. iv, figs. 2, 4, 5, 7; pl. vi, fig. 2. Senonian; Quedlinburg, Saxony.
- **dichotoma linearis**, Richter, 1906, p. 19, pl. iii, figs. 5, 8; pl. iv, figs. 6, 9, 9 a. Ibid.
- **dichotoma regularis**, Richter, 1906, p. 19, pl. iii, figs. 3, 4, 6, 10; pl. iv, figs. 1, 8; pl. v, figs. 9, 10; pl. vi, fig. 5. Ibid.
- **gracillima**, Richter, 1906, p. 19, pl. iii, figs. 12, 12 a. Ibid.

- Hausmannia Kohlmanni**, Richter, 1901, p. 21; & 1906, p. 21, pl. i, figs. 1-11; pl. ii, figs. 1, 3, 4, 5, 6, 8, 9; pl. v, figs. 1, 2, 5, 6, 7, 8; pl. vi, figs. 3, 6, 7, 9. Senonian; Quedlinburg, Saxony.
- **rigida**, Newberry, 1895, p. 35, pl. i, figs. 2, 3, 5. Amboy Clay; New Jersey, U.S.A.
- **Sewardi**, Richter, 1906, p. 22, pl. i, fig. 12; pl. v, figs. 3, 4; pl. vi, fig. 8. Lower Cretaceous; Strohberg.
- **spuria**, Richter, 1906, p. 23, pl. ii, fig. 2 (pl. v, fig. 12 (?)). Ibid.
- Hedera aquamara**, Ward, 1887, p. 59, pl. xxvi, fig. 7 (re-named *Quercus aquamara* (Ward), Knowlton, 1898). Laramie Formation; Wyoming, U.S.A.
- **Bruneri**, Ward, 1887, p. 58, pl. xxvi, fig. 6. Ibid.
- **cecilensis**, Berry, 1910 n, p. 28, pl. viii, fig. 2. Magothy Formation; Cecil County, U.S.A.
- **credneriæfolia**, Velenovsky, 1882 n, p. 30, pl. viii, fig. 8; pl. x, fig. 2. (= *Hederophyllum credneriæfolium*, Velenovsky, 1889.) Cenomanian; Vyserovic, Bohemia.
- **cretacea**, Lesquereux, 1892, p. 127, pl. xviii, fig. 1. Dakota Group; Kansas, U.S.A.
- **cuneata**, Heer, 1882, p. 83, pl. xxviii, fig. 12; pl. xlv, fig. 2. Atane Beds; Greenland.
- **decurrens**, Lesquereux, 1892, p. 130, pl. xviii, fig. 6. Dakota Group; Kansas, U.S.A.
- **microphylla**, Lesquereux, 1892, p. 127, pl. xviii, figs. 2, 3. Ibid.
- **obliqua**, Newberry, 1895, p. 113, pl. xxxvii, fig. 8; pl. xxxviii, fig. 5. Amboy Clay; Woodbridge, U.S.A.
- **orbiculata** (Heer), Lesquereux, 1892, p. 129, pl. xvii, figs. 12-14. (= *Chondrophyllum orbiculatum*, Heer, 1874 a.) Dakota Group; Kansas, U.S.A.
- **ovalis**, Lesquereux, 1874, p. 91, pl. xxv, fig. 3; pl. xxvi, fig. 4. Dakota Group; Nebraska, U.S.A.
- **platanoidea**, Lesquereux, 1876 n, p. 351, pl. iii, fig. 3. Dakota Group; Kansas, U.S.A.
- **primordialis**, Saporta, 1879, p. 200, text-figs. 29-1 & 2. (= *Hederophyllum primordiale*, in Velenovsky, 1889.) Cenomanian; Bohemia.
- **Schimper**, Lesquereux, 1876 a, p. 395; & 1876 n, p. 351, pl. vii, fig. 5. Dakota Group; Kansas, U.S.A.
- **simplex**, Hollick, 1906 a, p. 97, pl. xxxvii, fig. 9. Middle Cretaceous; Martha's Vineyard, U.S.A.
- **sp.?**, Hollick, 1898 n, p. 421, pl. xxxviii, fig. 5. Middle Cretaceous; Staten Island, U.S.A.
- Hederophyllum angulatum**, Fontaine, 1889, p. 324, pl. clxii, fig. 1. Potomac Formation; Virginia, U.S.A.
- **crenulatum**, Fontaine, 1889, p. 324, pl. clxii, fig. 3. Ibid.
- **peltatum**, Marik, 1901, p. 11, pl. ii, fig. 8. Cenomanian; Bohemia.

- Hederophyllum credneriaefolium**, Velenovsky, 1889, pp. 50, 54.
(= *Hedera credneriaefolia*, Velenovsky, 1882 B.)
- **primordiale** (Saporta), Velenovsky, 1889, pp. 50, 54. (= *Hedera primordialis*, Saporta, 1879.)
- Heterofilicites anceps**, Berry, 1906 B, p. 154, pl. xxvi.
Magothy Formation; New Jersey, U.S.A.
- Hexagonaria senonica**, Deecke, 1901, p. 473. Senonian; Rügen.
- Himantites alopecarus**, Debey & Ettingshausen, 1859 A, p. 212, pl. iii, fig. 1 a. Senonian; Aix, Rhenish Prussia.
- Hymenæa dakotana**, Lesquereux, 1892, p. 145, pl. lv, figs. 2, 3; pl. lvi, figs. 1, 2; pl. lxii, fig. 2. Dakota Group; Kansas, U.S.A.
- **elongata**, Velenovsky, 1882 A, p. 214; & 1886, p. 57, pl. xx, figs. 3, 5. Cenomanian; Bohemia.
- **inæqualis**, Velenovsky, 1886, p. 56, pl. xxi, figs. 5, 6. (= *Hymenæophyllum inæqualis*, Velenovsky, 1889.) Ibid.
- **Kuchlensis**, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- **latifolia**, Velenovsky, 1882 A, p. 214 [nomen nudum]. Ibid.
- **primigenia** [spelt **Hymeneæa**], Saporta, 1879, p. 199, text-fig. 28, 2; also Velenovsky, 1886, p. 56, pl. xx, fig. 4; pl. vi, figs. 1-4. Ibid.
- **virginiensis**, Fontaine, 1889, p. 320, pl. clxiii, fig. 6.
Potomac Formation; Virginia, U.S.A.
- Hymenæophyllum inæquale**, Velenovsky, 1889, p. 51. (= *Hymeneæa inæqualis*, Velenovsky, 1886.)
- Hymenocphyllites heterophyllus**, Unger, 1850 A, p. 527; & 1867, p. 650, pl. ii, figs. 3, 4. Upper Cretaceous; Austria.
- **macrocphyllus**, Goeppert, 1836, p. 262; & Unger, 1867, p. 650, pl. ii, fig. 5. Ibid.
- Hymenophyllum cretaceum**, Lesquereux, 1873, p. 421; & 1874, p. 45, pl. i, figs. 3-4; pl. xxix, fig. 6.
Dakota Group; Kansas, U.S.A.
- Hymenopteris psilotoides**, Stokes & Webb, 1824, p. 424, pl. xlv, fig. 7; pl. xlvii, fig. 2; Mantell, 1827, p. 55, pl. i, fig. 3; pl. iii, fig. 7; pl. iii*, fig. 2; pl. xx, figs. 1, 2. (= *Sphenopteris Mantelli*, Brongniart, 1828.) Wealden, Sussex.
- Hypoglossidium antiquum**, Heer, 1874 A, p. 129, pl. xxxviii, figs. 14, 15. Cretaceous; Spitzbergen.
- Hysterites dubius**, Debey & Ettingshausen, 1859 A, p. 213, pl. iii, fig. 5 d. Senonian; Aix, Rhenish Prussia.
- **protogæus** (Heer), Meschinelli, 1892, p. 774. (= *Hysterium protogæum*, Heer, 1882.)
- Hysterium protogæum**, Heer, 1882, p. 20, pl. xxiv, figs. 9, 9 b. (= *Hysterites protogæus*, Meschinelli, 1892.)
Atane Beds; Greenland.
- Ilex amboyensis** (Newberry), Berry, 1909, p. 259 (re-naming *Ilex ovata*, Newberry, 1896.) Raritan Formation; New Jersey, U.S.A.

- Ilex antiqua*, Heer, 1882, p. 97, pl. xxvii, fig. 1 a.
Atane Beds; Greenland.
- *armata*, Lesquereux, 1892, p. 176, pl. xxix, fig. 8.
Dakota Group; Kansas, U.S.A.
- *borealis*, Heer, 1883 A, p. 39, pl. lxiv, figs. 3, 4.
Patoot Beds; Greenland.
- *dakotensis*, Lesquereux, 1892, p. 178, pl. xxix, fig. 11.
Dakota Group; Kansas, U.S.A.
- ? *elongata*, Newberry, 1895, p. 98, pl. xviii, figs. 1, 5.
Amboy Clay; Sayreville, U.S.A.
- *Masoni*, Lesquereux, 1892, p. 179, pl. vii, fig. 6; pl. lxiii, fig. 6.
Dakota Group; Kansas, U.S.A.
- ? *ovata*, Newberry, 1895, p. 98, pl. xviii, fig. 2 (re-named *Ilex amboyensis*, Berry, 1909.)
Amboy Clay; Sayreville, U.S.A.
- *papillosa*, Lesquereux, 1892, p. 177, pl. xxix, figs. 9, 10; pl. lviii, fig. 3.
Dakota Group; Kansas, U.S.A.
- *patootensis*, Heer, 1883 A, p. 40, pl. lxiv, fig. 5.
Patoot Beds; Greenland.
- *Pernerii*, Bayer, 1893, pp. 20 & 42, fig. 14; & Bayer in Fritsch, 1893, p. 131, text-fig. 190.
Senonian; Priesen, Bohemia.
- *Scudleri*, Lesquereux, 1892, p. 178, pl. lviii, fig. 2.
Dakota Group; Kansas, U.S.A.
- *strangulata*, Lesquereux, 1876 A, p. 399; & 1876 B, p. 359, pl. viii, fig. 3.
Dakota Group (?); Colorado, U.S.A.
- Illicium deletum*, Velenovsky, 1886, p. 51, pl. xviii, fig. 5.
Cenomanian; Bohemia.
- Inga Cottai*, Ettingshausen, 1867 A, p. 262, pl. iii, fig. 18.
Cenomanian; Niederschoena, Saxony.
- *cretacea*, Lesquereux, 1892, p. 153, pl. lv, fig. 11.
Dakota Group; Kansas, U.S.A.
- *latifolia*, Velenovsky, 1886, p. 55, pl. xx, figs. 6, 7.
Cenomanian; Vyserovic, Bohemia.
- Inolepis affinis*, Heer, 1883 A, p. 11, pl. liii, figs. 2, 2 b.
Patoot Beds; Greenland.
- *bohemica*, Marik, 1901, p. 8, pl. i, fig. 19.
Cenomanian; Bohemia.
- *imbricata*, Heer, 1874 A, p. 72, pl. xvi, figs. 12-16; pl. xxiii, figs. 6 c, 7, 8.
Kome Beds; Greenland.
- Isoetes Choffati*, Saporta, 1894, p. 134, pl. xxiv, figs. 2 b, 9-11; pl. xxv, figs. 5-8; pl. xxvii, fig. 6.
Urgonian; Portugal.
- Isoetopsis Choffati*, Saporta, 1891, p. 250 [nomen nudum].
Albian; Portugal.
- Jeanpaulia arctica*, Heer, 1874 A, p. 131, pl. ii, figs. 15 & 16, name in pl. descr. given to *J. borealis*. (= *Hausmannia? arctica*, in Richter, 1906.)

- Jeanpaulia borealis*, Heer, 1871, p. 1181; & 1874 A, p. 57, pl. ii, figs. 15 & 16. (= *Hausmannia? arctica*, in Richter, 1906.)
Kome Beds; Greenland.
- *carinata*, Velenovsky, 1888 B, p. 18, pl. i, figs. 1-5.
Perucer Beds; Bohemia.
- *grandis*, Heer, 1871, p. 1181 [nomen nudum].
Kome Beds; Greenland.
- *lepida*, Heer, 1874 A, p. 58, pl. ii, figs. 1-4. (= *Sphenopteris lepida*, Heer, 1871.)
Ibid.
- Jordania ebenoides*, Schenk, 1883, p. 10, pl. iv, figs. 13-14.
Upper Cretaceous; Libyan Desert.
- Juglandinium longiradiatum*, Vater, 1884, p. 841, pl. xxviii, figs. 15-16.
Lower Senonian; Brunswick.
- Juglandiphyllum integrifolium*, Fontaine, 1889, p. 315, pl. clvii, figs. 3, 5, 6.
Potomac Formation; Virginia, U.S.A.
- Juglandites cretacea*, Dawson, 1886, p. 14.
Mill Creek Series; Canada.
- *elegans*, Goeppert, 1842 B, p. 156, pl. liv, fig. 18.
Senonian; Aix, Rhenish Prussia.
- *ellsworthianus*, Lesquereux, 1892, p. 70, pl. xxxvii, fig. 1.
Dakota Group; Kansas, U.S.A.
- *fallax*, Dawson, 1894, p. 59, pl. xi, fig. 48.
Upper Cretaceous; Vancouver Island, Canada.
- *Lacoei*, Lesquereux, 1892, p. 71, pl. xlviii, fig. 5.
Dakota Group; Kansas, U.S.A.
- *primordialis*, Lesquereux, 1892, p. 70, pl. xxxv, fig. 15.
Ibid.
- *sinuatus*, Lesquereux, 1892, p. 71, pl. xxxv, figs. 9-11.
Ibid.
- ? *sp.*, Dawson, 1894, p. 59, pl. x, fig. 43; pl. xiii, fig. 56.
Upper Cretaceous; Vancouver Island, Canada.
- Juglans arctica*, Heer, 1882, p. 71, pl. xl, fig. 2; pl. xli, fig. 4c; pl. xlii, figs. 1-3; pl. xliii, fig. 3.
Atane Beds; Greenland.
- *crassipes*, Heer, 1869 A, p. 23, pl. vi, fig. 3.
Cenomanian; Moletuin, Moravia.
- *Debeyana* (Heer), Lesquereux, 1868, p. 101. (= *Populus? Debeyana*, Heer, 1867.)
Dakota Group; Nebraska, U.S.A.
- *elongata*, Hollick, 1906 A, p. 55, pl. xi, figs. 3, 4 (for *Laurus Omalii*, Sap. & Mar.).
Middle Cretaceous; Long Island, U.S.A.
- *harwoodensis*, Dawson, 1883, p. 28, pl. viii, fig. 31.
Upper Cretaceous; Vancouver Island, Canada.
- *Humboldtana*, Stiehler, 1858, p. 63, pl. xi, fig. 13.
Cretaceous; New Granada.
- *Leconteana*, Lesquereux, 1871, p. 382; & 1878 B, p. 285, pl. liv, figs. 10-13.
Laramie Formation; Colorado, U.S.A.
- ? *missouriensis*, Knowlton, 1900 A, p. 12, pl. i, figs. 7-9.
Montana Formation; Missouri R., U.S.A.
- *ramnoides*, Lesquereux, 1872, p. 294.
Laramie Formation; Wyoming, U.S.A.

Juglans rugosus, Lesquereux, 1868, p. 206; & 1878 b, p. 286, pl. liv, figs. 4, 5; pl. lv, figs. 1-9; pl. lvi, figs. 1, 2.

Laramie Formation; Wyoming, U.S.A.

Jugloxylon hamacanam, Stopes & Fujii, 1910, pp. 62-63, pl. vii, fig. 48. Upper Cretaceous; Hokkaido, Japan.

Jungermannites vetustior, Saporta, 1894, p. 123, pl. xxxiii, fig. 14. Urgonian; Cereal, Portugal.

Juniperus hypnoides, Heer, 1882, p. 47, pl. xlv, figs. 3-4; pl. xlv, fig. 18. Atane Beds; Greenland.

— *macilenta*, Heer, 1882, p. 47, pl. xxxv, figs. 10-11. Ibid.

Juranyia hemiflabellata, Tuzson, 1908, p. 1, pl. i, figs. 1, 2; pl. ii, fig. 3. Upper Cretaceous; Hungary.

Kaidacarpium cretaceum, Heer, 1883 a, p. 19, pl. lxiv, fig. 9 b.

Patoot Beds; Greenland.

Kaidacarpum Bucklandi (Lindley & Hutton), Carruthers, 1868, p. 156, pl. ix, fig. 2. Upper Greensand; Wiltshire.

— *minus*, Carruthers, 1868, p. 156. Potton Sands; Cambridge.

Kalmia Brittoniana, Hollick, 1893, p. 34, pl. ii, figs. 6, 7, 8. Cretaceous; New York, U.S.A.

Keckia ambigua, Eichwald, 1865, p. 11, pl. iv, figs. 14, 15.

Neocomian; Russia.

— *annulata*, Glocker, 1841, p. 319, pl. iv. (= *Münsteria annulata*, Unger.) Quadersandstein; Austria.

— *cylindrica*, Otto, 1852, p. 5, pl. ii; pl. iii; pl. iv, fig. 2. Lower Quadersandstein; Wendischcarsdorf.

— *nodulosa*, Otto, 1852, p. 9, pl. iv, figs. 3, 6, 7. Ibid.

— *vesiculosa*, Otto, 1852, p. 8, pl. iv, fig. 1. Ibid.

— *sp.*, Lorenz, 1902, p. 23, pl. viii, fig. 3. Lower Cretaceous; Switzerland.

Kingthiophyllum primævum, Crie, 1889, p. 15 [nomen nudum].

Cretaceous; New Zealand.

Kirchnera arctica (Heer), Velenovsky, 1888 b, p. 16, pl. ii, figs. 12-16. Perucr Beds; Bohemia.

— *dentata*, Velenovsky, 1888 b, p. 18, pl. ii, figs. 1, 2. Ibid.

Kloedenia quercoides, Goepfert, 1839, p. 521, pl. viii b, figs. 1, 3, 4. Cretaceous; Silesia.

Knightiophyllum primævum, Ettingshausen, 1887 a, p. 185, pl. ix, fig. 12. Upper Cretaceous; New Zealand.

Kohlmannopteris insignis, Richter, 1899 a, p. 40 [nomen nudum]; & 1901, p. 20. Senonian; Quedlinburg.

Kosmogyra acanthica, Stache, 1889, p. 135, pl. iv, fig. 5. (= *Chara acanthica*, Stache, 1880.) Upper Cretaceous (?); Cognale.

— *cingulata*, Stache, 1889, p. 121, pl. ii, fig. 20. (= *Chara cingulata*, Stache, 1880.) Liburnian (Upper Cretaceous); Cosina.

— *guttifera*, Stache, 1889, p. 134, pl. iv, fig. 6. (= *Chara guttifera*, Stache, 1880.) Ibid.

— *ornata*, Stache, 1889, p. 134, pl. iv, figs. 4, a, b. (= *Chara ornata*, Stache, 1880.) Ibid.

Kosmogyræ perarmata, Stache, 1889, p. 134, pl. iv, figs. 3 a, b.
(= *Chara perarmata*, Stache, 1880.)

Liburnian (Upper Cretaceous); Cosina.

— **superba**, Stache, 1889, p. 134, pl. iv, figs. 2 a, b. (= *Chara superba*,
Stache, 1880.) Upper Cretaceous (?); Divacea.

Kcsmogyrella carinata, Stache, 1889, p. 121, pl. ii, fig. 19. (= *Chara*
carinata, Stache, 1880.) Ibid.

Kranneria mirabilis, Corda (*in litt.*), Velenovsky, 1885, p. 1, pl. i,
figs. 1-7, 10-13; pl. iv, figs. 1, 2, 4, 7, 8, 9. (= *Palæostrobus*
mirabilis, Renger, 1866.) Cenomanian; Bohemia.

Laccopteris Dunkeri, Schenk, 1871 A, p. 17, pl. viii, figs. 3-5; Velenov-
sky, 1888, p. 12, pl. ii, figs. 3-7. Cenomanian; Bohemia.

— **pulchella**, Heer, 1831, p. 16, pl. xv, figs. 7, 7 b.
Neocomian; Portugal.

Lagynophora articulata, Stache, 1889, p. 133, pl. vi, fig. 5.
Liburnian (Upper Cretaceous); Cosina.

— **folicosa**, Stache, 1889, p. 133, pl. iv, figs. 10, 11. Ibid.

— **liburnica**, Stache, 1889, p. 132, pl. iv, figs. 9, 14. (= *Chara*
liburnica, Stache, 1880.) Ibid.

— **nodulifera**, Stache, 1889, p. 133, pl. vi, fig. 3. Ibid.

— **symmetrica**, Stache, 1889, p. 133, pl. vi, figs. 1, 4. Ibid.

Lambertia dura, Velenovsky, 1883, p. 30, pl. x, fig. 16. (= *Drynaria*
dura (Velenovsky), Bayer, 1899.) Cenomanian; Bohemia.

Laminarites crenatus, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.

— **polystigma**, Debey & Ettingshausen, 1859 A, p. 192, pl. i, fig. 6.
Ibid.

— **spathulatus**, Debey, 1849, p. 299 [nomen nudum]. Ibid.

— **tuberculosus**, Sternberg, 1833, p. 35. (= *Fucoides tuberculosus*,
Brongniart, 1828.) Cretaceous; La Rochelle, France.

Lamprocarpites nitidus, Heer, 1882, p. 58, pl. viii, figs. 10 b, 12-14.
Atane Beds; Greenland.

Laricopsis angustifolia, Fontaine, 1889, p. 233, pl. cii, figs. 9, 10;
pl. ciii, figs. 1, 4. Potomac Formation; Virginia, U.S.A.

— **brevifolia**, Fontaine, 1889, p. 234, pl. cii, figs. 5, 6. Ibid.

— **longifolia**, Fontaine, 1889, p. 233, pl. cii, figs. 7, 8; pl. ciii, figs. 2,
3; pl. clxv, fig. 4; pl. clxviii, figs. 5, 6. Ibid.

— **longifolia latifolia**, Fontaine in Ward, 1905, p. 312, pl. lxxiii,
figs. 11-14. Kootanie Formation; Montana, U.S.A.

Laurelia primæva, Lesquereux, 1892, p. 108, pl. xx, fig. 8.
Dakota Group; Kansas, U.S.A.

Laurinium brunswicense, Vater, 1884, p. 845, pl. xxix, figs. 22-24.
Lower Senonian (?); Helmstedt.

Laurinophyllum austro-caledonicum, Crieé, 1889, p. 27 [nomen
nudum]. Cretaceous; New Caledonia.

Laurinoxylon albiense, Fliche, 1905, p. 356, pl. x, figs. 2-3.
Albian; Madagascar.

- Laurinoxylon primigenium**, Schenk, 1883, p. 11, pl. iii, fig. 10;
pl. v, figs. 15-16. Cenomanian (?); Cairo, Egypt.
- **uniseriatum**, Gothan, 1908 A, p. 16, pl. ii, figs. 3-11.
Cretaceous (?); Antarctica.
- **sp.**, Gothan, 1908 A, p. 19, pl. ii, figs. 12, 13. Ibid.
- Laurogene cretacea**, Ettingshausen in Reuss, 1854, p. 740 [nomen nudum].
Cenomanian; Molettein, Moravia.
- Laurophyllum angustifolium**, Newberry (*non* Lesquereux), 1895,
p. 86, pl. xvii, figs. 10, 11. Amboy Clay; New Jersey, U.S.A.
- **aquisgranense**, Lange, 1890, p. 669, pl. xxxiv, fig. 4.
Senonian; Gemmenich, Rhenish Prussia.
- **austro-caledonicum**, Crié, 1889, p. 27 [nomen nudum].
Cretaceous; New Caledonia.
- **debile**, Dawson, 1883, p. 22, pl. ii, figs. 7, 7 a.
Upper Cretaceous; North-West Territory, Canada.
- **elegans**, Hollick, 1906 A, p. 81, pl. xxvii, figs. 1-5.
Raritan Formation; Staten Island, U.S.A.
- **ellsworthianum**, Lesquereux, 1892, p. 95, pl. xiii, fig. 7. (= *Quercus ellsworthianus*, Lesquereux, 1868.)
Dakota Group; Kansas, U.S.A.
- **insigne**, Dawson, 1894, p. 61, pl. vii, figs. 24, 25.
Upper Cretaceous; Vancouver Island, Canada.
- **lanceolatum**, Newberry, 1895, p. 86, pl. xvii, figs. 1, 12.
Amboy Clay; New Jersey, U.S.A.
- **minus**, Newberry, 1895, p. 86, pl. xvi, figs. 7-9. Ibid.
- **nervillosum**, Hollick, 1906 A, p. 82, pl. xxvii, figs. 6, 7.
Raritan Formation; Staten Island, U.S.A.
- **reticulatum**, Lesquereux, 1873, p. 425; & 1874, p. 76, pl. xv,
figs. 4, 5. (= *Ficus reticulata* (Lx.), Hollick, 1898.)
Dakota Group; Kansas, U.S.A.
- Laurus affinis**, Hosius & von der Marck, 1880, p. 167, pl. xxxi, fig. 9.
Upper Senonian; Haldem, Westphalia.
- **affinis**, Velenovsky (*non* Hosius & von der Marck), 1887, p. 70,
pl. xxviii, figs. 4, 5, 7, 8. Cenomanian; Bohemia.
- **angusta**, Heer, 1882, p. 76, pl. xx, figs. 1 b, 7; pl. xliii, fig. 1 c.
Atane Beds; Greenland.
- **antecedens**, Lesquereux, 1892, p. 92, pl. xi, fig. 3.
Dakota Group; Kansas, U.S.A.
- **attenuata**, Saporta, 1894, p. ?, pl. xxxvii, fig. 9.
Cretaceous (?); Portugal.
- **Colleti**, Fliche, 1892, p. 1085; & 1896, p. 279, pl. xiii, fig. 7, text-
fig. 13. Albian; France.
- **crassinervis**, Dawson, 1883, p. 23.
Upper Cretaceous (?); North-West Territory, Canada.
- **cretacea**, Ettingshausen, 1867 A, p. 252, pl. ii, fig. 13.
Cenomanian; Niederschoena, Saxony.
- **Hollae**, Heer, 1882, p. 76, pl. xxxiii, fig. 13; pl. xlv, fig. 5 b;
pl. xlv, fig. 3. Atane Beds; Greenland.

- Laurus Hollickii**, Berry, 1905 E, p. 79, pl. lii, fig. 4.
Matawan Formation; New Jersey, U.S.A.
- **Knowltoni**, Lesquereux, 1892, p. 94, pl. 1, fig. 4.
Dakota Group; Kansas, U.S.A.
- **labrusca**, Unger, 1850 A, p. 423. (= *Stereulia labrusca*, Unger, 1851 B.)
Dakota Group; Nebraska, U.S.A.
- **macrocarpa**, Lesquereux, 1868, p. 98; & 1874, p. 74, pl. x, fig. 2.
Dakota Group; Nebraska, U.S.A.
- (**Carpites**) **microcarpa**, Lesquereux, 1892, p. 93, pl. xvi, fig. 8.
Dakota Group; Kansas, U.S.A.
- **modesta**, Lesquereux, 1883, p. 53, pl. xvi, fig. 4.
Dakota Group; Colorado, U.S.A.
- **Nebrascensis**, Lesquereux, 1868, p. 98; & 1874, p. 74, pl. x, fig. 1;
pl. xxviii, fig. 14. (= *Persea Nebrascensis*, Lx., 1869.)
Dakota Group; Nebraska, U.S.A.
- **Newberryana**, Hollick, 1894 A, p. 52, pl. clxxix, fig. 5.
Cretaceous; Long Island, U.S.A.
- **notandia**, Saporta, 1894, p. 202, pl. xxxvii, fig. 7.
Upper Albian; Portugal.
- **Oдини**, Heer, 1882, p. 77, pl. xxxvii, fig. 9.
Atane Beds; Greenland.
- **Omalii**, Saporta & Marion, 1873, p. 49, pl. vi, fig. 1. Eocene.
(Recorded from American Cretaceous, Hollick, 1894 A.)
- **palæocretacea**, Saporta, 1894, p. 202, pl. xxxvii, fig. 12.
Albian; Portugal.
- **plutonia**, Heer, 1882, p. 75, pl. xix, figs. 1 d, 2-4; pl. xx, figs. 3 a,
4-6; pl. xxiv, fig. 6 b; pl. xxviii, figs. 10-11; pl. xlii, fig. 4 b.
Atane Beds; Greenland.
- **plutonina**, Ettingshausen, 1893, p. 149; & 1895, p. 26, pl. ii, fig. 15.
Cretaceous; Australia.
- **præstans**, Lesquereux, 1876 c; & 1878 B, p. 215, pl. lxiii, fig. 7.
Laramie Formation; Wyoming, U.S.A.
- **prætavia**, Saporta & Marion, 1885, p. 117, text-fig. 125 D.
Turonian; France.
- **primigenia**, Unger, 1850 A, p. 423.
Eocene; Sotzka.
(Recorded by Heer, in Meek & Hayden, 1859, p. 265, for
Dakota Cretaceous, U.S.A.)
- **proteæfolia**, Lesquereux, 1876 A, p. 393; & 1876 B, p. 342, pl. v,
figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- **sccialis**, Lesquereux, 1878 B, p. 213, pl. xxxvi, figs. 1, 2, 3, 4, 7.
Laramie Formation; Wyoming, U.S.A.
- **teliformis**, Lesquereux, 1892, p. 94, pl. 1, fig. 9.
Dakota Group; Kansas, U.S.A.
- **titahensis**, Lesquereux, 1878 B, p. 216, pl. xxvi, fig. 11.
Laramie Formation; Wyoming, U.S.A.
- ? **sp.**, Knowlton, 1900 A, p. 13, pl. i, fig. 4.
Montana Formation; Missouri R., U.S.A.

- Leguminosites albizzioides**, Menzel, 1909, p. 400, pl. 2, figs. 1, 2, 3.
Senonian; Cameroons.
- **amissus**, Heer, 1874 A, p. 119, pl. xxxii, fig. 19.
Atane Beds; Greenland.
- **atanensis**, Heer, 1874 A, p. 119, pl. xxxiv, fig. 6. Ibid.
- **cassiæformis**, Heer, 1874 A, p. 119, pl. xxxiv, fig. 12. (= *Cassia antiquorum*, Heer, 1882.) Ibid.
- **cassioides**, Lesquereux, 1878 B, p. 300, pl. lix, figs. 1-4 (name pre-occupied, re-named *Leguminosites Lesquereuxiana*, Knowlton, 1898).
Laramie Formation; Montana, U.S.A.
- **cassioides**, Engelhardt (*non* Lesquereux), in Vanhöffen, 1897, pp. 365 & 371, fig. 29. Atane Beds; Greenland.
- **constrictus**, Lesquereux, 1892, p. 151, pl. xlv, fig. 3.
Dakota Group; Kansas, U.S.A.
- **convolutus**, Lesquereux, 1892, p. 151, pl. xlv, fig. 4. Ibid.
- **coronilloides** (?), Heer, 1874 A, p. 119, pl. xxxiv, fig. 4. (= *Colutea coronilloides*, Heer, 1882.) Atane Beds; Greenland.
- **cretaceus**, Engelhardt, 1892 A, p. 104, pl. ii, figs. 11, 11 a.
Cenomanian; Niederschoena, Saxony.
- **cultriformis**, Lesquereux, 1883, p. 86, pl. x, fig. 4.
Dakota Group; U.S.A.
- **dakotensis**, Lesquereux, 1892, p. 150, pl. xxxviii, fig. 5.
Dakota Group; Kansas, U.S.A.
- **Dalageri**, Heer, 1882, p. 104, pl. xxxvii, fig. 10.
Atane Beds; Greenland.
- **dentatus**, Heer, 1883 A, p. 44, pl. lxx, figs. 10-11.
Patoot Beds; Greenland.
- **frigidus**, Heer, 1883 A, p. 44, pl. lxx, fig. 13; pl. lv, figs. 21 a, 22.
Patoot Beds; Greenland.
- **hymenophyllus**, Lesquereux, 1892, p. 152, pl. lv, figs. 7-9; pl. lvi, fig. 3. Dakota Group; Kansas, U.S.A.
- **infracretacicus**, Saporta, 1894, p. 208, pl. xxxvii, fig. 14.
Upper Albian; Portugal.
- **insularis**, Heer, 1882, p. 103, pl. xlv, fig. 6.
Atane Beds; Greenland.
- **lanceolatus**, Schenk, 1876, p. 170, pl. xxix, fig. 16.
Upper Cretaceous; Brandenburg, Tyrol.
- **Lesquereuxiana**, Knowlton, 1898, p. 131. (= *Leguminosites cassioides*, Lesquereux, 1878.)
Laramie Formation; Montana, U.S.A.
- **macilentus**, Heer, 1882, p. 104, pl. xxvii, fig. 15.
Atane Beds; Greenland.
- **Marcouanus**, Heer in Meek & Hayden, 1859, p. 265. (= *Liriodendron Meekii Marcouana*, Heer, 1882, & *Bumelia Marcouana*, Lesquereux, 1874.) Dakota Group; Nebraska, U.S.A.
- **omphalobioides**, Lesquereux, 1892, p. 149, pl. xxxviii, fig. 4.
Dakota Group; Kansas, U.S.A.

- Leguminosites orbiculatus*, Heer, 1882, p. 104, pl. xx, fig. 3 c.
Atane Beds; Greenland.
- *ovalifolius*, Heer, 1882, p. 103, pl. xxvii, fig. 14. Ibid.
- *ovatus*, Schenk, 1876, p. 170, pl. xxix, fig. 15.
Upper Cretaceous; Brandenburg, Tyrol.
- *pachyphyllus*, Ettingshausen, 1893, p. 151; & 1895, p. 52, pl. iv, fig. 18.
Cretaceous; Australia.
- *patootensis*, Heer, 1883 A, p. 44, pl. lxx, figs. 12, 12 b.
Patoot Beds; Greenland.
- *phaseolites*, Heer, 1874 A, p. 118, pl. xxxiv, figs. 7-11.
Atane Beds; Greenland.
- *podogonialis*, Lesquereux, 1892, p. 148, pl. xiii, fig. 11; pl. xxxviii, fig. 16.
Dakota Group; Kansas, U.S.A.
- *prodromus*, Heer, 1871, p. 1184; & 1874 A, p. 118, pl. xxxiv, fig. 13.
Atane Beds; Greenland.
- *raritensis*, Berry, 1909, p. 257, pl. xviii, fig. 4.
Raritan Formation; New Jersey, U.S.A.
- *robiniifolia*, Berry, 1910 A, p. 196.
Middle Cretaceous; North Carolina, U.S.A.
- *truncatus*, Lesquereux, 1892, p. 150, pl. xxi, fig. 7.
Dakota Group; Kansas, U.S.A.
- Lemna? bullata*, Lesquereux, 1875, p. 363.
Laramie Formation; Wyoming, U.S.A.
- *scutata*, Dawson, 1875.
(Recorded from Montana Group by Lesquereux, 1878.)
- Lepidocaryopsis Westphaleni*, Stur, 1873, p. 2.
Cenomanian; Bohemia.
- Lepidodendron punctatum*, Sternberg, 1820, p. 23, pl. iv, figs. 1, 2.
(= *Protopteris punctata*, Presl.)
- Leptospermum cretaceum*, Velenovsky, 1889, p. 21, pl. i, fig. 10.
(= *Leptospermum macrocarpum*, Vel.) Cenomanian, Bohemia.
- *macrocarpum*, Velenovsky, 1889, p. 27, pl. i, fig. 10. Ibid.
- Leptostrobus? alatus*, Ward, 1899 B, p. 673, pl. clxiii, figs. 16, 17.
Lower Cretaceous; Black Hills, U.S.A.
- *foliosus*, Fontaine, 1889, p. 230, pl. ci, fig. 4; pl. ciii, fig. 5; pl. civ, fig. 1.
Potomac Formation; Virginia, U.S.A.
- *longifolius*, Fontaine, 1889, p. 228, pl. ci, figs. 2, 3; pl. cii, figs. 1-4; pl. cii, figs. 6-12; pl. civ, fig. 6. Ibid.
- *? multifidus*, Fontaine, 1889, p. 230, pl. clxv, fig. 6. Ibid.
- *? ovalis*, Ward, 1905, p. 514, pl. cviii, figs. 9, 10 (naming *Leptostrobus? (b)* sp. Fontaine, 1889, p. 231, pl. cxxxvi, figs. 10, 10 a). Ibid.
- *? sp. (a)* Fontaine, 1889, p. 231, pl. cxxxv, fig. 6. Ibid.
- *? sp. (b)* Fontaine, 1889, p. 231, pl. cxxxvi, fig. 10. Ibid.
- *? sp. (c)* Fontaine, 1889, p. 232, pl. cxxxvi, fig. 11. Ibid.
- Libocedrus cretacea*, Heer, 1882, p. 49, pl. xxix, figs. 1-3; pl. xliii, fig. 1 d.
Atane Beds; Greenland.

- Libocedrus salicornioides cretacea**, Velenovsky, 1885, p. 28, pl. viii, fig. 2. Perucur Beds; Bohemia.
- **Veneris**, Velenovsky, 1885, p. 28, pl. x, figs. 13, 15, 17. Pläner; Bohemia.
- Limnophyllum lanceolatum**, Hosius & von der Marck, 1880, p. 184, pl. xxxviii, fig. 154. Lower Senonian; Westphalia.
- **primaevum**, Hosius & von der Marck, 1880, p. 183, pl. xxxviii, fig. 153. Ibid.
- Lindera Masconi**, Lesquereux, 1892, p. 96, pl. xviii, figs. 9, 10 (re-named *Benzoin Masoni*, Knowlton, 1898). Dakota Group; Kansas, U.S.A.
- **venusta**, Lesquereux, 1892, p. 95, pl. xvi, figs. 1, 2 (re-named *Benzoin venustum*, Knowlton, 1898). Ibid.
- Liquidambar integrifolium**, Lesquereux, 1868, p. 93; & 1874, p. 56, pl. ii, figs. 1-3; pl. xxiv, fig. 2; pl. xxix, fig. 8. Dakota Group; Nebraska, U.S.A.
- **obtusilobatus** (Heer), Hollick in Newberry, 1898, p. 101, pl. v, fig. 4; pl. xii, fig. 4. (= *Phyllites obtusilobatus*, Heer, 1859.) Dakota Group; Kansas, U.S.A.
- Liriodendron acuminatum**, Lesquereux, 1884, p. 227. Dakota Group; Kansas, U.S.A.
- **acuminatum**, var. **bilobatum**, Lesquereux, 1892, p. 207, pl. xxviii, fig. 4. Ibid.
- **alatum**, Newberry MS., Hollick, 1894 E, p. 467, pl. ccxx. Laramie Formation; Colorado, U.S.A.
- **attenuatum**, Hollick, 1906 A, p. 68, pl. xxi, figs. 9-11. Middle Cretaceous; Long Island, U.S.A.
- **Celakovskii**, Velenovsky, 1882 A, p. 213; & 1883, p. 43, pl. xiv, fig. 2. Cenomanian; Bohemia.
- **cruciforme**, Lesquereux, 1884, p. 227. (= *L. giganteum cruciforme*, Lx., 1892.) Dakota Group; Kansas, U.S.A.
- **dubium**, Berry, 1907, p. 196, pl. xiv, fig. 3. Middle Cretaceous; North Carolina, U.S.A.
- **giganteum**, Lesquereux, 1868, p. 99; & 1874, p. 93, pl. xxii, fig. 2. Dakota Group; Nebraska, U.S.A.
- **giganteum cruciforme**, Lesquereux, 1892, p. 206, pl. xxviii, figs. 1, 2. (= *L. cruciforme*, Lx., 1884.) Dakota Group; Kansas, U.S.A.
- **intermedium**, Lesquereux, 1868, p. 99; & 1874, p. 93, pl. xx, fig. 5. Dakota Group; Nebraska, U.S.A.
- **laramiensis**, Ward, 1885, p. 556, pl. lx, fig. 1. Montana Formation; Wyoming, U.S.A.
- **Meekii**, Heer in Meek & Hayden, 1859, p. 265; & Heer, 1867, p. 21, pl. iv, fig. 3. Dakota Group; Nebraska, U.S.A.
- **Meekii genuina**, Heer, 1882, p. 98, pl. xvii, figs. 12-13; pl. xxiii, fig. 6. Atane Beds; Greenland.

- Liriodendron Meekii Marcouana**, Heer, 1882, p. 82, pl. xxii
figs. 4-7, 11; pl. xxiii, fig. 3; pl. xlv, fig. 13 *a, b*. (= *Leguminosites*
Marcouanus, Heer, 1859.) Atane Beds; Greenland.
- **Meekii mucronulata**, Heer, 1882, p. 88, pl. xxii, figs. 3-10.
Ibid.
- **Meekii obcordata**, Heer, 1882, p. 88, pl. xxii, figs. 1 *b*, 2; pl. xxiii,
fig. 4. (= *Phyllites obcordatus*, Heer, 1859.) Ibid.
- **Meekii primæva**, Heer, 1882, p. 88, pl. xviii, fig. 4 *c*; pl. xxii,
fig. 9; pl. xxiii, fig. 5. Ibid.
- **morganensis**, Berry, 1906 *v*, p. 176; and 1906 *z*, p. 150, pl. xxi,
figs. 2-4; pl. xxiii, fig. 1; pl. xxv.
Magothy Formation; New Jersey, U.S.A.
- **oblongifolium**, Newberry, 1887, p. 5, pl. lxi, fig. 1.
Amboy Clay; New Jersey, U.S.A.
- **pinnatifidum**, Lesquereux, 1884, p. 227.
Dakota Group; Kansas, U.S.A.
- **prætulipiferum**, Dawson, 1894, p. 63, pl. viii, fig. 27.
Upper Cretaceous; Vancouver Island, Canada.
- **primævum**, Newberry, 1870, p. 12; & 1878, pl. vi, fig. 7 (not
named); & Lesquereux, 1892, p. 203, pl. xxiv, fig. 4; pl. xxvi, figs. 1-4.
Dakota Group; Nebraska, U.S.A.
- **quercifolium**, Newberry, 1887, p. 6, pl. lxii, fig. 1.
Amboy Clay; New Jersey, U.S.A.
- **Schwarzii**, Richter, 1904, p. 16, pl. ii, fig. 10; 1905, p. 6, pl. i,
fig. 10. Senonian; Quedlinburg.
- **semi-alatum**, Lesquereux, 1884, p. 227; & 1892, p. 204, pl. xxv,
figs. 2-4; pl. xxix, fig. 3. Dakota Group; Kansas, U.S.A.
- **simplex**, Newberry, 1887, p. 6, pl. lxii, figs. 2-4. (= *Liriodendropsis*
simplex & *L. angustifolia*, Newberry, 1896.)
Amboy Clay; New Jersey, U.S.A.
- **Snowii**, Lesquereux, 1892, p. 209, pl. xxix, figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- **succedens**, Dawson, 1894, p. 62, pl. viii, fig. 26.
Upper Cretaceous; Vancouver Island, Canada.
- **Wellingtonii**, Lesquereux, 1892, p. 208, pl. xxviii, fig. 3.
Dakota Group; Kansas, U.S.A.
- Liriodendropsis angustifolia**, Newberry, 1895, p. 84, pl. liii, fig. 8.
(= *Liriodendron simplex*, Newberry, 1887, in part.)
Amboy Clay; New Jersey, U.S.A.
- **constricta** (Ward), Hollick, 1906 *A*, p. 71, pl. xxii, fig. 7; pl. xxvi,
figs. 6-15; pl. xl, fig. 15.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- **lacerata** (Sap.), Ward, 1896, p. 540. (= *Chondrophyton laceratum*,
Saporta, 1894.) Cenomanian; Portugal.
- **retusa** (Heer), Hollick, 1906 *A*, p. 72, pl. xxv, figs. 8, 9. (= *Sapo-*
tacites retusus, Heer, 1883 *A*.)
Middle Cretaceous; New Jersey, U.S.A.

- Liriodendropsis simplex**, Newberry, 1895, p. 83, pl. xix, figs. 2, 3; pl. liii, figs. 1-4, 7. (= *Liriodendron simplex*, Newberry, 1887, in part.)
Amboy Clay; New Jersey, U.S.A.
- **simplex constricta**, Ward, 1896, p. 540.
Middle Cretaceous; New Jersey, U.S.A.
- **spectabilis**, Hollick, 1906 A, p. 73, pl. xxii, figs. 1-6.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Liriophyllum Beckwithii**, Lesquereux, 1878 A, p. 482; & 1883, p. 76, pl. x, fig. 1.
Dakota Group; Colorado, U.S.A.
- **obcordatum**, Lesquereux, 1883, p. 77; & 1892, p. 210, pl. xxviii, fig. 7.
Ibid.
- **populoides**, Lesquereux, 1878 A, p. 483; & 1883, p. 76, pl. xi, figs. 1, 2.
Ibid.
- Lithothamnium amphiroaeformis**, Rothpletz, 1891 B, p. 314, pl. xvi, figs. 10, 14.
Turonian; France.
- **cenomanicum**, Rothpletz, 1891 B, p. 313, pl. xvi, figs. 1, 2, 16.
Cenomanian; France.
- **Goldfussi**, Gümbel, 1871, p. 285, pl. D iii, fig. L 3 a-b.
Upper Cretaceous; Maestricht.
- **gosaviense**, Rothpletz, 1891 B, p. 314, pl. xvii, figs. 1, 3.
Senonian; France.
- **mamillosum**, Gümbel, 1871, p. 41, pl. ii, figs. 7 a-b.
Danian; Maestricht.
- **palmatum**, Goldfuss in Gümbel, 1871, p. 284, pl. Div, fig. L 1 a, 1 b.
Upper Cretaceous; France.
- **parisiense**, Gümbel, 1871, p. 42, pl. ii, figs. 8 a-b.
Danian; Maestricht.
- **perulatum**, Gümbel, 1871, p. 44, pl. ii, figs. 11 a-c.
Ibid.
- **procaenum**, Gümbel, 1871, p. 43, pl. ii, figs. 10 a-b.
Ibid.
- **racemosum**, Goldfuss in Gümbel, 1871, p. 284, pl. Div, figs. L 2 a, 2 b.
Upper Cretaceous; France.
- **turonicum**, Rothpletz, 1891 B, p. 313, pl. xvi, figs. 9, 13.
Turonian; France.
- Litsaea bohemica**, Engelhardt, 1892, p. 101, pl. i, fig. 8.
Cretaceous; Kaunitz, Bohemia.
- **carbonensis**, Ward, 1885, p. 553, pl. xvi, fig. 11. (= *Malapoenna carbonensis* (Lx.), Knowlton, 1898, renamed.)
Laramie Formation; Wyoming, U.S.A.
- **expansa**, Saprota & Marion, 1878, p. 63, pl. ii, figs. 1, 2.
(Recorded from Cenomanian of Argentina.)
- **laurinoides**, Hosius & von der Marek, 1880, p. 189, pl. xl, fig. 157. (= *Phyllites laurinoides*, Hosius, 1870.)
Lower Senonian; Westphalia.
- Litsea cretacea**, Lesquereux, 1892, p. 96, pl. xv, fig. 2. (= *Malapoenna cretacea* (Lx.), Knowlton, 1898, renamed.)
Dakota Group; Kansas, U.S.A.

- Litsea falcifolia*, Lesquereux, 1892, p. 97, pl. xi, fig. 5. (= *Malapoenna falcifolia* (Lx.), Knowlton, 1893, renamed.)
Dakota Group; Kansas, U.S.A.
- *Weediana*, Knowlton in Weed & Knowlton, 1893, p. 55. (= *Malapoenna Weediana*, Knowlton, 1898.)
Laramie Formation; Montana, U.S.A.
- Lochmophycus caulerpoides*, Debey & Ettingshausen, 1859 A, p. 198, pl. ii, figs. 1-5.
Senonian; Aix, Rhenish Prussia.
- Lomatia Saportanea*, Lesquereux, 1876 B, p. 346.
Dakota Group; Kansas, U.S.A.
- ? *Saportanea*, var. *longifolia*, Lesquereux, 1883, p. 52.
Dakota Group; Colorado, U.S.A.
- Lomatites palæo-Ilex*, Ettingshausen, 1867 A, p. 255, pl. iii, fig. 16.
Cenomanian; Niederschoena, Saxony.
- Lomapteris superstes*, Saporta, 1879, p. 198, text-fig. 27, i.
Turonian; Toulon, France.
- Lonchopteris Huttoni*, Presl in Sternberg, 1838, p. 166. (= *Lonchopteris Mantelli*, Lindley & Hutton, p. 59, pl. clxxi.)
Lower Greensand; England.
- *Iusitanica*, Saporta, 1890 A, p. 814; & 1894, p. 80, pl. xvi, fig. 13.
Valanginian; Portugal.
- *Mantelli*, Brongniart, 1823, p. 369, pl. cxxxi, figs. 4 & 5. (= *Poly-podites Mantelli*, Goeppert, 1836.)
Lower Cretaceous; Beauvais.
- Lycopodites Francheti*, Saporta, 1894, p. 131, pl. xxiii, fig. 13; pl. xxv, fig. 21.
Urgonian; Portugal.
- *gracilis*, Brongniart (*non* Oldham & Morris), 1828 A, p. 84 [nomen nudum].
Lower Cretaceous; Bavaria.
- *gracillimus*, Saporta, 1894, p. 131, pl. xxvi, fig. 5.
Urgonian; Portugal.
- *insignis*, Reich in Geinitz, 1842, p. 98.
Cenomanian; Niederschoena, Saxony.
- *Limai*, Saporta, 1894, p. 132, pl. xxiii, figs. 16-17; pl. xxiv, fig. 11 a; pl. xxvi, fig. 1.
Urgonian; Cercal, Portugal.
- ? *montanensis*, Fontaine in Ward, 1905, p. 302, pl. lxxii, figs. 15, 16.
Kootanie Formation; Montana, U.S.A.
- Lycopodium cretaceum*, Berry, 1910 G, p. 276, text-fig. 1.
Upper Cretaceous; S. Carolina, U.S.A.
- *Lesquereuxii*, Knowlton, 1897, p. 154; & 1900 A, p. 24.
Montana Formation; Wyoming, U.S.A.
- *redivivum*, Heer, 1874 A, p. 60, pl. xiii, fig. 1.
Kome Beds; Greenland.
- Lygodites patulatus*, Schulze, 1888, p. 26 [nomen nudum].
Upper Quader; Germany.
- Lygodium compactum*, Lesquereux, 1878 B, p. 64, pl. v, fig. 9.
Laramie Formation; Colorado, U.S.A.
- *cretaceum*, Debey & Ettingshausen, 1859 B, p. 198, pl. ii, figs. 18-21; pl. iii, fig. 28.
Senonian; Aix, Rhensish Prussia.

- Lygodium trichomanoides**, Lesquereux, 1874, p. 45, pl. i, fig. 2.
Dakota Group; Kansas, U.S.A.
- Macclintockia appendiculata**, Heer, 1882, p. 71, pl. xxxvii, fig. 1.
Atane Beds; Greenland.
- **cretacea**, Heer, 1882, p. 70, pl. xxxvi, figs. 1, 2a; pl. xxxvii, figs. 2-4. Ibid.
- **trinervis**, Heer, 1868, p. 115, pl. xv, figs. 7-9. Tertiary.
(Recorded Dawson, 1894, Cretaceous.)
- Macrotaeniopteris vancouverensis**, Dawson, 1894, p. 55, pl. v, figs. 1-3. Upper Cretaceous; Vancouver Island, Canada.
- Magnolia alternans**, Heer in Capellini & Heer, 1867, p. 20, pl. iii, figs. 2-4; pl. iv, figs. 1-2. Dakota Group; Nebraska, U.S.A.
- **amplifolia**, Heer, 1869 A, p. 21, pl. viii, figs. 1, 2; pl. ix, fig. 1. Cenomanian; Moravia.
- **angustifolia**, Newberry, 1883, p. 513.
Laramie Formation; New Mexico, U.S.A.
- **auriculata**, Newberry MS. in Hollick, 1894 A, p. 61, pl. clxxix, figs. 6, 7; & Newberry, 1895, p. 75, pl. xli, fig. 13; pl. lviii, figs. 1-11. (Re-named *Magnolia Hollicki*, Berry, 1909.)
Cretaceous; Long Island, U.S.A.
- **Boulayana**, Lesquereux, 1892, p. 202, pl. lx, fig. 2. Dakota Group; Kansas, U.S.A.
- **Capellinii**, Heer in Capellini & Heer, 1867, p. 21, pl. iii, figs. 5, 6. Dakota Group; Nebraska, U.S.A.
- **cenomanensis**, Saporta & Marion, 1878, p. 13. Cenomanian; Bohemia.
- **cuneata**, Newberry in Reynold's Expl., 1869, p. 163 [nomen nudum]. Cretaceous; Orcas Island, U.S.A.
- **Delgadoi**, Saporta, 1894, p. 194, pl. xxxv, fig. 5. Albian (?); Portugal.
- **ensifolia**, Lesquereux, 1872, p. 302. (= *Celastrophyllum ensifolium*, Lx., 1874.) Dakota Group; Kansas, U.S.A.
- **glaucoides**, Newberry MS. in Hollick, 1894 A, p. 60, pl. clxxv, figs. 1, 7; & Newberry, 1895, p. 74, pl. lvii, figs. 1-4. Cretaceous; Long Island, U.S.A.
- **Hollicki**, Berry, 1909, p. 253 (re-naming *Magnolia auriculata*, Newberry in Hollick, 1894 A). Raritan Formation; New Jersey, U.S.A.
- **Isbergiana**, Heer, 1882, p. 91, pl. xxxvi, fig. 3. Atane Beds; Greenland.
- **Lacoeana**, Lesquereux, 1892, p. 201, pl. lx, fig. 1. Dakota Group; Kansas, U.S.A.
- **longifolia**, Newberry MS. in Hollick, 1893, p. 36, pl. iii, fig. 9; & Newberry, 1895, p. 76, pl. lv, figs. 3, 5; pl. lvi, figs. 1-4. (= *Magnolia Newberryi*, Berry, 1907.) Raritan (?) Formation; Woodbridge, U.S.A.

- Magnolia longipes**, Newberry MS. in Hollick, 1894 A, p. 60, pl. clxxviii, figs. 1, 3; & Newberry, 1895, p. 76, pl. liv, figs. 1-3.
Cretaceous; Long Island, U.S.A.
- **magnifica**, Dawson, 1883, p. 22, pl. iii, fig. 11.
Upper Cretaceous; North-West Territory, Canada.
- **Marbodi**, Krasser & Kubart, 1906 B, p. 47 [nomen nudum].
Cenomanian; Moletain, Moravia.
- **Newberryi**, Berry, 1907, p. 195, pl. xiii, fig. 6. (Re-naming *M. longifolia*, Newberry.)
Middle Cretaceous; North Carolina, U.S.A.
- **obovata**, Newberry, 1870, p. 15. See *Nyssa vetusta*, Newberry, 1868.
- **obtusata**, Heer, 1882, p. 90, pl. xv, fig. 12; pl. xxi, fig. 3.
Atane Beds; Greenland.
- **occidentalis**, Dawson, 1894, p. 63, pl. x, fig. 36.
Upper Cretaceous; Vancouver Island, Canada.
- **palæocretacica**, Saporta, 1894, p. 222, pl. xxxix, fig. 26.
Cenomanian (?); Portugal.
- **palæopetala**, Hollick, 1903, p. 102, text-fig. A on p. 103.
Dakota Group; Kansas, U.S.A.
- **pseudoacuminata**, Lesquereux, 1892, p. 199, pl. xxiv, fig. 2.
Ibid.
- **pulchra**, Ward, 1885, p. 556, pl. lx, figs. 2, 3.
Montana Formation; Wyoming, U.S.A.
- **rotundifolia**, Newberry, 1883, p. 513.
Laramie Formation; New Mexico, U.S.A.
- **? sarthacensis**, Crié, 1879 (?), p. 24 [nomen nudum].
Cretaceous; Sainte-Croix, France.
- **speciosa**, Heer, 1869 A, p. 20, pl. vi, fig. 1; pl. ix, fig. 2; pl. x; pl. xi, fig. 1.
Cenomanian; Moletain, Moravia.
- **telonensis**, Saporta, 1879, p. 198, text-fig. 27, 3.
Turonian; Toulon, France.
- **tenuifolia**, Lesquereux, 1868, p. 100; & 1874, p. 92, pl. xxi, fig. 1.
Dakota Group; Nebraska, U.S.A.
- **Van Ingeni**, Hollick, 1894 A, p. 61, pl. clxxv, fig. 6.
Cretaceous; Long Island, U.S.A.
- **woodbridgensis**, Hollick in Newberry, 1895, p. 74, pl. xxxvi, fig. 11; pl. lvii, figs. 5-7.
Amboy Clay; New Jersey, U.S.A.
- Magnoliæphyllum ? sp.**, Krasser, 1896, p. 131, pl. xvii, fig. 12.
Cretaceous; Kunstadt, Moravia.
- Majanthemophyllum cretaceum**, Heer, 1882, p. 57, pl. xxx, fig. 22.
Atane Beds; Greenland.
- **lanceolatum**, Heer, 1882, p. 58, pl. xxx, fig. 21.
Patoot Beds; Greenland.
- **pusillum**, Heer, 1883 A, p. 18, pl. lv, figs. 17, 17 b. Ibid.
- Malapoenna carbonensis** (Ward), Knowlton, 1898, p. 142 (re-naming *Litsæa carbonensis*, Ward, 1885).
Laramie Formation; Wyoming, U.S.A.

- Malapoenna cretacea** (Lesquereux), Knowlton, 1898, p. 142 (re-naming *Litsea cretacea*, Lx., 1892). Dakota Group; Kansas, U.S.A.
- **falcifolia** (Lesquereux), Knowlton, 1898, p. 142 (re-naming *Litsea falcifolia*, Lx., 1892). Ibid.
- **horrellensis**, Berry, 1910 A, p. 198, pl. xxiv, figs. 1-9.
Middle Cretaceous; North Carolina, U.S.A.
- **macrophylloides**, Knowlton, 1900 A, p. 57, pl. xiv, figs. 4, 5.
Montana Formation; Wyoming, U.S.A.
- **Weediana**, Knowlton, 1898, p. 142 (re-naming *Litsea Weediana*, Knowlton, 1893).
- **sp.**, Hollick, 1906 A, p. 78, pl. xxxi, fig. 4.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Malpighiastrum cretaceum**, Ettingshausen, 1893, pp. 138, 150; & 1895, p. 47, pl. iv, fig. 1. Cretaceous; Australia.
- Mammacites Francheti**, Fliche, 1896, p. 283, pl. xiii, figs. 7 and 71.
Cenomanian; Chaudefontaine, France.
- Manihotites georgiana**, Berry, 1910 B, p. 507, text-figs. 1, 2.
Upper Cretaceous; Georgia, U.S.A.
- Mantellia? Babbagensis**, Woodward, 1885, p. 290, pl. vii, figs. 1-3.
Cretaceous (?); Mt. Babbage, Australia.
- **inclusa**, Carruthers, 1870, p. 703, pl. lxiii, figs. 2, 3. (= *Cyca-deoidea inclusa* (Carr.), Schimper, 1874.)
Lower Greensand; Potton.
- Marattia cretacea**, Velenovsky, 1888 B, p. 9, pl. i, fig. 13.
Perucer Beds; Bohemia.
- **minor**, Saporta, 1894, p. 83, pl. xvi, fig. 14.
Valanginian; Portugal.
- Marattites desideratus**, Marion & Laurent, 1898, p. 189, pl. i, fig. 1.
Cretaceous; Roumania.
- Marsilea Andersoni**, Hollick, 1905 C, p. 409, pl. lxxi, figs. 1-3.
Cretaceous; Long Island, U.S.A.
- **attenuata**, Hollick, 1894 D, p. 256, pl. ccv, fig. 10. (= *Salvinia attenuata*, Lesquereux, 1876 C.)
Laramie Formation; Wyoming, U.S.A.
- Marsilia cretacea**, Heer, 1882, p. 39, pl. xvi, figs. 11, 12.
Atane Beds; Greenland.
- **? (Baiera) grandis**, Heer, 1874, p. 37, pl. iii, fig. 4. Ibid.
- **perucensis** (Velenovsky), Bayer in Fritsch & Bayer, 1901, p. 86, text-fig. 34. (= *Marsilia sp.*, Velenovsky.)
Perucer Beds; Vyserovic, Bohemia.
- **? sp.**, Velenovsky, 1888, p. 597, figs. 11-13. (= *M. perucensis* (Velenovsky), Bayer.) Ibid.
- Matonia Wiesneri**, Krasser, 1896, p. 119, pl. xi, fig. a 1; pl. xii, figs. 1 & 2; pl. xvi, fig. 10. (= *Matonidium Wiesneri*, Krasser, 1889.)
Cenomanian; Kunstadt, Moravia.
- Matonidium Althausii** (Dunker), Ward, 1899 B, p. 653, pl. 160, figs. 5-8.
Lower Cretaceous; Black Hills, U.S.A.

- Matonidium Wiesneri**, Krasser, 1889, p. 34 [nomen nudum]. (= *Matonia Wiesneri*, Krasser, 1896.) Cenomanian; Kunstadt, Moravia.
- Megalozamia falciformis**, Hosius & von der Marck, 1880, p. 203, pl. xliii, figs. 181-183 a & b. Lower Gault; Westphalia.
- Melastomites cuneiformis**, Hosius & von der Marck, 1880, p. 190, pl. xl, fig. 159. Lower Senonian; Westphalia.
- **parvula**, Unger, 1865, p. 377, pl. i, fig. 8. Cretaceous (?); Deva, Transylvania.
- Melophytum cyclostigma**, Debey & Eitingshausen, 1859 n, p. 241, pl. vii, figs. 28-30. Senonian; Aix, Rhenish Prussia.
- Menispermities acerifolia**, Lesquereux, 1874, p. 96, pl. xx, figs. 2, 3. (= *Acerites menispermifolia*, Lesquereux, 1868 = *Menispermities menispermifolia*, Knowlton, 1898.) Dakota Group; Nebraska, U.S.A.
- **acutibolus**, Lesquereux, 1883, p. 78, pl. xiv, fig. 2. Dakota Group; Kansas, U.S.A.
- **borealis**, Heer, 1882, p. 91, pl. xxxix, fig. 2. Atane Beds; Greenland.
- **Brysoniana**, Hollick, 1894 A, p. 59, pl. clxxx, fig. 10. Cretaceous; Long Island, U.S.A.
- **? bunzlaviensis**, Roemer, 1889, p. 144, pl. xii, fig. 9. Senonian; Bunzlau, Silesia.
- **californicus**, Fontaine in Ward, 1905, p. 268, pl. lxix, figs. 12-14. Shasta Formation; California, U.S.A.
- **cercidifolius**, Saporta, 1894, p. 191, pl. xxxv, fig. 7. Albion; Portugal.
- **cyclophyllus**, Lesquereux, 1876 n, p. 358, pl. vi, fig. 4. Dakota Group; Kansas, U.S.A.
- **dentatus**, Heer, 1882, p. 92, pl. xxxviii, fig. 4. Atane Beds; Greenland.
- **grandis**, Lesquereux, 1883, p. 80, pl. xv, figs. 1-2. Dakota Group; Kansas, U.S.A.
- **Knightii**, Knowlton, 1900 A, p. 61, pl. xv, fig. 2. Montana Formation; Wyoming, U.S.A.
- **menispermifolius** (Lesquereux), Knowlton, 1898, p. 143 (renaming *Acerites menispermifolius*, Lx., 1868 & *Menispermities acerifolia*, Lesquereux, 1874). Dakota Group; Nebraska, U.S.A.
- **obtusiloba**, Lesquereux, 1874, p. 94, pl. xxv, figs. 1-2; pl. xxvi, fig. 3. Dakota Group; Kansas, U.S.A.
- **obtusiloba**, var. (?), Lesquereux, 1874, p. 95, pl. xxii, fig. 1. (= *Dombeyopsis obtusiloba*, Lesquereux, 1868.) Ibid.
- **ovalis**, Lesquereux, 1876 A, p. 398; & 1876 B, p. 357, pl. v, fig. 4. Ibid.
- **populifolius**, Lesquereux, 1876 n, p. 357, pl. v, fig. 3. Ibid.
- **reniformis**, Dawson, 1883, p. 23, pl. iv, fig. 12. Upper Cretaceous; North-West Territory, Canada.
- **rugosus**, Lesquereux, 1892, p. 196, pl. xxix, fig. 7. Dakota Group; Kansas, U.S.A.

- Menispermities salinæ** (Lesquereux), Knowlton, 1898, p. 143 (re-naming *Populites Salinæ*, Lx., 1873). Dakota Group; Kansas, U.S.A.
 — **salinensis**, Lesquereux, 1874, p. 95, pl. xx, figs. 1, 4. Ibid.
 — **tenuinervis**, Fontaine, 1889, p. 322, pl. clxxii, fig. 8.
 Potomac Formation; Maryland, U.S.A.
 — **virginiensis**, Fontaine, 1889, p. 321, pl. clxi, figs. 1, 2.
 Potomac Formation; Virginia, U.S.A.
 — **Wardianus**, Hollick in Newberry, 1895, p. 85, pl. xxix, figs. 9, 11.
 Amboy Clay; New Jersey, U.S.A.
 — **sp.**, Dawson, 1894, p. 62, pl. xi, fig. 50.
 Upper Cretaceous; Vancouver Island, Canada.
 — **sp.**, Hollick, 1906 A, p. 62, pl. xii, fig. 7 (for *Hedera* sp.?, Hollick).
 Cretaceous; Staten Island, U.S.A.
Menispermophyllum Celakovskianum, Velenovsky, 1889, p. 54
 [nomen nudum]. (= *Menispermophyllum Celakovskii*, Velenovsky.)
 Perner Beds; Bohemia.
 — **Celakovskii**, Velenovsky, in Fritsch & Bayer, 1901, p. 128. Ibid.
Menispermum (Cocculus) assimile, Marion, 1890, p. 1054.
 Turonian; Martigues, France.
Mertensia Kurriana (Heer), Engelhardt, 1892 A, p. 84. (= *Gleichenia Kurriana*, Heer.)
 Cretaceous; Bohemia.
 — **Zippei**, Engelhardt, 1892 A, p. 83. (= *Pecopteris Zippei*, Corda = *Gleichenia Zippei*, Heer.) Ibid.
Metrosideros perigrinus, Heer, 1874 A, p. 116, pl. xxvii, fig. 22.
 Atane Beds; Greenland.
Microdictyon regale, Richter, 1899 A, p. 40.
 Senonian; Quedlinburg, Saxony.
Microlepiea pluripartita, Saporta, 1894, p. 83, pl. xv, fig. 17; pl. xvi, fig. 24.
 Valanginian; Portugal.
Microlepidium striatulum, Velenovsky, 1889, p. 11, pl. i, figs. 25-27.
 Cenomanian; Bohemia.
Microzamia? dubia, Berry, 1905 A, p. 43, pl. i, fig. 2.
 Cretaceous; New Jersey, U.S.A.
 — **gibba**, Corda in Reuss, 1846, p. 85, pl. xvi, figs. 1-10. (= *Conites gibbus*, Reuss, 1844.)
 Quader & Pläner; Bohemia.
Mimusops ballotæoides, Engelhardt, 1892 A, p. 98, pl. ii, fig. 13.
 Quader; Saxony.
Mitropicea Decheni, Debey, 1848 A, p. 120 [nomen nudum].
 Senonian; Aix, Rhenish Prussia.
 — **Noeggerathii**, Debey, 1848 A, p. 120 [nomen nudum]. Ibid.
Monheimia aquisgranensis, Debey & Ettingshausen, 1859 B, p. 211, pl. iv, figs. 3-10. Ibid.
 — **polypodioides**, Debey & Ettingshausen, 1859 B, p. 211, pl. iii, figs. 34-36; pl. iv, figs. 1-2, 21. Ibid.
Monimia præ-vestita, Ettingshausen, 1893, pp. 135, 149; & 1895, p. 23, pl. ii, figs. 13, 14.
 Cretaceous; Australia.
Moriconia americana, Berry, 1910 B, p. 20.
 Magothy Formation; Maryland, U.S.A.

- Moricona cyclotoxon**, Debey & Ettingshausen, 1859 b, p. 239, pl. vii, figs. 23-27. Senonian; Aix, Rhenish Prussia.
- Morinium populifolium**, Ettingshausen in Reuss, 1854, p. 740 [nomen nudum]. Cenomanian; Moletain, Moravia.
- Munieria baconica**, von Hantken in Deecke, 1883, p. 9, pl. i, figs. 4-10. Cretaceous; Austria.
- Münsteria annulata**, Unger. (= *Keckia annulata*, Glocker, 1841.)
- **cretacea**, W. A. Ooster, 1871, p. 69, pl. xi, fig. 24. Upper Cretaceous; Switzerland.
- ? **Goepperti**, Geinitz, 1850 A, p. 266. Quadersandstein; Silesia.
- **Keckii**, Unger, 1850 A, p. 14. (= *Muensteria annulata*, Unger = *Keckia annulata*, Glocker, 1841.)
- **Massalongiana**, Zigno, 1864, p. 525 [nomen nudum]. Neocomian; Italy.
- **rugosa**, Zigno, 1864, p. 525 [nomen nudum]. Ibid.
- **Schneideriana**, Goeppert, 1842 A, p. 115, pl. li, fig. 3. Quadersandstein; Silesia.
- **spiralis**, Massalongi, 1857, p. 778 [nomen nudum]. Neocomian; Italy.
- Muscites cretaceus**, Debey & Ettingshausen, 1859 b, p. 185, pl. i, fig. 6. Senonian; Aix, Rhenish Prussia.
- Myrica acuta**, Hollick in Newberry, 1895, p. 65, pl. xlii, fig. 35. Amboy Clay; U.S.A.
- **aspera**, Lesquereux, 1892, p. 66, pl. ii, fig. 11. Dakota Group; Kansas, U.S.A.
- **Brittoniana**, Berry, 1905 A, p. 46. (= *Myrica Heerii*, Berry, non Boulay, 1903.) Cretaceous; New Jersey, U.S.A.
- **Brookensis**, Fontaine, 1889, p. 310, pl. cl, fig. 11; pl. clvi, fig. 10. Potomac Formation; Virginia, U.S.A.
- **Campei**, Marion, 1890, p. 1054 [nomen nudum]. Turonian; Martigues, France.
- **cinnamomifolia**, Newberry, 1895, p. 64, pl. xxii, figs. 9-14. Amboy Clay; New Jersey, U.S.A.
- **cliffwoodensis**, Berry, 1904 A, p. 73, pl. iv, fig. 1. Matawan Formation; Cliffwood, New Jersey, U.S.A.
- **cretacea**, Heer (non Lesquereux), 1871 A, p. 10, pl. iii, fig. 2 a-c. Senonian; Quedlinburg, Saxony.
- **cretacea**, Lesquereux, 1876 A, p. 392; & 1876 B, p. 339, pl. iii, fig. 4. (Name pre-occupied, = *M. dakotensis*, Lx., 1883.) Dakota Group; Kansas, U.S.A.
- **dakotensis**, Lesquereux, 1883, p. 35, pl. iv, fig. 9. (= *M. cretacea*, which name was preoccupied by Heer.) Ibid.
- **Davisii**, Hollick, 1893, p. 32, pl. ii, fig. 3. Cretaceous; Kreischerville, New York, U.S.A.
- **elegans**, Berry, 1907, p. 191, pl. xi, figs. 1-4, 6. Middle Cretaceous; North Carolina, U.S.A.
- **emarginata**, Heer, 1882, p. 66, pl. xli, fig. 2; pl. xlii, fig. 12 e. Atane Beds; Greenland.

- Myrica fenestrata**, Newberry, 1895, p. 63, pl. xlii, fig. 32.
Amboy Clay; New Jersey, U.S.A.
- **fragiliformis** (Zenker), Engelhardt, 1892 A, p. 98. (= *Salix fragiliformis*, Zenker, 1833.) Cenomanian; Niederschöna; Saxony.
- **Gaudryi**, Marion, 1890, p. 1054 [nomen nudum].
Turonian; Martigues, France.
- **gracilior**, Saporta, 1894, p. 218, pl. xxxviii, figs. 10-11.
Cenomanian; Portugal.
- **grandifolia**, Hollick, 1893, p. 32, pl. iii, fig. 1. (Name pre-occupied
= *M. Hollicki*, Ward, 1893.)
Raritan (?) Formation; Staten Island, U.S.A.
- **Heerii**, Berry, 1903 c, p. 682, text-figs. 7, 8, p. 678. (= *Myrica Brittoniana*, Berry, 1905 A.)
Matawan Formation; New Jersey, U.S.A.
- **Hollicki**, Ward, 1893, p. 437. (= *M. grandifolia*, Hollick, 1893.)
- **indigena**, Krasser, 1889, p. 34; & 1896, p. 129, pl. xv, fig. 1.
Cenomanian; Kunstadt, Moravia.
- **lacera**, Saporta, 1894, p. 200, pl. xxxvi, figs. 6, 7.
Upper Albian; Portugal.
- **leiophylla**, Hosius & von der Marck, 1880, p. 155, pl. xxviii, fig. 47.
Upper Senonian; Haldem, Westphalia.
- **Lessigiana**, Lesquereux. (= *Artocarpus Lessigiana*, Knowlton.)
- **longa**, Heer, 1882, p. 65, pl. xviii, fig. 9 b; pl. xxix, figs. 15-17;
pl. xxxiii, fig. 10; pl. xli, fig. 4 b, d. (= *Proteooides longus*, Heer,
1874 A.)
Atane Beds; Greenland.
- **Newberryana**, Hollick in Newberry, 1895, p. 63, pl. xlii, fig. 5.
Amboy Clay; New Jersey, U.S.A.
- **obliqua**, Knowlton in Lesquereux, 1892, p. 68, pl. lxiv, fig. 16.
Dakota Group; Kansas, U.S.A.
- **obtusa**, Lesquereux, 1874, p. 63, pl. xxiv, fig. 10. Ibid.
- (*Comptonia*) **parvula**, Heer, 1883 A, p. 20, pl. lv, figs. 1-3.
Patoot Beds; Greenland.
- **præcox**, Heer, 1883 A, p. 21, pl. lv, fig. 4. Ibid.
- **primæva**, Hosius & von der Marck, 1880, p. 155, pl. xxviii, fig. 46.
Upper Senonian; Haldem, Westphalia.
- **pseudo-lignitum**, Ettingshausen, 1893, pp. 136, 148; & 1895,
p. 15, pl. i, figs. 4, 5. Cretaceous; Australia.
- **razitanensis**, Hollick in Newberry, 1895, p. 65, pl. xlii, fig. 34.
? Amboy Clay; U.S.A.
- **revisenda**, Saporta, 1894, p. 201, pl. xxxvi, fig. 5.
Upper Albian; Portugal.
- **Rougoni**, Marion, 1890, p. 1054 [nomen nudum].
Turonian; Martigues, France.
- **Schenkiana**, Heer, 1871 A, p. 11, pl. iii, fig. 1.
Senonian; Quedlinburg, Saxony.
- **Schimperi**, Lesquereux, 1892, p. 66, pl. ii, fig. 12.
Dakota Group; Kansas, U.S.A.

- Myrica* ? *semina*, Lesquereux, 1874, p. 63, pl. xxvii, figs. 4, 4 a.
Dakota Group; Nebraska, U.S.A.
- *serrata*, Penhallow (*non* Velenovsky), 1907, p. 309, text-fig. 4.
Cretaceous; British Columbia, Canada.
- *serrata*, Velenovsky, 1883, p. 34, pl. x, figs. 1-8. (= *Myricophyllum serratum*, Velenovsky, 1889.)
Cenomanian; Bohemia.
- *Sternbergii*, Lesquereux, 1883, p. 35.
Dakota Group; Kansas, U.S.A.
- *thulensis*, Heer, 1871, p. 1183; & 1874 A, p. 107, pl. xxxi, fig. 1.
Atane Beds; Greenland.
- *Torreyi*, Lesquereux, 1873, p. 302; & 1878, p. 129, pl. xvi, figs. 3-10.
Montana Formation; Wyoming, U.S.A.
- ? *trifoliata*, Newberry, 1898, p. 37, pl. xiv, fig. 2.
Dakota Group; New Mexico.
- *vernassiensis*, Bozzi, 1891, p. 378, pl. xvi, fig. 5.
Cretaceous (?); Vernasso, Italy.
- *Zenkeri* (Ettingshausen), Heer, 1874 A, p. 108, pl. xxxi, fig. 2.
Atane Beds; Greenland.
- *sp.*, Hollick, 1906 A, p. 54, pl. vii, fig. 22.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- (fruit ?), Heer, 1882, p. 66, pl. xlv, fig. 26.
Cenomanian; Greenland.
- ? *sp.*, Knowlton, 1900 A, p. 34, pl. vi, fig. 4.
Montana Formation; Wyoming, U.S.A.
- ? *sp.*, Lesquereux, 1874, p. 63, pl. xxvii.
Dakota Group; Kansas, U.S.A.
- *sp.*, Velenovsky, 1883, p. 11, pl. v, figs. 6-8. [Catkins and fruit.]
Cenomanian; Bohemia.
- Myricæphyllum dentatum*, Fontaine, 1889, p. 316, pl. clvi, fig. 6.
Potomac Formation; Virginia, U.S.A.
- Myricanthium amentaceum*, Velenovsky, 1883, p. 11, pl. v, figs. 9-12.
Cenomanian; Vyserovic, Bohemia.
- Myricophyllum asplenioides*, Lange, 1890, p. 667, pl. xxxiv, figs. 1, 2.
Senonian; Wingertsberg, Rhenish Prussia.
- *glandulosum*, Velenovsky, 1889, p. 15, pl. vi, figs. 6-8.
Cenomanian; Bohemia.
- *longepetiolatum*, Ettingshausen, 1893, pp. 139, 148; & 1895, p. 16, pl. i, fig. 29.
Cretaceous; Australia.
- *serratum*, Velenovsky, 1889. (= *Myrica serrata*, Velenovsky, 1883.)
- Myrsine borealis*, Heer, 1874 A, p. 113, pl. xxxii, fig. 23; & 1882, p. 81, pl. xxiv, figs. 7 b, 8; pl. xxvii, fig. 1 b; pl. xlv, figs. 19, 20.
Atane Beds; Greenland.
- *caloneura*, Bayer, 1893, pp. 23-43, fig. 16; & Bayer in Fritsch, 1893, p. 131, text-fig. 188.
Senonian; Priesen, Bohemia.
- *crassa*, Lesquereux, 1892, p. 114, pl. lii, figs. 2, 3.
Dakota Group; Kansas, U.S.A.

- Myrsine elongata**, Newberry MS. in Hollick, 1894 A, p. 54, pl. clxxvii, fig. 2; & Newberry, 1895, p. 122, pl. xxii, figs. 1-3 (re-named *Myrsine Gaudini*, Berry, 1909.) Cretaceous; Long Island, U.S.A.
- **Gaudini** (Lesquereux), Berry, 1909, p. 263. (Re-naming *Myrsinites? Gaudini*, Lesquereux, 1892, and *Myrsine elongata*, Hollick, 1894 A.)
Baritan Formation; New Jersey, U.S.A.
- **manifesta**, Bayer, 1893, pp. 21, 43, fig. 15; & Bayer in Fritsch, 1893, p. 130, text-fig. 187. Senonian; Priesen, Bohemia.
- **oblongata**, Hollick in Newberry, 1895, p. 122, pl. xlii, fig. 15.
? Amboy Clay; U.S.A.
- Myrsinites? Gaudini**, Lesquereux, 1892, p. 115, pl. lii, fig. 4 (re-named *Myrsine Gaudini*, Berry, 1909).
Dakota Group; Kansas, U.S.A.
- Myrsinophyllum revisendum**, Saporta, 1894, p. 186, pl. xxxiv, fig. 10. Albian; Portugal.
- **varians**, Velenovsky, 1889, p. 25, pl. iv, figs. 8, 9; pl. v, fig. 12; pl. vi, figs. 10, 11. Cenomanian; Bohemia.
- **venulosum**, Saporta, 1894, p. 204, pl. xxxvii, fig. 15.
Upper Albian; Portugal.
- Myrtophyllum antiquorum**, Heer, 1871, p. 1184 [nomen nudum].
Atane Beds; Greenland.
- (**Eucalyptus?**) **Geinitzi**, Heer, 1869 A, p. 22, pl. xi, figs. 3, 4.
(= *Eucalyptus Geinitzi*, Heer, 1882.)
Cenomanian; Moletuin, Moravia.
- **latifolium**, Ettingshausen, 1893, p. 150; & 1895, p. 50, pl. iv, fig. 19. Cretaceous; Australia.
- **longifolium**, Velenovsky, 1882 A, p. 214 [nomen nudum].
Cretaceous; Bohemia.
- **parvulum**, Heer, 1882, p. 94, pl. xv, fig. 11; pl. xxi, fig. 4.
Atane Beds; Greenland.
- **pusillum**, Heer, 1871 A, p. 14, pl. iii, fig. 10.
Senonian; Quedlinburg, Saxony.
- (**Eucalyptus?**) **Schuebleri**, Heer, 1869 A, p. 23, pl. xi, fig. 2.
Cenomanian; Moletuin, Moravia.
- **Warderi**, Lesquereux, 1892, p. 136, pl. liii, fig. 10.
Dakota Group; Kansas, U.S.A.
- Nageiopsis acuminata**, Fontaine, 1889, p. 201, pl. lxxxv, fig. 11.
Potomac Formation; Virginia, U.S.A.
- **angustifolia**, Fontaine, 1889, p. 202, pl. lxxxvi, figs. 8, 9; pl. lxxxvii, figs. 2-6; pl. lxxxviii, figs. 1, 3, 4, 6-8; pl. lxxxix, fig. 2.
Ibid.
- **crassicaulis**, Fontaine, 1889, p. 193, pl. lxxix, figs. 2, 6; pl. lxxxii, fig. 1; pl. lxxxiv, figs. 3, 9, 11. Ibid.
- **decrescens**, Fontaine, 1889, p. 199, pl. lxxvii, fig. 3. Ibid.
- **heterophylla**, Fontaine, 1889, p. 201, pl. lxxxiv, fig. 4; pl. lxxxvi, figs. 6, 7; pl. lxxxviii, figs. 2, 5. Ibid.

Nageiopsis inaequilateralis, Fontaine, 1889, p. 200, pl. lxxxv, fig. 6.
(=*Podozamites inaequilateralis*, Berry, 1910 c.)

Potomac Formation; Virginia, U.S.A.

— **latifolia**, Fontaine, 1889, p. 198, pl. lxxxii, fig. 3. (= *Podozamites latifolius*, Berry, 1910 c.) Ibid.

— **longifolia**, Fontaine, 1889, p. 195, pl. lxxv, fig. 1; pl. lxxvi, figs. 2-6; pl. lxxvii, figs. 1, 2; pl. lxxviii, figs. 1-5; pl. lxxix, fig. 7; pl. lxxxv, figs. 1, 2, 8, 9. Ibid.

— **microphylla**, Fontaine, 1889, p. 201, pl. lxxxiv, fig. 6; pl. lxxxv, fig. 14; pl. lxxxvi, figs. 1-3, 5. Ibid.

— **montanensis**, Fontaine in Ward, 1905, p. 312, pl. lxxiii, fig. 10.
Kootanie Formation; Montana, U.S.A.

— **obtusifolia**, Fontaine, 1889, p. 200, pl. lxxxv, fig. 7.

Potomac Formation; Virginia, U.S.A.

— **ovata**, Fontaine, 1889, p. 199, pl. lxxvii, fig. 4; pl. lxxx, fig. 5.

Ibid.

— **recurvata**, Fontaine, 1889, p. 197, pl. lxxv, fig. 2; pl. lxxix, fig. 4; pl. lxxx, fig. 3. Ibid.

— **subfalcata**, Fontaine, 1889, p. 203, pl. clxviii, fig. 4. Ibid.

— **zamioides**, Fontaine, 1889, p. 196, pl. lxxix, figs. 1, 3; pl. lxxx, figs. 1, 2, 4; pl. lxxxi, figs. 1-6. Ibid.

Nathorstia angustifolia, Heer, 1880 b, p. 7, pl. i, figs. 1-6.

Kome Beds; Greenland.

— **firma**, Heer, 1880 b, p. 7, pl. i, fig. 7. (= *Danaeites firmus*, Heer.) Ibid.

— **latifolia**, Nathorst, 1908, p. 17, pl. iii. Cenomanian; Greenland.

Nathorstiana arborea, Richter, 1909, p. 3, pl. viii, figs. 1, 2, 3, 5, 8, 13 a (14); pl. x, figs. 11, 15. Senonian; Quedlinburg, Saxony.

— **gracilis**, Richter, 1909, p. 4, pl. viii, figs. 4, 6, 7, 9-12; pl. ix, figs. 3, 8 (9); pl. x, figs. 1-4, 6, 7 (12, 14, 16). Ibid.

— **squamosa**, Richter, 1909, p. 6, pl. ix, figs. 1, 2, 4-7, 10-13; pl. x, fig. 17. Ibid.

Nechalea fluitans, Debey in Murlon, 1881, p. 133 [nomen nudum].

Upper Cretaceous; Belgium.

— **lobata**, Debey, 1849, p. 299 [nomen nudum].

Senonian; Aix, Rhenish Prussia.

— **minor**, Debey in Murlon, 1881, p. 133 [nomen nudum].

Upper Cretaceous; Belgium.

— **petiolata**, Debey, 1849, p. 299 [nomen nudum].

Senonian; Aix, Rhenish Prussia.

— **serrata**, Debey, 1848 a, p. 115 [nomen nudum]. Ibid.

Nectandra imperfecta, Hollick, 1906 a, p. 76, pl. xxvii, figs. 13, 14.

Middle Cretaceous; Martha's Vineyard, U.S.A.

Negundoides acutifolia, Lesquereux, 1863, p. 101; & 1874, p. 97, pl. xxi, fig. 5. Dakota Group; Nebraska, U.S.A.

Nelumium arcticum, Heer, 1882, p. 92, pl. xl, fig. 6.

Atane Beds; Greenland.

— **Choffati**, Saporta, 1894 a, p. 836. Lower Cretaceous; Portugal.

- Nelumbium gallo-provinciale**, Saporta, 1882, p. 924.
Cretaceous of Fuveau; Trest, France.
- **lusitanicum**, Saporta, 1894 A, p. 835. Lower Albian; Portugal.
- **provinciale**, Saporta in Saporta & Marion, 1885, p. 125, text-fig. p. 128; & 1890, p. 5, pl. xii, fig. 1; pl. xiii, fig. 1; pl. xiv, figs. 1-6 a. Danian (?); France.
- **Saskatchuense**, Dawson, 1888, p. 35.
Belly River Series; Canada.
- **tenuifolium**, Lesquereux, 1873, p. 402; & 1878 B, p. 253, pl. xlv, fig. 3. Laramie Formation; Colorado, U.S.A.
- Nelumbo Dawsoni**, Hollick, 1894 D, p. 309 (re-naming *Brasenia antiqua*, Dawson, 1885). Belly River Series; Canada.
- **intermedia**, Knowlton, 1900 A, p. 53, pl. xiii, figs. 3-5.
Montana Formation; Wyoming, U.S.A.
- **Kempii**, Hollick, 1905 C, p. 412, pl. lxxiv, figs. 1, 2; pl. lxxv; pl. lxxvi; pl. lxxvii, fig. 1. (= *Serenopsis Kempii*, Hollick, 1893 B.)
Cretaceous; Long Island, U.S.A.
- **Laramiense**, Hollick, 1894 D, p. 307, text-fig. on p. 308.
Laramie Formation; Colorado, U.S.A.
- **primæva**, Berry, 1905 E, p. 75, pl. xliii, fig. 1.
Matawan Formation; New Jersey, U.S.A.
- ? **sp.**, Knowlton, 1900 A, p. 54, pl. xiii, fig. 6.
Montana Formation; Wyoming, U.S.A.
- Neomeris** *cf.* **annulus**, Parker & Jones in Böhm, 1903, p. 97.
Cenomanian; Guldtscha, Asia.
- (**Herouvalina**) **cretacea**, Steinmann, 1899, p. 200, text-figs. 42-46.
Upper Cretaceous; Mexico.
- Nerium Roehli**, von der Marck, 1864, p. 78, pl. xiii, fig. 3. 2-4.
Upper Senonian; Westphalia.
- Neuropteridium spinulosum**, Saporta, 1894, p. 79, pl. xvi, fig. 7 b.
Valanginian; Portugal.
- **torresianum**, Saporta, 1894, p. 79, pl. xviii, fig. 14. Ibid.
- Neuropteris angulata**, Newberry in Ives, 1861, p. 131, pl. iii, fig. 5.
Cretaceous; Arizona, U.S.A.
- **Castor**, Dawson, 1883, p. 24, pl. iv, figs. 14, 14 a.
Upper Cretaceous; Vancouver Island, Canada.
- **heterophylla cretacea**, Penhallow, 1902 B, p. 36.
Lower Cretaceous; Queen Charlotte Island, Canada.
- Neurosporangium foliaceum**, Debey & Ettingshausen, 1859 A, p. 190, pl. i, fig. 5. Senonian; Aix, Rhenish Prussia.
- **undulatum**, Debey & Ettingshausen, 1859 A, p. 192, pl. i, fig. 7. Ibid.
- Nicolia ægyptiaca**, Unger, 1859, p. 213, pl. i, figs. 1, 2; & 1866 A, p. 291, pl. i. Cretaceous or Tertiary; Egypt & Abyssinia.
- **Moresneti**, Hovelacque, 1889, p. 505; & 1890, p. 63, text-figs. 2-8, pl. iii, fig. 2. (= *Achenosaurus multidentis*, in part.)
Senonian; Moresnet, Belgium.

- Nicolia Owenii*, Carruthers, 1870 A, p. 310, pl. xiv, figs. 1, 2.
Cretaceous (?); Cairo.
- Nilssonia bohémica*, Velenovsky, 1885, p. 11, pl. ii, figs. 25-28;
pl. iii, fig. 12. Cenomanian; Bohemia.
- *californica*, Fontaine, 1905 A, p. 252, pl. lxvii, fig. 7.
Neocomian; California, U.S.A.
- *densinerve* (Fontaine), Berry, 1910 D, p. 639 (spelt *Nilsonia*;
= *Platypterigium densinerve*, Fontaine.)
Potomac Formation; Virginia, U.S.A.
- *Gibbsii* (Newberry), Hollick in Newberry, 1898, p. 16, pl. xv,
figs. 2, 2 a (re-naming *Tanopteris Gibbsii*, Newberry, 1863).
Cretaceous; Washington, U.S.A.
- *Johnstrupi*, Heer, 1882, p. 44, pl. vi, figs. 1-6.
Atane Beds; Greenland.
- *lata*, Dawson, 1883, p. 24, pl. iv, figs. 15 bis, 15 a.
Upper Cretaceous; Vancouver Island, Canada.
- *nigracollensis*, Wieland in Ward, 1905, p. 319, pl. lxxiii,
figs. 15 a-d. Dakota Group; S. Dakota, U.S.A.
- *oregonensis* (Fontaine), Berry, 1910 D, p. 637. (= *Sepindopsis*
oregonensis, Fontaine, 1905 A.)
Potomac Formation; Virginia, U.S.A.
- *orientalis*, Heer, internal anatomy discovered in Upper Cretaceous
of Japan by Stopes, 1910 A.
- *pasaytensis*, Penhallow, 1907, p. 307, text-fig. 3.
British Columbia, Canada.
- *polymorpha cretacea*, Penhallow, 1902 B, p. 42.
Lower Cretaceous; Queen Charlotte Island, Canada.
- *pterophylloides*, Yokoyama, 1894, p. 228, pl. xxii, figs. 8-10;
pl. xxv, fig. 7. Neocomian; Japan.
- ? *sambucensis*, Ward, 1905, p. 254, pl. lxvii, fig. 8.
Shasta Formation; California, U.S.A.
- *schaumbergensis* (Dunker), Nathorst, 1890, p. 45, pl. i, figs. 6-9;
& Yokoyama, 1894, p. 227, pl. xx, figs. 12, 14; pl. xxi, fig. 14;
pl. xxii, figs. 5-7. Neocomian; Japan.
- *Stantoni*, Ward, 1905, p. 251, pl. lxvii, figs. 5, 6.
Shasta Formation; California, U.S.A.
- Nipadites cretaceus*, Massalonge, 1857, p. 778 [nomen nudum].
Neocomian; Verona, Italy.
- *provincialis*, Saporta, 1868, p. 303, text-fig. 2.
Upper Cretaceous; France.
- Niponophyllum cordaitiforme*, Stopes & Fujii, 1910, pp. 16-23,
pl. iii, figs. 14-16, text-figs. 5-7.
Upper Cretaceous; Hokkaido, Japan.
- Nitella cosinensis*, Stache, 1889, p. 121, pl. iii, figs. 4, 5, 6.
Liburnian (Upper Cretaceous); Cosina.
- *devisita*, Stache, 1889, p. 136, pl. iii, figs. 56, a, b.
Upper Cretaceous (?); Corgnale.

- Nitella?** (*Chara*) **globulus**, Stache, 1889, p. 121, pl. i a, fig. 16.
Liburnian; Cosina.
- **robusta**, Stache, 1889, p. 136, pl. iv, fig. 7. (= *Chara robusta*,
Stache, 1880.) Upper Cretaceous (?); Divaca.
- **Stacheana** (Unger), Stache, 1889, p. 135, pl. iii, figs. 54, a, b, c, d;
pl. iv, figs. 1 a, b; pl. v, figs. 40 a, b. (= *Chara Stacheana*, Unger,
1860.) Upper Cretaceous (?); Cosina.
- **subimpressa**, Stache, 1889, p. 136, pl. iii, figs. 55 a, b.
Upper Cretaceous (?); Trieste.
- Noeggerathiopsis Robinsi**, Dawson, 1894, p. 56, pl. vi, fig. 7.
Upper Cretaceous; Vancouver Island, Canada.
- Nordenskiöldia borealis**, Heer, 1870 A, p. 65, pl. vii, figs. 1-13.
Recorded Lesquereux, 1892, p. 219, in Dakota Group, Kansas.
- Nothopogoxylon scalariforme**, Gothan, 1908 A, p. 20, pl. ii, figs. 14
-18. Cretaceous (?); Antarctica.
- Nullipora gracilis**, Reuss, 1846, p. 66, pl. xvi, figs. 4-6.
Upper Cretaceous; Bohemia.
- Nulliporites granulosus**, Heer, 1865, p. 190, text-fig. 103; & 1877,
p. 143, pl. lviii, figs. 10, 11. Cretaceous; Switzerland.
- Nyssa Buddiana**, Ward, 1887, p. 53, pl. xlvii, fig. 7.
Laramie Formation; Wyoming, U.S.A.
- **cuneatus**, Newberry, 1863, p. 524; & 1898, p. 125, pl. xvii, figs. 4-6.
Cretaceous (?); Washington, U.S.A.
- **Snowiana**, Lesquereux, 1892, p. 126, pl. lii, fig. 11.
Dakota Group; Kansas, U.S.A.
- **vetusta**, Newberry, 1870, p. 11; & 1878, pl. ii, fig. 2 (as *Magnolia*
obovata); & 1898, p. 125, pl. i, fig. 2; pl. iv, fig. 4.
Dakota Group; Nebraska, U.S.A.
- Octea nassauensis**, Hollick, 1906 A, p. 76, pl. xxvii, fig. 8.
Middle Cretaceous; Long Island, U.S.A.
- Olea?** **myricoides**, Saporta, 1894, p. 220, pl. xxxix, fig. 25.
Cenomanian; Portugal.
- Oleandra arctica**, Heer, 1874 A, p. 38, pl. xii, figs. 3-11; pl. xxi, fig. 2 b.
Kome Beds; Greenland.
- **graminaefolia**, Knowlton, 1907, p. 113, pl. xi, figs. 5, 5 A, 6, 6 A.
Kootanie Formation; Montana, U.S.A.
- Oleandrium tenerum**, Saporta, 1890 A, p. 814; & 1894, p. 85, pl. xv,
fig. 3; pl. xvi, fig. 18. Valanginian; Portugal.
- Oncopteris Kauniciana** (Dormitzer), Velenovsky, 1888 B, p. 22, pl. v,
fig. 1. (= *Alsophilina Kauniciana*, Dormitzer, 1853.)
Perucer Beds; Bohemia.
- **Nettwalli**, Dormitzer in Krejci, 1853, p. 28, pl. ii.
Cenomanian; Bohemia.
- Onoclea inquirenda**, Hollick, 1906 A, p. 32, pl. i, figs. 1-7. (= *Caulin-*
ites inquirendus, Hollick, 1905 c.)
Magothy Formation; Long Island, U.S.A.

- Onoclea minima**, Knowlton, 1899 B, p. 656, pl. lxxvii, figs. 11-15.
Laramie Formation; Yellowstone Park, U.S.A.
- **sp.**, Nathorst, 1891, p. 24, pl. i, figs. 5-7.
Mecklenburg, nr. Rostock.
- Onychiopsis capsulifera** (Velenovsky), Nathorst, 1890, p. 55. (= *Thyrsopteris capsulifera*, Velenovsky, 1888.)
Peruczer Beds; Bohemia.
- **elegans**, Yokoyama, 1894, p. 215, pl. xxviii, figs. 7, 7 a.
Neocomian (?) ; Japan.
- **elongata** (Geyler), Yokoyama, 1889, recorded by Krasser, 1896, p. 121, pl. vii, fig.
Cenomanian; Kunstadt, Moravia.
- **Mantelli** (Brongniart), Seward, 1894, p. 41. (= *Sphenopteris Mantelli*, Brongn., 1828.)
Recorded Potomac Formation, Maryland, U.S.A., Fontaine, 1889, p. 91.
- **psilotoides** (Stokes & Webb), Ward, 1905, p. 155, and recorded p. 518, pl. cxiii, fig. 1. (= *O. Mantelli*, Seward, 1894.)
Older Potomac Formation; Virginia, U.S.A.
- Opegrapha antiqua**, Lesquereux, 1873, p. 390; & 1878 B, p. 36, pl. i, figs. 1-1 c.
Laramie Formation; Wyoming, U.S.A.
- Opegraphites striato-punctatus**, Debey, 1848 A, p. 116 [nomen nudum; cancelled, Debey, 1850].
Senonian; Aix, Rhenish Prussia.
- Ophioglossum granulatum**, Heer, 1883 A, p. 8, pl. lvii, figs. 8-9.
Patoot Beds; Greenland.
- Oreodaphne cretacea**, Lesquereux, 1874, p. 84, pl. xxx, fig. 5.
Dakota Group; Kansas, U.S.A.
- Oreodoxites plicatus**, Lesquereux, 1883, p. 122, pl. xviii, figs. 1-4.
Laramie Formation; Colorado, U.S.A.
- Osmunda arctica**, Heer, 1883 A, p. 7, pl. xlix, figs. 4-7; pl. i, figs. 6-8.
Patoot Beds; Greenland.
- **delawarensis**, Berry, 1906 B, p. 164, pl. viii, figs. 2-4.
Magothy Formation; Delaware, U.S.A.
- **Dicksonioides**, Fontaine, 1889, p. 146, pl. xli, fig. 5; pl. lviii, fig. 9; pl. lix, figs. 1, 4, 8, 9, 11; pl. lx, figs. 2, 4, 5, 9; pl. lxi, figs. 1, 2.
Potomac Formation; Virginia, U.S.A.
- **Dicksonioides latipennis**, Fontaine, 1889, p. 147, pl. lx, figs. 1, 3; pl. lxi, fig. 3.
Ibid.
- **Gerini**, Saporta, 1890 B, p. 183, pl. xi, fig. 1.
Upper Cretaceous; France.
- **haldemiana**, Hosius & von der Mark, 1880, p. 140, pl. xxv, fig. 18.
Upper Senonian; Haldem, Westphalia.
- **major**, Lesquereux, 1883, p. 121, pl. xviii, fig. 5.
Laramie Formation; Colorado, U.S.A.
- **montanensis**, Knowlton in Stanton & Hatcher, 1905, p. 129, pl. xiv, fig. 6.
Judith River Beds; Montana, U.S.A.
- **novæ-caesaræ**, Berry, 1907 A, p. 671, text-fig. 2.
Magothy Formation; New Jersey, U.S.A.
- **Oebergiana**, Heer, 1874 A, p. 98, pl. xxvi, figs. 9, 9 b; pl. xxxii, fig. 7 a.
Atane Beds; Greenland.

- Osmunda petiolata*, Heer, 1874 A, p. 57, pl. iii, figs. 1 c, 2 b; pl. xx, fig. 10 c. Kome Beds; Greenland.
- *retinenda*, Saporta, 1894, p. 107, pl. xix, figs. 16-17. Lower Cretaceous (?); Portugal.
- *sphenopteroides*, Fontaine, 1889, p. 145, pl. xxv, fig. 13. Potomac Formation; Virginia, U.S.A.
- Osmundites skidegatensis*, Penhallow, 1902 A, p. 3, pls. i-v, text-figs. 1, 3 on pp. 6, 10; & 1902 B, p. 52, pls. vii-xi. Lower Cretaceous; Queen Charlotte Island, Canada.
- Osmundophyllum cretaceum*, Velenovsky, 1889, p. 6, pl. ii, fig. 21. Cenomanian; Bohemia.
- Otozamites* (?) *grönlandicus*, Heer, 1874 A, p. 99, pl. xxvi, fig. 2. Atane Beds; Greenland.
- Pachyphyllum Heerianum*, Saporta, 1894, p. 108, pl. xix, fig. 25. Lower Cretaceous (?); Portugal.
- Pachypteris cretacea*, Debey, 1849, p. 299 [nomen nudum]. Senonian; Aix, Rhenish Prussia.
- *dalmatica*, Kerner, 1896, p. 47, pl. ii; pl. iii, figs. 1, 4; pl. v, fig. 10. Cenomanian; Dalmatia.
- *dalmatica dentata*, Kerner, 1896, p. 47, pl. iii, fig. 3. Ibid.
- *dimorpha*, Kerner, 1896, p. 49, pl. iii, fig. 2; pl. v, fig. 8. Ibid.
- Pagiophyllum dubium*, Fontaine, 1894, p. 271, pl. xxxix, figs. 2-11. Trinity Division; Texas, U.S.A.
- *sp.*, Dawson, 1893, p. 90, text-fig. 14. Kootanie Formation; North-West Territory, Canada.
- Palæanthus* (*Williamsonia*) *problematica*, Newberry, 1895, p. 125, pl. xxxv, figs. 1-9. (= *Williamsonia problematica* (Newby.), Ward, 1895.) Amboy Clay; New Jersey, U.S.A.
- Palæobromelia Jugleri*, Ettingshausen, 1852 B, p. 3, pl. i, fig. 1; pl. ii, figs. 1-3. [Probably egg-cases of fishes (Stopes).] Wealden; Hanover.
- Palæocassia angustifolia*, Ettingshausen, 1867 A, p. 261, pl. iii, figs. 6-7. (= *Cassia angusta*, Heer, 1882.) Cenomanian; Niederschoena, Saxony.
- *lanceolata*, Ettingshausen, 1867 A, p. 262, pl. i, fig. 8; pl. iii, fig. 8. Ibid.
- *laurinea*, Lesquereux, 1892, p. 147, pl. lxiv, fig. 12. Dakota Group; Kansas, U.S.A.
- *phaseolitoides*, Ettingshausen, 1887 A, p. 189, pl. ix, fig. 17. (= *Cassia præ-phaseolitoides*, Ettingshausen, 1895.) Upper Cretaceous; New Zealand.
- Palæocyparis flexuosa*, Saporta, 1894, p. 108, pl. xix, figs. 19-20, 26; pl. xx, figs. 1-5. Lower Cretaceous (?); Portugal.
- *obscura*, Saporta, 1894, p. 178, pl. xxxiv, fig. 8. Albian; Portugal.
- Palæohillia arkansana*, Knowlton, 1895, p. 387, text-figs. 1-3. Trinity Division; Arkansas, U.S.A.

- Palæochillia* sp., Holm, 1896, p. 207, pl. xvii, figs. 1, 2. [Structure of *P. arkansana*.] Trinity Division; Arkansas, U.S.A.
- Palæolepis bicornuta*, Saporta, 1894, p. 179, pl. xxxiii, fig. 4 c. Albain; Portugal.
- *cheiromorpha*, Saporta, 1894, p. 215, pl. xxxix, figs. 15-16. Cenomanian; Portugal
- *emarginata*, Saporta, 1894, p. 179, pl. xxxiii, fig. 15. Albain; Portugal.
- *multipartita*, Saporta, 1894, p. 215, pl. xxxix, figs. 17-18. Cenomanian; Portugal.
- Palæostrobus crassipes*, Renger, 1866, p. 137, pl. i, figs. 2, 3, 4, 5. (= *Dammartites crassipes*, Goepfert.) Upper Cretaceous; Horice.
- *mirabilis*, Renger, 1866, p. 134, pl. i, fig. 1. (= *Krannera mirabilis*, Corda.) Upper Cretaceous; Novy Bydzov.
- Palæospathe Sarthacensis*, Crié, 1879 (?), p. 20. Cretaceous; Sainte-Croix, France.
- Paleodictyon giganteum*, Peruzzi, 1881, p. 1, pl. i, fig. 3. Cretaceous; Italy.
- Palinosphaeria* sp., Reinsch, 1905, p. 402, text-fig. 1. Upper Cretaceous.
- Paliurus affinis*, Heer, 1883 A, p. 42, pl. lxii, figs. 16-19. Patoot Beds; Greenland.
- *anceps*, Lesquereux, 1892, p. 166, pl. xxxv, fig. 4. Dakota Group; Kansas, U.S.A.
- *cretaceus*, Lesquereux, 1892, p. 165, pl. xxxv, fig. 3. Ibid.
- *integrifolius*, Hollick, 1894 A, p. 57, pl. clxxvii, figs. 5, 8, 12. Cretaceous; Long Island, U.S.A.
- *membranaceus*, Lesquereux, 1868, p. 101; & 1874, p. 108, pl. xx, fig. 6. Dakota Group; Nebraska, U.S.A.
- *minimus*, Knowlton, 1899 B, p. 659, pl. lxxvii, figs. 7-9. Laramie Formation; Yellowstone National Park, U.S.A.
- *montanus*, Dawson, 1886, p. 14. Mill Creek Series; Canada.
- *Neillii*, Dawson, 1894, p. 62, pl. xi, figs. 44, 45. Upper Cretaceous; Vancouver Island, Canada.
- *obovatus*, Lesquereux, 1892, p. 165, pl. xxxv, fig. 6. Dakota Group; Kansas, U.S.A.
- *ovalis*, Dawson, 1886, p. 14, pl. iv, figs. 4, 8. Mill Creek Series; Canada.
- *populiferus*, Berry, 1906 B, p. 177; & 1906 B, p. 153, pl. xx, fig. 1. Magothy Formation; New Jersey, U.S.A.
- *zizyphoides*, Lesquereux, 1873, p. 397; & 1878 B, p. 274, pl. li, figs. 1-6. Laramie Formation; Wyoming, U.S.A.
- Palmacites canaliculatus*, Schlotheim (*non* Heer), 1820, p. 326, pl. xvi, fig. 2. Quadersandstein; Gotha, Germany.
- *horridus*, Heer, 1869 A, p. 15, pl. v, fig. 1. Cenomanian; Moletein, Moravia.
- *obsoletus*, Schlotheim, 1820, p. 396, pl. xvi, fig. 3. Quadersandstein; Gotha, Germany.

- Palmacites Reichi*, Geinitz, 1850 A, p. 270.
Quadersandstein; Dittersbach.
- *rimosus*, Heer, 1876 B, p. 11, figs. 21, 22 [fruits].
Upper Cretaceous; Egypt.
- *varians*, Corda in Reuss, 1846, p. 87, pl. xlvii, figs. 7-9. (= *Fasciculites varians*, Unger, considered to be *Protopteris Sternbergi*, Corda, by Feistmantel, 1872.)
Pläner; Bohemia.
- *Zittelii*, Schenk, 1880 A, p. 657 [nomen nudum].
Nubian Sandstone; Cairo.
- ? *sp.*, Otto, 1852, p. 9, pl. iv, figs. 4, 5.
Quadersandstein; Saxony.
- Palmocarpus cretaceus*, Miquel, 1853, p. 51, pl. vii.
Senonian; Limburg.
- Palmophyllum moleteianum*, Krasser & Kubart, 1906 B, p. 47 [nomen nudum].
Cenomanian; Moletein, Moravia.
- Palmoxylon Andegavense*, Crié, 1892, p. 101, unnumb. pl., fig. 4.
Turonian (?); France.
- *angiorhiza*, Stenzel, 1904, p. 267, pl. xxiv, figs. 287-289.
Cenomanian (?); Tlacolula, Mexico.
- *Aschersoni*, Schenk, 1883, p. 6, pl. v, fig. 4.
Cretaceous (?); Egypt.
- *astron*, Stenzel, 1904, p. 257, pl. xx, figs. 208-213.
Cenomanian (?); Tlacolula, Mexico.
- *astron radicum*, Stenzel, 1904, p. 260, pl. xx, figs. 214-222.
Ibid.
- *astron verum*, Stenzel, 1904, p. 258, pl. xx, figs. 208-213.
Ibid.
- *Boxbergæ* (Geinitz), Stenzel, 1904, p. 195, pl. ix, figs. 75-78.
Turonian; France.
- *cheyennense*, Wieland, 1903 A, p. 216 [nomen nudum].
Upper Cretaceous; Fort Pierre, S. Dakota, U.S.A.
- *Cottæ libycum*, Stenzel, 1904, p. 221, pl. xv, figs. 136, 137.
Upper Senonian; Egypt.
- *Lingerinum*, Crié, 1892, p. 99, unnumb. pl., figs. 1-3.
Turonian (?); France.
- *parvifasciculosum*, Vater, 1884, p. 830, pl. xxvii, figs. 2-3.
Lower Senonian; Harzburg.
- *parvifasciculatum*, Schenk, 1890, p. 888. (Same as *P. parvifasciculosum*.)
- *radiatum*, Vater, 1884, p. 831, pl. xxvii, fig. 4.
Lower Senonian (?); Helmstedt.
- *scleroticum*, Vater, 1884, p. 829, pl. xxvii, fig. 1.
Lower Senonian (?); Brunswick.
- *tenue*, Stenzel, 1904, p. 228, pl. xvi, figs. 145-151.
Cenomanian (?); Tlacolula, Mexico.
- *texense*, Stenzel, 1904, p. 185, pl. viii, figs. 61-63.
Cenomanian (?); Texas, U.S.A.
- *variabile*, Vater, 1884, p. 832, pl. xxvi, fig. 5.
Lower Senonian (?); Helmstedt.

- Palmoxylon variabile verum*, Stenzel, 1904, p. 238, pl. xviii, figs. 188-196; pl. xix, fig. 197. Lower Senonian; North Germany.
- *Zitteli*, Schenk, 1883, p. 5, pl. ii, figs. 5-6. Cenomanian (?); Libyan Desert.
- Panax cretacea*, Heer, 1871, p. 1183; & 1874 A, p. 114, pl. xxxii, figs. 9, 10. Atane Beds; Greenland.
- *globulifera*, Heer, 1883 A, p. 35, pl. lxii, fig. 8 b. Patoot Beds; Greenland.
- *macrocarpa*, Heer, 1883 A, p. 35, pl. lxii, fig. 8. Ibid.
- Pandanus Aquisgranensis*, Debey in Mourlon, 1881, p. 133 [nomen nudum]. Senonian; Limburg.
- *austriacus*, Ettingshausen, 1852, p. 492, pl. xxiii, fig. 1. Senonian; Austria.
- *pseudo-inermis*, Ettingshausen, 1852, p. 493, pl. xxiv, figs. 1-2; pl. xxv, fig. 1. Ibid.
- *Simildæ*, Stiehler, 1857, p. 454; & 1858, p. 75, pl. xiv. Senonian; Quedlinburg, Saxony.
- *trinervis*, Ettingshausen, 1852, p. 494, pl. xxvi, fig. 1. Senonian; Austria.
- Paracallipteris Potonie*, Richter, 1904, p. 17, pl. i, fig. 13; & 1905, p. 6, pl. i, fig. 13. Senonian; Quedlinburg, Saxony.
- Paracedroxylon scituate*, Sinnott, 1909, p. 171, pls. lxxx, lxxxi. Cretaceous (?); Massachusetts, U.S.A.
- Paracredneria Fritschii*, Richter, 1905, p. 15, pl. ii, fig. 14; pl. iii, fig. 9. Upper Cretaceous; Westerhausen.
- Parathinnfeldia dubia*, Richter, 1904, p. 14, pl. i, fig. 3; & 1905, p. 5, pl. i, fig. 3. Senonian; Quedlinburg, Saxony.
- Parrotia Canfieldi*, Lesquereux, 1892, p. 141, pl. xxx, fig. 6. Dakota Group; Kansas, U.S.A.
- *grandidentata*, Lesquereux, 1892, p. 140, pl. xxxix, figs. 2-4. Ibid.
- ? *Winchelli*, Lesquereux, 1892, p. 140, pl. xxix, figs. 5, 6. Dakota Group; Minnesota, U.S.A.
- Passiflora antiqua*, Newberry, 1895, p. 109, pl. xxiii, fig. 7. Amboy Clay; Woodbridge, U.S.A.
- Pecopteris Andersoniana*, Heer, 1874 A, p. 41, pl. iii, figs. 7, 7 b. Kome Beds; Greenland.
- *angustipennis*, Fontaine, 1889, p. 87, pl. xxi, fig. 10. Potomac Formation; Virginia, U.S.A.
- *arctica*, Heer, 1868, p. 80, pl. i, fig. 13; pl. xliii, fig. 5. Kome Beds; Greenland.
- *arenaria*, Hampe, 1855, p. 12 [nomen nudum]. Senonian; Quedlinburg, Saxony.
- *argutula*, Heer, 1871, p. 1182; & 1874 A, p. 96, pl. xxvi, figs. 8, 8 b. Atane Beds; Greenland.
- *atyrkanensis*, Heer, 1878, p. 29, pl. viii, figs. 9-17. Lower Cretaceous; Siberia.

- Pecopteris Bayeana**, Heer, 1871, p. 1181 [nomen nudum].
Kome Beds; Greenland.
- **bohémica**, Corda in Reuss, 1846, p. 95, pl. xlix, fig. 1.
Quadersandstein; Bohemia.
- **Bolbroeana**, Heer, 1871, p. 1181; & 1874 A, p. 41, pl. iii, fig. 6.
Kome Beds; Greenland.
- **borealis**, Brongniart, 1828, p. 351, pl. cxix, fig. 12; & Heer, 1868, p. 81, pl. i, fig. 14; pl. xlv, figs. 5, a, b. Ibid.
- **brevipennis**, Fontaine, 1889, p. 86, pl. xxi, figs. 1-3.
Potomac Formation; Virginia, U.S.A.
- **Browniana**, Dunker. (= *Cladophlebis Browniana*, recorded from Canadian Kootanie, Dawson, 1893.)
- **Buftoni**, Johnston, 1896, p. 60, fig. 3. Cretaceous(?); Tasmania.
- **Choffatiana**, Heer, 1881, p. 15, pl. xi, figs. 9, 9 b.
Cretaceous; Portugal.
- **constricta**, Fontaine, 1889, p. 89, pl. xx, figs. 1, 2, 4.
Potomac Formation; Virginia, U.S.A.
- **cuspidata**, Schulze, 1888, p. 26 [nomen nudum].
Senonian; Heimbürg, Germany.
- **cycloloba**, Newberry in Ives, 1861, p. 130, pl. iii, figs. 3, 4, 4 a.
Base of Cretaceous(?); Arizona, U.S.A.
- **denticulata**, Heer, 1874 A, p. 95, pl. xxvi, figs. 7, 7 b.
Atane Beds; Greenland.
- **dilacerata**, Saporta, 1894, p. 78, pl. xviii, fig. 13.
Valanginian; Portugal.
- **dispersa**, Saporta, 1894, p. 170, pl. xxviii, fig. 13.
Albian; Portugal.
- **Geinitzii**, var., Fontaine in Weed & Pirsson, 1898, p. 481.
Kootanie Formation; Montana, U.S.A.
- **Geyleriana**, Nathorst, 1890, p. 48, pl. iv, fig. 1; pl. vi, fig. 1.
Lower Cretaceous; Japan.
- *cf.* **Haidingeri** (Debey & Ettingshausen), Zeiller, 1905, p. 331, pl. vii, fig. 6. Upper Cretaceous; Seltzi Izvor, Balkans.
- **hyperborea**, Heer, 1868, p. 81, pl. xlv, fig. 4.
Kome Beds; Greenland.
- **incerta**, Debey, 1848 A, p. 117 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **komensis**, Heer, 1882, p. 6, pl. iii, fig. 5. Kome Beds; Greenland.
- **kudlisetensis**, Heer, 1874 A, p. 97, pl. xxvi, fig. 18.
Upper Cretaceous; Greenland.
- **latiloba**, Heer, 1878, p. 28, pl. viii, figs. 5-6.
Lower Cretaceous(?); Siberia.
- **lobifolia**, Corda (*non* (Phill.) Lindley & Hutton) in Reuss, 1846, p. 95, pl. xlix, figs. 4, 5. Lower Quader; Bohemia.
- **microdonta**, Fontaine, 1889, p. 85, pl. xix, fig. 8; pl. xx, figs. 5, 11.
Potomac Formation; Virginia, U.S.A.
- **minor**, Velenovsky, 1888 n, p. 19, pl. iii, fig. 18.
Perucer Beds; Bohemia.

- Pecopteris minutula*, Saporta, 1894, p. 170, pl. xviii, fig. 14.
Albian; Portugal.
- *montanensis*, Fontaine, 1893, p. 492, pl. lxxxiii, figs. 1, 1 a.
Kootanie Formation; Montana, U.S.A.
- *nebraskana*, Heer, 1868, p. 332, text-fig. 8 (re-named *Cyatheites?*
nebraskana, Knowlton, 1898). Dakota Group; Nebraska, U.S.A.
- *Nordströmi*, Heer, 1871, p. 1182 [nomen nudum].
Atane Beds; Greenland.
- *Oebergiana*, Heer, 1871, p. 1182 [nomen nudum]. Ibid.
- *osmundacea*, Schulze, 1888, p. 26 [nomen nudum].
Senonian; Heimbürg, Germany.
- *ovatodentata*, Fontaine, 1889, p. 85, pl. xv, fig. 8; pl. xxii, fig. 12;
pl. xxiii, fig. 1. Potomac Formation; Virginia, U.S.A.
- *pachyphylla*, Fontaine, 1889, p. 88, pl. xxvi, figs. 4, 5. Ibid.
- *Pfaffiana*, Heer, 1871, p. 1182; & 1874 a, p. 95, pl. xxvi, fig. 5.
Atane Beds; Greenland.
- *polypodioides*, Debey (*non* Brongniart), 1849, p. 299 [nomen
nudum]. Senonian; Aix, Rhenish Prussia.
- *Reichiana*, Brongniart (*non* Goeppert), Schimper, *nec* Presl, 1828,
p. 302, pl. cxvi, fig. 7. Cenomanian; Niederschoena, Saxony.
- *Reichiana*, Presl in Sternberg, 1838, p. 155, pl. xxxvii, fig. 2.
Quadersandstein; Bavaria.
- *reticulata*, Stokes & Webb, 1824, p. 424, pl. xlv, fig. 5; pl. xlvii,
fig. 3. (= *Weichselia reticulata* (Stokes & Webb), Ward, 1899.)
Wealden; Tilgate Forest.
- *Schonæ*, Reich in Cotta, 1836, p. 586.
Quadersandstein; Freiberg, Saxony.
- *serrulata*, Heer (*non* Hartt, *nec* (Lesquereux) Schimper), 1871,
p. 1182 [nomen nudum]. Atane Beds; Greenland.
- (*Pteris?*) *socialis*, Heer, 1882, p. 34, pl. vii, fig. 4; pl. viii, fig. 15;
pl. xxxii, fig. 9. Ibid.
- *striata*, Presl in Sternberg, 1838, p. 155, pl. xxxvii, figs. 3, 4.
Quadersandstein; Bavaria.
- *strictinervis*, Fontaine, 1889, p. 84, pl. xiii, figs. 6-8; pl. xix,
fig. 9; pl. xx, fig. 3; pl. xxii, fig. 13; pl. clxx, figs. 5, 6.
Potomac Formation; Virginia, U.S.A.
- *tenella*, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- *virginiensis*, Fontaine, 1889, p. 82, pl. viii, figs. 1-7; pl. ix,
figs. 1-6; pl. xxiv, fig. 2; pl. clxix, fig. 3.
Potomac Formation; Virginia, U.S.A.
- *Zippei*, Corda in Reuss, 1846, p. 95, pl. xlix, figs. 2, 3. (= *Gleichenia*
Zippei, Heer, 1868 = *Mertensia Zippei*, Engelhardt, 1892 =
Gleichenites Zippei, Seward, 1910.) Quader; Bohemia.
- *sp.*, Dawson, 1883, p. 25.
Upper Cretaceous; Vancouver Island, Canada.
- *sp.*, Heer, 1878, p. 29, pl. viii, figs. 17 c, 17 d
Lower Cretaceous; Siberia.

- Pecopteris* sp., Zeiller, 1905, p. 333, pl. vii, figs. 2, 3.
Upper Cretaceous; Balkans.
- sp., Zeiller, 1905, p. 335, pl. vii, figs. 7, 7 a.
Upper Cretaceous; Balkans.
- (*Cladophlebis*?) sp., Zeiller, 1905, p. 335, pl. vii, fig. 4. Ibid.
- Periploca cretacea*, Hollick, 1906 A, p. 105, pl. xl, fig. 16.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Persea Hayana*, Lesquereux, 1892, p. 103, pl. xvi, fig. 6.
Dakota Group; Kansas, U.S.A.
- *Lecconteana*, Lesquereux, 1874, p. 75, pl. xxviii, fig. 1. (= *Sassafras*?
Lecconteanum, Lesquereux, 1869, and *Quercus Benzoin*, Lesquereux,
1859.) Ibid.
- *nebrascensis*, Lesquereux, 1869, p. 431, pl. xxiii, figs. 9, 10.
(= *Laurus nebrascensis*, Lesquereux, 1868.)
Dakota Group; Nebraska, U.S.A.
- *Schimperii*, Lesquereux, 1892, p. 103, pl. xvi, fig. 5.
Dakota Group; Kansas, U.S.A.
- *Sternbergii*, Lesquereux, 1874, p. 76, pl. vii, fig. 1. (= *Ficus Stern-*
bergii, Lesquereux, 1873.) Ibid.
- *Suessi*, Krasser in Krasser & Kubart, 1906 B, p. 47 [nomen nudum].
Cenomanian; Moletuin, Moravia.
- *valida*, Hollick, 1906 A, p. 76, pl. xxix, figs. 8, 9.
Middle Cretaceous; Long Island, U.S.A.
- Perseophyllum Hauthalianum*, Kurtz, 1902, p. 52.
Equivalent of Dakota Group; Patagonia.
- Persoonia Lesquereuxii*, Knowlton in Lesquereux, 1892, p. 89, pl. xx,
figs. 10-12. Dakota Group; Kansas, U.S.A.
- *spatulata*, Hollick in Newberry, 1895, p. 71, pl. xlii, fig. 14.
? Amboy Clay; U.S.A.
- Petrosphaeria japonica*, Stopes & Fujii, 1910, pp. 4-6, pl. i, figs. 1-6.
Upper Cretaceous; Hokkaido, Japan.
- Peuce aquisgranensis*, Endlicher, 1847 B, p. 294. (= *Pinites aquis-*
granensis, Goeppert, 1842 B.) Senonian; Aix, Rhenish Prussia.
- *cretacea*, Endlicher, 1847 B, p. 32. (= *Pinites cretacea*, Corda,
1846.) Greensand; Bohemia.
- Peucedanites antiquissimus*, Saporta, 1890 B, p. 180, text-fig. 35.
Albian (?); Portugal.
- *primordialis*, Saporta, 1894, p. 194, pl. xxxv, fig. 13. Ibid.
- Phacidites communis* (Feistmantel), Meschinelli, 1892, p. 779.
(= *Phacidium commune*, Feistmantel, 1874.)
Perucur Beds; Bohemia.
- Phacidium circumscriptum*, Bayer, 1899, p. 7, text-fig. 3. Ibid.
- *commune*, Feistmantel, 1874, pp. 262, 266. (= *Phacidites*
communis, Meschinelli, 1892.) Ibid.
- *myrtophylli*, Engelhardt, 1892 A, p. 80, pl. ii, fig. 10 a, b.
Quader; Freiberg, Saxony.
- *Palæccassiae*, Ettingshausen, 1867 A, p. 242, pl. i, figs. 8, 8 b.
Cenomanian; Niederschoena, Saxony.

- Phaseolites elegans*, Hollick, 1906 A, p. 85, pl. xxxii, fig. 4.
Middle Cretaceous; Long Island, U.S.A.
- *formus*, Lesquereux, 1892, p. 147, pl. lv, figs. 5, 6, 12.
Dakota Group; Kansas, U.S.A.
- *Manhassetensis*, Hollick, 1905 c, p. 414, pl. lxxviii, figs. 1-2.
Cretaceous; Long Island, U.S.A.
- Phegopteris Grothiana*, Heer, 1883 A, p. 3, pl. xlviii, figs. 12-13.
Patoot Beds; Greenland.
- *Jörgenseni*, Heer, 1882, p. 32, pl. xxxv, figs. 1-3.
Atane Beds; Greenland.
- *Kornerupi*, Heer, 1883 A, p. 3, pl. xlix, fig. 3.
Patoot Beds; Greenland.
- Phillyria Engelhardti*, Velenovsky, 1887, p. 68, pl. xxvii, figs. 2-5.
Cenomanian; Bohemia.
- Phlebomeris? falciformis*, Saporta, 1894, p. 84, pl. xviii, fig. 8 &
p. 169, pl. xxx, figs. 2-4. Valanginian & Albian; Portugal.
- *spectanda*, Saporta, 1894, p. 168, pl. xxix, fig. 14; pl. xxx, fig. 1.
Albian; Portugal.
- *Wilkommi*, Saporta, 1894, p. 168, pl. xxx, fig. 5. Ibid.
- Phlebopteris dubia*, Richter, 1899 A, p. 40.
Neocomian; Quedlinburg, Saxony.
- Phragmites? Cliffwoodensis*, Berry, 1905 e, p. 99, pl. xlvii, fig. 5.
Matawan Formation; New Jersey, U.S.A.
- *cordaiformis*, Dawson, 1883, p. 26, pl. v, fig. 22.
Upper Cretaceous; Vancouver Island, Canada.
- *cretaceus*, Lesquereux, 1874, p. 55, pl. i, figs. 13, 14; pl. xxix,
fig. 7. (= *Arundo cretaceus*, Lx., 1868.)
Dakota Group; Kansas, U.S.A.
- *falcata*, Knowlton, 1899 b, p. 658, pl. lxxviii, fig. 5.
Laramie Formation; Yellowstone Park, U.S.A.
- *laramianus*, Cockerell, 1909, p. 141.
Laramie Formation; Colorado, U.S.A.
- *Prattii*, Berry, 1910 A, p. 191. (= *Phragmites* sp., Berry, 1907,
p. 190, pl. xi, fig. 5.) Middle Cretaceous; North Carolina, U.S.A.
- *Rumpfii*, Stur, 1871, p. 503. Cretaceous; Piber, Austria.
- sp., Berry, 1907, p. 190, pl. xi, fig. 5. (= *Phragmites Prattii*, Berry,
1910 A.) Middle Cretaceous; N. Carolina, U.S.A.
- Phycodes sericeus*, Debey & Ettingshausen, 1859 A, p. 200.
Senonian; Aix, Rhenish Prussia.
- Phyllites actinoneuron*, Ettingshausen, 1893, p. 151; & 1895, p. 54,
pl. iv, fig. 24. Cretaceous; Australia.
- *acuminatus*, Goepfert, 1847, p. 361 (naming fig. 4, pl. I of
Goepfert, 1842 A). Quadersandstein; Silesia.
- *amissus*, Lesquereux, 1892, p. 217, pl. lxii, fig. 1.
Dakota Group; Kansas, U.S.A.
- *amorphus*, Lesquereux, 1868, p. 102; & 1874, p. 113, pl. xxii,
figs. 3-4. Dakota Group; Nebraska, U.S.A.

- Phyllites angustus*, Crié, 1877, p. 232 [nomen nudum].
Cretaceous; Mans, France.
- (*Comptonia*?) *antiqua*, Nilsson, 1832, p. 346, pl. i, fig. 8.
Greensand; Sweden.
- *aristolochiæformis*, Lesquereux, 1892, p. 217, pl. lix, fig. 8.
Dakota Group; Kansas, U.S.A.
- *betulæfolius*, Lesquereux, 1869, p. 430, pl. xxiii, fig. 4.
Dakota Group; Nebraska, U.S.A.
- *bipartitus*, Velenovsky, 1887, p. 73, pl. xxix, fig. 4.
Cenomanian; Vyserovic, Bohemia.
- *celastroides*, Heer, 1871 A, p. 14, pl. iii, fig. 12.
Senonian; Quedlinburg, Saxony.
- *celatus*, Lesquereux, 1892, p. 215, pl. lxi, fig. 1.
Dakota Group; Kansas, U.S.A.
- *cenomanensis*, Crié, 1877, p. 232 [nomen nudum].
Cenomanian; Ste. Croix, France.
- *cliffwoodensis*, Berry, 1905 A, p. 47, pl. ii, fig. 8.
Cretaceous; Cliffwood, U.S.A.
- *coriaceus*, Newberry in Ives, 1861, p. 132, pl. iii, figs. 7, 7 a.
Lower Cretaceous (?); Arizona, U.S.A.
- *cotinus*, Lesquereux, 1876 B, p. 364.
Dakota Group; Nebraska, U.S.A.
- (*Acer*?) *cretaceum*, Nilsson, 1832, p. 345, pl. i, figs. 1, 2.
Greensand; Sweden.
- *curvinervis*, Hosius, 1870 A, p. 101, pl. xvi, fig. 30. (= *Viburnum subrepandum*, Hosius & v. d. Mark, 1880.)
Senonian; Legden, Westphalia.
- *denticulatus*, Knowlton in Stanton & Hatcher, 1905, p. 148, pl. xix, fig. 2.
Judith River Beds; Montana, U.S.A.
- *durescens*, Lesquereux, 1892, p. 218, pl. lxi, fig. 5; pl. lxii, figs. 3, 4.
Dakota Group; Kansas, U.S.A.
- *Ehrlichi*, Unger, 1850 A, p. 503; & 1867, p. 651, pl. ii, figs. 9, 10.
Upper Cretaceous; Austria.
- *ellipticus*, Newberry, 1895, p. 130, pl. xxiv, fig. 9. (Re-named by Berry, 1909, *Diospyros amboyensis*.)
Amboy Clay; Woodbridge, U.S.A.
- *emarginatus*, Goeppert, 1847, p. 361, pl. xxxvii, fig. 11.
Quadersandstein; Silesia.
- *enervis*, Goeppert, 1847, p. 361, pl. xxxvii, figs. 8, 9 a. Ibid.
- *erosus*, Lesquereux, 1892, p. 216, pl. lxi, fig. 4.
Dakota Group; Kansas, U.S.A.
- (*Alnus*) *Friesii*, Nilsson, 1832, p. 346, pl. i, fig. 7.
Greensand; Sweden.
- *Geinitzensis*, Goeppert, 1865 c, p. 12. (= *Debeya serrata*, Miquel.)
- *Geinitzianus*, Goeppert, 1847, p. 361, pl. xxxvii, figs. 5-7.
Quadersandstein; Silesia.
- *grönlandicus*, Engelhardt in Vanhöffen, 1897, pp. 363, 372, text-fig. 30.
Kome Beds; Greenland.

- Phyllites ilicifolius*, Lesquereux, 1892, p. 213, pl. x, fig. 9.
Dakota Group; Kansas, U.S.A.
- *incurvatus*, Heer, 1882, p. 105, pl. xxvii, fig. 13.
Atane Beds; Greenland.
- *indeterminatus*, Sternberg, 1825, p. 40, pl. xliii, fig. 1.
Greensand; Sweden.
- *inflexinervis*, Saporta, 1894, p. 204, pl. xxxvi, fig. 12 b.
Upper Albian; Portugal.
- *innectens*, Lesquereux, 1892, p. 219, pl. lxx, fig. 6.
Dakota Group; Kansas, U.S.A.
- *intricata*, Knowlton in Stanton & Hatcher, 1905, p. 148, pl. xviii, fig. 6.
Judith River Beds; Montana, U.S.A.
- *lacerus*, Saporta, 1894, pl. xxxv, fig. 11.
Albian (?); Portugal.
- *Lacoei*, Lesquereux, 1892, p. 213, pl. xlv, fig. 6.
Dakota Group; Kansas, U.S.A.
- *lævigatus*, Heer (*non* Miquel), 1874 A, p. 120, pl. xxxiv, fig. 16.
Atane Beds; Greenland.
- *lævigatus*, Miquel (*non* Heer), 1853, p. 41, pl. i, fig. 2.
Senonian; Limburg.
- *latifolius*, Berry, 1910 c, p. 194. (= *Nageiopsis latifolia*, Fontaine, 1889.)
Patuxent Formation; Virginia, U.S.A.
- *laurencianus*, Lesquereux, 1892, p. 215, pl. xlv, fig. 5.
Dakota Group; Kansas, U.S.A.
- *laurinoides*, Hosius, 1870 A, p. 101, pl. xvi, fig. 31. (= *Litsæa laurinoides*, Hosius & v. d. Marck, 1880.)
Senonian; Legden, Westphalia.
- *linguæformis*, Heer, 1874 A, p. 120, pl. xxxiv, fig. 15.
Atane Beds; Greenland.
- *longepetiolatus*, Heer, 1882, p. 105, pl. xv, fig. 8.
Ibid.
- *monocotyleus*, Miquel, 1853, p. 53, pl. i, fig. 3.
Cretaceous; Belgium.
- *multinervis*, Hosius, 1870 A, p. 101, pl. xvii, fig. 34.
Senonian; Legden, Westphalia.
- *nervulosus*, Sternberg, 1825, p. 39, pl. xlii, fig. 2.
Greensand; Hör, Sweden.
- *obcordatus*, Heer in Meek & Hayden, 1859, p. 266. (= *Liriodendron Meekii obcordata*, Heer, 1882.)
Dakota Group; Nebraska, U.S.A.
- *obscura* [us], Hollick in Newberry, 1895, p. 131, pl. xlii, fig. 33.
Amboy Clay; New Jersey, U.S.A.
- *obtusilobatus*, Heer in Meek & Hayden, 1859, p. 266. (= *Liquidambar obtusilobatus*, Hollick in Newberry, 1898.)
Dakota Group; Nebraska, U.S.A.
- *orbicularis*, Newberry, 1895, p. 130, pl. xxiv, figs. 7, 8. (= *Populus orbicularis*, Berry, 1909.)
Amboy Clay; Sayreville, U.S.A.
- *pachyphyllus*, Fontaine, 1889, p. 325, pl. cxlix, fig. 2.
Potomac Formation; Virginia, U.S.A.

- Phyllites pelagicus*, Unger, 1850, p. 503; & 1867, p. 653, pl. ii, fig. 13.
Upper Cretaceous; Austria.
- *perplexus*, Lesquereux, 1892, p. 215, pl. xxxviii, fig. 15.
Dakota Group; Kansas, U.S.A.
- *platanoides*, Bozzi, 1891, p. 379, pl. xxi, fig. 8.
Senonian (?); Vernasso, Italy.
- *poinsettioides*, Hollick, 1893, p. 37, pl. i, fig. 10.
Raritan Formation; New York, U.S.A.
- *problematicus*, Saporta, 1894, p. 97, pl. xvi, fig. 10.
Valanginian; Portugal.
- *proteaceus*, Bozzi, 1891, p. 379, pl. xvi, figs. 6-7.
Senonian (?); Vernasso, Italy.
- *proteoides*, Unger, 1867, p. 652, pl. ii, fig. 11.
Upper Cretaceous; Austria.
- *quinquenervis*, Hosius, 1870, p. 101, pl. xvii, fig. 33.
Senonian; Legden, Westphalia.
- *ramosinervis*, Heer, 1871 A, p. 15, pl. iii, figs. 13, 14.
Senonian; Quedlinburg, Saxony.
- *repandus*, Sternberg (*non* Engelhardt), 1822, p. 29, pl. xxv, fig. 1.
Quadersandstein; Germany.
- *Reussi*, Unger, 1867, p. 653, pl. ii, fig. 12.
Upper Cretaceous; Austria.
- *rhoifolius*, Lesquereux, 1868, p. 101; & 1874, p. 111, pl. xxii, figs. 5, 6.
Dakota Group; Nebraska, U.S.A.
- *rhomboideus*, Lesquereux, 1874, p. 112, pl. vi, fig. 8. (= *Ficus rhomboidea*, Lesquereux, 1868.)
Ibid.
- *sinuatus*, Lange, 1890, p. 671, pl. xxxiv, fig. 9.
Senonian; Spitalgarten, Prussia.
- *Snowii*, Lesquereux, 1892, p. 214, pl. xxxviii, fig. 2.
Dakota Group; Kansas, U.S.A.
- *stipulæformis*, Lesquereux, 1892, p. 216, pl. lxi, fig. 2.
Ibid.
- *Sturi*, Unger, 1865, p. 378, pl. i, figs. 10-11.
Cretaceous (?); Deva, Transylvania.
- *tesselatus*, Sternberg, 1825, pl. xlii, fig. 4 (described as from Cretaceous; removed from Cretaceous by Debey & Ettingshausen, 1859 B, p. 245).
- *testaceus*, Goepfert, 1847, p. 361 (naming figs. 9 & 10, pl. liii of Goepfert, 1842 A).
Quadersandstein; Silesia.
- *Thierensi*, Bosquet MS. in Debey, 1851, p. 569 [nomen nudum]. (= *Delessertites Thierensi*, Miquel, 1853.)
Danian; Maestricht.
- *triloba*, Knowlton, 1900 A, p. 74, pl. xviii, fig. 2.
Montana Formation; Wyoming, U.S.A.
- *triplinervis*, Hosius (*non* Saporta), 1870 A, p. 101, pl. xvi, fig. 32.
Senonian; Legden, Westphalia.
- *triplinervis*, Saporta (*non* Hosius), 1894, p. 209, pl. xxxvi, fig. 7 a.
Upper Albian; Portugal.
- *umbonatus*, Lesquereux, 1868, p. 102; & 1874, p. 113, pl. xix, fig. 4.
Dakota Group; Nebraska, U.S.A.

- Phyllites undulatus**, Newberry, 1895, p. 131, pl. xxiv, fig. 10.
Amboy Clay; Woodbridge, U.S.A.
- **Vanonæ**, Heer in Capellini & Heer, 1867, p. 22, pl. i, fig. 8.
Dakota Group; Nebraska, U.S.A.
- **venosissimus**, Newberry in Ives, 1861, p. 131, pl. iii, fig. 6.
Base of Cretaceous (?); Arizona, U.S.A.
- (**Salix?**) **Wahlbergii**, Nilsson, 1832, p. 345, pl. i, figs. 3-6.
Greensand; Sweden.
- **Winkleri**, Debey, 1848, p. 124 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **zamiaeformis**, Lesquereux, 1892, p. 38, pl. ii, fig. 7.
Dakota Group; Kansas, U.S.A.
- **sp.**, Corda in Reuss, 1846, p. 140, pl. l, figs. 4-10; pl. li, figs. 1-8.
Lower Quader; Bohemia.
- **sp.**, Dawson, 1887, p. 32, pl. ii, fig. 20.
Laramie Formation; Canada.
- **sp.**, Knowlton, 1899 b, pl. lxxviii, fig. 7 (not in text).
Yellowstone Park, U.S.A.
- **sp.**, Knowlton, 1900 A, p. 75, pl. ix, fig. 4.
Montana Formation; Wyoming, U.S.A.
- **sp.**, Knowlton, 1900 A, p. 75, pl. xix, fig. 5. Ibid.
- **sp.**, Lange, 1890, p. 672, pl. xxxiv, fig. 10.
Senonian; Aix, Rhenish Prussia.
- **sp.**, Lesquereux, 1892, p. 216, pl. lix, fig. 7.
Dakota Group; Kansas, U.S.A.
- Phyllocladites crenatus**, Schulze, 1888, p. 21 [nomen nudum].
Senonian; Quedlinburg, Saxony.
- **rotundifolius**, Heer, 1874 A, p. 124, pl. xxxv, figs. 17-21.
Cretaceous; Spitzbergen.
- Phyllocladopsis heterophylla**, Fontaine, 1889, p. 204, pl. lxxxiv, fig. 5; pl. clxvii, fig. 4. Potomac Formation; Virginia, U.S.A.
- Phyllocladus laciniosa**, Schulze, 1888, p. 26 [nomen nudum].
Upper Cretaceous; Heimburg, Germany.
- **subintegrifolius**, Lesquereux, 1868, p. 92; & 1874, p. 54, pl. i, fig. 12. (Re-named *Thinnfeldia subintegrifolia* (Lx.), Knowlton, 1898.)
Dakota Group; Nebraska, U.S.A.
- Phyllocladoxylon antarcticum**, Gothan, 1908 A, p. 4, pl. i, figs. 4-8.
Cretaceous (?); Antarctica.
- Phyllotæmia costulata**, Saporta, 1894, p. 221, pl. xxxix, fig. 20.
Cenomanian (?); Portugal.
- **demersa**, Saporta, 1894, p. 216, pl. xxxviii, fig. 6. Ibid.
- **elongata**, Saporta, 1894, p. 216, pl. xxxix, fig. 21. Ibid.
- **nervosa**, Saporta, 1894, p. 216, pl. xxxviii, figs. 7-8. Ibid.
- **stipulacea**, Saporta, 1894, p. 217, pl. xxxviii, figs. 12-13. Ibid.
- Phymatoderma Lemerianum**, Brongniart, 1849 A, p. 10.
Gault; France.
- Picea albertensis**, Penhallow, 1908, p. 82.
Judith River Series; Alberta, Canada.

- Picea cliffwoodensis*, Berry, 1906 b, p. 169 ; & 1906 b, p. 143, pl. xix.
Magothy Formation ; New Jersey, U.S.A.
- *cretacea*, Velenovsky, 1889, p. 14, pl. i, figs. 4, 5.
Cenomanian ; Vyserovic, Bohemia.
- Piceites exogyrus*, Goeppert. See Hildebrand, 1861, p. 251.
Middle Cretaceous ; Bohemia.
- Pinites aquisgranensis*, Goeppert, 1842 b, p. 151, pl. liv, figs. 1-5,
10-12. (= *Cycadopsis aquisgranensis*, Debey, 1848 b.)
Senonian ; Aix, Rhenish Prussia.
- *Andræi* (Coemans), Gardner, 1886 b, p. 3, pl. x, fig. 1. (= *Pinus*
Andræi, Coemans, 1867.) Gault ; Folkestone.
- *Benstedii*, Endlicher, 1847 A, p. 19. (= *Abies Benstedii*, Mantell,
1846.) Lower Greensand ; Maidstone.
- *cretacea*, Corda in Reuss, 1846, p. 91, pl. xlvii, figs. 1-6. (= *Peuce*
cretacea, Endlicher, 1847 b.) Pläner ; Bohemia.
- *cretaceus*, Dunker (*non* Corda), Goeppert, 1856, p. 182, pl. xxxv,
fig. 2. Quadersandstein ; Blankenburg, Saxony.
- *cyclopterus*, Saporta, 1894, p. 178, pl. xxviii, fig. 20.
Lower Albian ; Portugal.
- *cylindroides*, Gardner, 1886 b, p. 245, & plate, figs. 2, 2 a.
Lower Greensand ; Potton.
- *exogyra* (Corda), Geinitz, 1850 A, p. 272. (= *Pinus exogyra*, Corda
in Reuss, 1846.) Quadersandstein ; Bohemia.
- *gracilis*, Carruthers, 1869, p. 2, pl. i, fig. 9. Gault ; Folkestone.
- *hexagonus*, Carruthers, 1871, p. 540, pl. xv. Ibid.
- *Leckenbyi*, Carruthers, 1869, p. 2, pl. i, figs. 1-5.
Lower Greensand ; Isle of Wight.
- *Leei*, Fontaine in Ward, 1905, p. 570, pl. cxix, figs. 6, 7.
Older Potomac Formation ; Maryland, U.S.A.
- *lepidodendroides*, Roemer, 1870, p. 290, pl. xxvii, fig. 7 ;
pl. xxviii, fig. 3. Perucer Beds ; Silesia.
- *Mantellii*, Carruthers, 1866 b, p. 543, pl. xxi, fig. 3.
Greensand ; Maidstone.
- *oblongus*, Williamson, 1887, p. 193, pl. ix. (= *Abies oblonga*,
Lindley & Hutton, 1834.) Greensand ; Dorset.
- *patens*, Miquel (*non* Carruthers), 1853, p. 41, pl. ii.
Senonian ; Limburg.
- *patens*, Carruthers (*non* Miquel), 1866 b, p. 543, pl. xxi, fig. 4.
Greensand ; Maidstone.
- *pottoniensis*, Gardner, 1886 b, p. 245, fig. 3.
Lower Greensand ; Potton.
- *Pricei*, Carruthers, 1875, p. 280, text-fig. Gault ; Folkestone.
- ? *Quenstedtii*, Feistmantel, 1874, p. 272. Perucer Beds ; Bohemia.
- *Reussii*, Endlicher, 1847 b, p. 287. (= *Pinus Reussii*, Corda, 1846.)
Cretaceous ; Bohemia.
- *sussexiensis*, Carruthers, 1866 b, p. 541. (= *Zamia sussexiensis*,
Mantell.) Lower Greensand ; Sussex.
- *undulatus*, Eichwald, 1853, p. 231. Lower Cretaceous ; Russia.

- Pinites* sp., Roemer, 1870, p. 301, pl. xxx, fig. 2.
Turonian; Upper Silesia.
- Pinostrobus prolongatus*, Feistmantel, 1874, p. 272.
Perucer Beds; Bohemia.
- *validus*, Feistmantel, 1874, p. 272. Ibid.
- Pinus Andræi*, Coemans, 1867, p. 12, pl. iv, fig. 4; pl. v, fig. 1.
(=*Pinites Andræi* (Coemans), Gardner, 1886 B.)
Cretaceous; La Louvière, Belgium.
- *anthraciticus*, Dawson, 1893, p. 89, text-fig. 10.
Kootanie Formation; North-West Territory, Canada.
- *Argonnensis*, Fliche, 1896, p. 217, pl. ix, fig. 3. [Cones.]
Albian; Clermont, France.
- *aspera*, Cornuel, 1866, p. 671, pl. xii, figs. 6-12.
Neocomian; France.
- *Briarti*, Coemans, 1867, p. 10, pl. iv, fig. 2.
Cretaceous; La Louvière, Belgium.
- (*Cedrus*) *Corneti*?, Coemans, 1867, p. 11, pl. iv, fig. 3. Ibid.
- *Crameri*, Heer, 1868, p. 84, pl. xlv, figs. 7-18.
Kome Beds; Greenland.
- *delicatulus*, Berry, 1904 A, p. 68, pl. i, fig. 12.
Matawan Formation; New Jersey, U.S.A.
- *depressa*, Coemans, 1867, p. 15, pl. v, fig. 4.
Cretaceous; La Louvière, Belgium.
- *Eirikiana*, Heer, 1874 A, p. 85, pl. ii, fig. 1; pl. xvii, figs. 6, 7;
pl. xviii, fig. 2 b; pl. xxiii, fig. 16. Kome Beds; Greenland.
- *elongata*, d'Orbigny, 1852, p. 617; & Cornuel, 1882, p. 260, pl. vii,
fig. 1. Aptian; Saint-Dizier.
- *exogyra*, Corda in Reuss, 1846, p. 91, pl. xlviii, figs. 16-18.
(=*Pinites exogyra*, Geinitz, 1850 A.) Greensand; Bohemia.
- *familiaris*, Brongniart, 1828 A, p. 107. (= *Conites familiaris*,
Sternberg, 1825.)
- *gibbosa*, Coemans, 1867, p. 13, pl. v, fig. 2.
Cretaceous; La Louvière, Belgium.
- *gracilis*, Cornuel, 1866, p. 671, pl. xii, figs. 4, 5.
Neocomian; France.
- *Guill(ie)ri*, Crieé, 1884, p. 512. Cretaceous; France.
- *Heeri*, Coemans, 1867, p. 14, pl. v, fig. 3.
Cretaceous; La Louvière, Belgium.
- *lingulata*, Heer, 1874 A, p. 84, pl. xii, fig. 10 d; pl. xxxiii, fig. 18.
Kome Beds; Greenland.
- *longissima*, Velenovsky, 1885, p. 29, pl. i, figs. 14-17.
Perucer Beds; Bohemia.
- *macrostrobilina*, Menzel, 1908, p. 27, pl. ii, figs. 1, 1 a, 1 b.
Quadersandstein; Bohemia.
- *mammilifer*, Saporta, 1880, p. 654, pl. iv, fig. 2.
Aptian; Cauville, France.
- *mattewanensis*, Berry, 1903 c, p. 680, text-fig. 4 on p. 678.
Matawan Formation; New Jersey, U.S.A.

- Pinus monasteriensis*, Hosius & von der Marck, 1880, p. 141, pl. xxvi, fig. 19. Upper Senonian; Westphalia.
- *Nathorsti*, Conwentz, 1892, p. 13, pl. i, fig. 1; pl. ii, figs. 1-5; pl. iii, figs. 1-3; pl. vi, figs. 1-10; pl. vii, figs. 1-4. Senonian; Sweden.
- *Nordenskiöldi*, Heer, 1876 c, p. 45, pl. ix, figs. 1-6. (Recorded Canadian Kootanie, Dawson, 1893.) Jurassic; Spitzbergen.
- *Olafiana*, Heer, 1874 A, p. 85, pl. xx, fig. 10 b; pl. xxiii, figs. 19, 19 b. Kome Beds; Greenland.
- *Omalii*, Coemans, 1867, p. 10, pl. iv, fig. 1. (= *Abies Omalii*, Schimper, 1872.) Cretaceous; La Louvière, Belgium.
- *oxyptera*, Saporta, 1890, p. 4, pl. xiii, fig. 7. Danian (?); France.
- *Parsyi*, Saporta, 1880, p. 655, pl. iv, fig. 3. Albian; Bléville, France.
- *Peterseni*, Heer, 1868, p. 84, pl. xlv, fig. 19. Kome Beds; Greenland.
- *Præhalepensis*, Fliche, 1896, p. 225, pl. x, fig. 2. Albian; Thibaudette, France.
- *Præmonticola*, Fliche, 1896, p. 220, pl. ix, fig. 4. Albian; Clermont, France.
- *protopicea*, Velenovsky, 1885, p. 31, pl. vii, figs. 1, 2, 4, 6; pl. iii, fig. 4. Cenomanian; Vyserovic, Bohemia.
- *Quenstedti*, Heer, 1869 A, p. 13, pl. ii, figs. 5-9; pl. iii. Cenomanian; Moletain, Moravia.
- *quinquefolia*, Hollick & Jeffrey, 1909, p. 16, pl. xxii, fig. 2. Raritan Formation; Staten Island, U.S.A.
- *raritanensis*, Berry, 1909, p. 247 (naming *Pinus* sp., Newberry). Raritan Formation; New Jersey, U.S.A.
- *Reussii*, Corda in Reuss, 1846, p. 90, pl. xlv, figs. 22-25. (= *Pinites Reussi*, Endlicher.) Greensand; Bohemia.
- *rhombifera*, Cornuel, 1866, p. 670, pl. xii, fig. 19. Neocomian; France.
- *Saportana*, Fliche, 1896, p. 232, pl. x, figs. 5, 6, text-fig. 9. Albian; Islettes, France.
- *schista*, Ward, 1905, p. 531, pl. cxii, figs. 13-15. Older Potomac Formation; Maryland, U.S.A.
- *shastensis*, Ward, 1905, p. 262, pl. lxix, figs. 1-3. Shasta Formation; California, U.S.A.
- *Staratschini*, Heer, 1874 A, pp. 104, 129, pl. xxviii, fig. 15; pl. xxxiv, fig. 1 c; pl. xxxviii, figs. 6-7. Atane Beds; Greenland.
- *submarginata*, Cornuel, 1866, p. 669, pl. xii, figs. 13-18 & 20, 21. Neocomian; France.
- *sulcata*, Velenovsky (*non* Eudes-Deslongchamps), 1885, p. 30, pl. vii, figs. 5, 11. Pläner; Bohemia.
- *susquaensis*, Dawson, 1883, p. 23, pl. iii, fig. 36. Cretaceous; North-West Territory, Canada.
- *tetraphylla*, Jeffrey, 1908, p. 214, pl. xiv, fig. 17. Raritan Formation; Staten Island, U.S.A.

- Pinus* (*Ptero-Cembra*) *Toillezi*, Coemans, 1867, p. 16, pl. v, fig. 5.
Cretaceous; La Louvière, Belgium.
- *triphylla*, Hollick & Jeffrey, 1909, p. 14, pl. iii, figs. 6, 7 (?);
pl. xxii, fig. 1. Raritan Formation; Staten Island, U.S.A.
- *ucranicus*, Goeppert, 1865 c, p. 12 [nomen nudum].
Danian; Maestricht.
- *umbonatus*, Lesquereux, 1868, p. 102.
Cretaceous; Nebraska, U.S.A.
- (*Abies*) *Upervikensis*, Heer, 1882, p. 56, pl. ix, figs. 5-7.
Atane Beds; Greenland.
- *vaginalis*, Heer, 1874 A, p. 103, pl. xxvii, fig. 15 b. Ibid.
- *vernonensis*, Ward, 1905, p. 497, pl. cix, figs. 4-6.
Older Potomac Formation; Virginia, U.S.A.
- *Wohlgemuthi*, Fliche, 1896, p. 223, pl. x, fig. 1.
Albian; Clermont, France.
- *yezoensis*, Stopes & Kershaw, 1910, p. 399, pl. xxvii, figs. 3-5;
pl. xxviii, fig. 2. Upper Cretaceous; Hokkaido, Japan.
- *sp.*, Cornuel, 1866, p. 673, pl. xii, figs. 2-3. Neocomian; France.
- *sp.*, Engelhardt, 1892, p. 105, pl. ii, fig. 16.
Cenomanian; Niederschoena, Saxony.
- ? *sp.*, Fontaine, 1889, p. 272, pl. clxx, fig. 4.
Lower Cretaceous; Maryland, U.S.A.
- *sp.*, Heer, 1874 A, p. 129, pl. xxxvii, fig. 5. Cretaceous; Spitzbergen.
- *sp.*, Hollick, 1893, p. 31, pl. i, figs. 13, 19, 20, 22.
Middle Cretaceous; Staten Island, U.S.A.
- *sp.*, Hollick & Jeffrey, 1909, p. 16, pl. ix, figs. 11, 12; pl. xxiii,
fig. 6. Ibid.
- *sp.*?, Hollick & Jeffrey, 1909, p. 17, pl. iii, fig. 8; pl. xxii, fig. 5.
Ibid.
- *sp.*?, Newberry, 1895, p. 47, pl. ix, figs. 5-8, 17, 18.
Amboy Clay; New Jersey, U.S.A.
- Pisonia atavia*, Velenovsky, 1887, p. 67, pl. xxxi, figs. 13, 14.
Cenomanian; Bohemia.
- *cretacea*, Berry, 1910 A, p. 191.
Middle Cretaceous; North Carolina, U.S.A.
- Pistacia Aquehongensis*, Hollick, 1898 B, p. 421, pl. xxxvi, fig. 5.
Cretaceous; Staten Island, U.S.A.
- *oblanceolata* (Lesquereux), Knowlton, 1898, p. 167 (re-naming
Ficus oblanceolata, Lesquereux, 1873).
Laramie Formation; Wyoming, U.S.A.
- Pistia corrugata*, Lesquereux, 1876 c, p. 299; & 1878 B, p. 103, pl. lxi,
figs. 1, 3-6, 11. Montana Formation; Wyoming, U.S.A.
- *Mazeli*, Saporta & Marion, 1885, p. 37, text-fig. 114, C. D.
Senonian; France.
- *Nordenskiöldi* (Heer), Berry, 1910 A, p. 189 (re-naming *Chondro-*
phyllum Nordenskiöldi, Heer). Atane Beds; Greenland.
- Pistites loriformis*, Hosius & von der Marck, 1880, p. 182, pl. xxxviii,
figs. 151, 152. Lower Senonian; Westphalia.

- Pitcarnia primæva*, Hosius & von der Marek, 1880, p. 217, pl. xlv, figs. 210, 211. Neocomian; Westphalia.
- Pityoidolepis statenensis*, Hollick & Jeffrey, 1909, p. 53, pl. ix, figs. 13, 14; pl. xxvii, figs. 1-3. Raritan Formation; Staten Island, U.S.A.
- Pityoxylon Argonnense*, Fliche, 1896, p. 253. Albian; France.
- *Hollicki*, Knowlton in Hollick, 1898, p. 134, text-figs. 1, 2. Upper Cretaceous; New Jersey, U.S.A.
- *infracretaceum*, Fliche, 1896, p. 251, pl. xii, fig. 1; pl. xvi, figs. 1, 2. Albian; Clermont, France.
- *piceoides (cretaceum)*, Vater, 1884, p. 821. Lower Senonian (?); Brunswick.
- *scituatense*, Jeffrey & Chrysler, 1906, p. 11, pl. ii, figs. 8-12. Cretaceous (?); Massachusetts, U.S.A.
- *statenense*, Jeffrey & Chrysler, 1906, p. 8, pl. i; pl. ii, fig. 7. Middle Cretaceous; Staten Island, U.S.A.
- *Thomasi*, Fliche, 1896, p. 254, pl. xvi, fig. 3. Albian; Triancourt, France.
- Pitys familiaris*, Unger, 1845, p. 197. (= *Conites familiaris*, Sternberg, 1825.)
- Planera antiqua*, Heer, 1883, p. 26, pl. iv, figs. 11-12. Patoot Beds; Greenland.
- *antiqua*, Newberry, 1895, p. 69, pl. xlii, figs. 1-4. (Name pre-occupied, = *P. Knowltoniana*, Hollick.) Amboy Clay; New Jersey, U.S.A.
- *betuloides*, Hollick, 1906, p. 57, pl. viii, fig. 22. Middle Cretaceous; Martha's Vineyard, U.S.A.
- *cretacea*, Berry, 1907, p. 193, pl. xi, figs. 7, 8. Middle Cretaceous; North Carolina, U.S.A.
- *Knowltoniana*, Hollick in Newberry, 1895, p. 69, pl. xlii, figs. 1-4. Amboy Clay; New Jersey, U.S.A.
- Plantaginopsis marylandica*, Fontaine in Ward, 1905, p. 561, pl. cxvii, fig. 7; pl. cxviii, figs. 1, 2. Older Potomac Formation; Maryland, U.S.A.
- Platæanthus (Williamsonia) problematicus*, Newberry, 1895, p. 125, pl. xxxv, figs. 1-9. Amboy Clay; New Jersey, U.S.A.
- Plataninium subaffine*, Vater, 1884, p. 843, pl. xxix, figs. 19-21. Lower Senonian (?); Brunswick.
- Platanophyllum crassinerve*, Fontaine, 1889, p. 316, pl. clviii, fig. 5. Potomac Formation; Virginia, U.S.A.
- Platanus aceroides? latior* (Goepfert), Lesquereux, 1868, p. 97. (= *P. latior*, Knowlton, 1898.) Dakota Group; Nebraska, U.S.A.
- *acute-triloba*, Krasser, 1889, p. 34; & 1896, p. 142, pl. xiii, fig. 2. Cenomanian; Kunstadt, Moravia.
- *affinis*, Lesquereux, 1874, p. 71, pl. iv, fig. 4. (= *Populites affinis*, Lx., 1873, & *Cissites affinis*, Lx., 1876.) Dakota Group; Kansas, U.S.A.

- Platanus affinis ampla*, Dawson, 1836, p. 12. (Re-named *Cissites affinis ampla* (Dn.), Knowlton.) Mill Creek Series; Mill Creek, Canada.
- *Aquehongensis*, Hollick, 1893, p. 32, pl. iv.
Raritan Formation (?); Staten Island, U.S.A.
- *araliæformis*, Krasser, 1893, p. 144, pl. xii, fig. 6.
Cenomanian; Kunstadt, Moravia.
- *betulæfolia*, Krasser, 1893, p. 34; & 1896, p. 143, pl. xiv, fig. 4.
Ibid.
- *cissoides*, Lesquereux, 1892, p. 75, pl. lxi, fig. 3.
Dakota Group; Kansas, U.S.A.
- *cuneiformis*, Krasser, 1896, p. 141, pl. xii, fig. 5; pl. xiv, fig. 3.
Cenomanian; Kunstadt, Moravia.
- *diminutiva*, Lesquereux, 1868, p. 98 (-us); & 1874, p. 73, pl. viii, fig. 5.
Dakota Group; Nebraska (?), U.S.A.
- *Heerii*, Lesquereux, 1872, p. 303; & 1874, p. 70, pl. viii, fig. 4; pl. ix, figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- *irregularis*, Krasser, 1896, p. 143, pl. xvi, fig. 2.
Cenomanian; Kunstadt, Moravia.
- *Kuemmelii*, Berry, 1906 B, p. 176; & 1906 E, p. 146, pl. xxiii, figs. 2, 3; pl. xxiv.
Magothy Formation; New Jersey, U.S.A.
- *lævis*, Velenovsky, 1889, p. 16, pl. i, figs. 1, 2. (= *Credneria lævis*, Vel., 1882 A.)
Cenomanian; Bohemia.
- *latiloba*, Newberry, 1870, p. 23; 1878, pl. ii, fig. 4; & 1898, p. 105, pl. i, fig. 4.
Dakota Group; Nebraska, U.S.A.
- *laticus* (Lesquereux), Knowlton, 1898, p. 170 (re-naming *Platanus primæva*, Lx., 1874, & *P. aceroides* (?), Goëpp., var. *laticus*, Lx., 1868.)
Dakota Group; Kansas, U.S.A.
- *moravica*, Krasser, 1896, p. 140, pl. xiii, fig. 3; pl. xv, fig. 3.
Cenomanian; Kunstadt, Moravia.
- ? *Newberryana*, Heer in Capellini & Heer, 1867, p. 16, pl. i, fig. 4.
Dakota Group; Nebraska, U.S.A.
- *obtusiloba*, Lesquereux, 1868, p. 97; & 1874, p. 69, pl. vii, figs. 3, 4.
Ibid.
- *onomastus*, Bayer, 1896, p. 22, text-figs. 17, 18, & p. 35.
Cenomanian; Kieslingswalda, Bohemia.
- *platanoides* (Lesquereux), Knowlton, 1898, p. 171 (re-naming *Viburnum platanoides*, Lesquereux, 1878 B).
Laramie Formation; Wyoming, U.S.A.
- *primæva*, Lesquereux, 1868, p. 97; & 1874, p. 69, pl. vii, fig. 2; pl. xxvi, fig. 2. (Re-named *P. laticus* by Knowlton, 1898.)
Dakota Group; Kansas, U.S.A.
- *primæva grandidentata*, Lesquereux, 1892, p. 73, pl. ix, figs. 1, 2. (Re-named *P. laticus grandidentata* by Knowlton, 1898.) Ibid.
- *primæva integrifolia*, Lesquereux, 1892, p. 74, pl. xlix, fig. 4. (Re-named *P. laticus integrifolia* by Knowlton, 1898.) Ibid.
- *primæva subintegrifolia*, Lesquereux, 1892, p. 73, pl. ix, figs. 3, 4. (Re-named *P. laticus subintegrifolia* by Knowlton, 1898.) Ibid.

- Platanus pseudo-Guillelmæ**, Krasser, 1896, p. 139, pl. xiv, figs. 1, 2.
Cenomanian; Kunstadt, Moravia.
- **Raynoldsii integrifolia**, Lesquereux, 1878 n, p. 185, pl. xxvi, figs. 4, 5.
Laramie Formation; Wyoming, U.S.A.
- **recurvata**, Lesquereux, 1874, p. 71, pl. x, figs. 3-5. (= *Sassafras* (*Araliopsis*) *recurvatum*, Lesquereux, 1873.)
Dakota Group; Kansas, U.S.A.
- **rhomboidea**, Velenovsky, 1889, p. 17, pl. ii, fig. 10; pl. vi, figs. 2, 3. (= *Credneria rhomboidea*, Vel., 1882.)
Cenomanian; Bohemia.
- **Velenovskyana**, Krasser, 1896, p. 138, pl. xv, fig. 2.
Cenomanian; Kunstadt, Moravia.
- **vyserovicensis**, Marik, 1901, p. 10, pl. ii, figs. 4, 5.
Cenomanian; Bohemia.
- ? **Wardii**, Knowlton, 1900 a, p. 14, pl. ii, figs. 1-4.
Montana Formation; Missouri R., U.S.A.
- **sp.**, Heer, 1883 a, p. 258, pl. liv, fig. 1 d. Patoot Beds; Greenland.
- **sp.**, Hollick, 1906 a, p. 83, pl. xxxi, fig. 5.
Cretaceous; Staten Island, U.S.A.
- Platyceriphyllum cretaceum**, Velenovsky, 1889, p. 29, pl. v, fig. 16.
(= *Platycerium cretaceum*, Vel. = *Hausmannia cretacea* (Velenovsky), Richter, 1906.)
Cenomanian; Vyserovic, Bohemia.
- Platycerium cretaceum**, Velenovsky, 1889, p. 5, pl. v, fig. 16.
(= *Dipteriphyllum cretaceum* (Velenovsky), Krasser.)
Ibid.
- Platypterigium densinerve**, Fontaine, 1889, p. 169, pl. xxx, fig. 8;
pl. xxxi, figs. 1, 4; pl. xxxii, figs. 1, 2; pl. xxxiii, fig. 1; pl. xxxiv, fig. 1;
pl. xxxv, figs. 1, 2. (Re-named *Nilssonia densinerve* by Berry, 1910 n.)
Potomac Formation; Virginia, U.S.A.
- **Rogersianum**, Fontaine, 1889, p. 171, pl. xxxi, fig. 2; pl. xxxiii, fig. 2;
pl. xxxiv, fig. 2. Ibid.
- Pleospores Shirainus**, Suzuki, 1910, p. 191, pl. vii, fig. 6, text-figs. 2 & 3.
Upper Cretaceous; Hokkaido, Japan.
- Plutonia cretacea**, Velenovsky, 1889, p. 11, pl. ii, figs. 11-20; pl. iii, figs. 1, 2.
Cenomanian; Vyserovic, Bohemia.
- Pocites acicularis**, Saporta, 1894, p. 146, pl. xxvi, fig. 7 b.
Urgonian; Portugal.
- **borealis**, Heer, 1874 a, p. 86, pl. xxiv, fig. 5.
Kome Beds; Greenland.
- **cercalinus**, Saporta, 1894, p. 146, pl. xxvii, fig. 17.
Urgonian; Portugal.
- **cretaceus**, Marik, 1901, p. 13, pl. ii, figs. 13-14.
Cenomanian; Bohemia.
- **gemillinervis**, Saporta, 1894, p. 96, pl. xv, fig. 36.
Valanginian; Portugal.
- **lævis**, Saporta, 1894, p. 180, pl. xxxii, fig. 10 (antedated by A. Braun, 1851).
Albian; Portugal.
- **Nelsonicus**, Ettingshausen, 1887 a, p. 180, pl. vii, figs. 22, 22 a.
Upper Cretaceous; New Zealand.

- Poacites paucinervis*, Saporta (*non* Heer, *nee* Watelet), 1894, p. 95, pl. xvi, fig. 33. Valanginian; Portugal.
- *plurinervis*, Saporta, 1894, p. 116, pl. xxii, fig. 8. Aptian; Portugal.
- *plurinervulosus*, Saporta, 1894, p. 146, pl. xxvi, fig. 4. Urgonian; Portugal.
- *striatifolius*, Saporta, 1894, pp. 57, 110, pl. xx, figs. 11, 12. Lower Cretaceous (?); Portugal.
- *tenellus*, Saporta, 1894, p. 95, pl. xvi, fig. 8; pl. xvii, fig. 3. Valanginian; Portugal.
- ? *sp.*, Hollick, 1894 A, p. 64, pl. clxxx, figs. 2, 12. Middle Cretaceous; Martha's Vineyard, U.S.A.
- *sp.*, Hollick, 1905 C, p. 411, pl. lxxiii, fig. 1. Cretaceous Shale; Long Island, U.S.A.
- Podalyriophyllum brochidodromum*, Ettingshausen, 1893, p. 151; & 1895, p. 51, pl. iv, fig. 17. Cretaceous; Australia.
- Podocarpites Tyrrellii*, Dawson, 1888, p. 35. Belly River Series; Canada.
- Podocarpium cupressinum*, Ettingshausen, 1887 A, p. 177, pl. vii, fig. 11. Upper Cretaceous; New Zealand.
- *prædacrydioides*, Ettingshausen, 1887 A, p. 178, pl. vii, fig. 12. Ibid.
- *tenuifolium*, Ettingshausen, 1887 A, p. 177, pl. vii, figs. 8-10, 10 A. Ibid.
- *Ungeri*, Ettingshausen, 1887 A, p. 177, pl. vii, figs. 13-15. Ibid.
- Podocarpoxylon aparenchymatosum*, Gothan, 1908 A, p. 8, pl. i, fig. 11. Cretaceous (?); Antarctica.
- Podocarpus cretacea*, Velenovsky, 1885, p. 13, pl. xii, figs. 5-11. (= *Podocarpus sp.*, Velenovsky, 1888 A.) Perucer Beds; Menlik.
- *sarthacensis*, Crie, 1879 (?), p. 30 [nomen nudum]. Cretaceous; Sainte-Croix, France.
- *sp.*, Velenovsky, 1888 A, p. 596, fig. 2. Cenomanian; Bohemia.
- Podogonium Americanum*, Lesquereux, 1878 B, p. 298, pl. lix, fig. 5; pl. lxiii, fig. 5; pl. lxv, fig. 6. Montana Formation; Wyoming, U.S.A.
- Podozamites acuminatus*, Hollick in Newberry, 1895, p. 45, pl. xiii, fig. 7. Amboy Clay; New Jersey, U.S.A.
- *acutifolius*, Fontaine, 1889, p. 181, pl. lxxx, fig. 6; pl. lxxxv, figs. 10, 15; pl. lxxxvii, fig. 1; pl. clxx, fig. 2. Potomac Formation; Virginia, U.S.A.
- ? *acutus*, Saporta, 1894, p. 87, pl. xvi, fig. 28. Valanginian; Portugal.
- *affinis* (Schenk), Schimper, 1870, p. 161. Urgonian; Silesia.
- *alcantarina*, Saporta, 1894, p. 212, pl. xxxix, fig. 1. Cenomanian; Portugal.
- *angustifolius* (Eichwald), Schimper, 1870, p. 160; recorded Lesquereux, 1884, p. 28. (Re-named *Podozamites Knowltoni* by Berry, 1909.) Dakota Group; Kansas, U.S.A.

- Podozamites austro-caledonica**, Crié, 1889, p. 27 [nomen nudum].
Cretaceous; Portes de Fer, New Caledonia.
- **caudatus**, Lesquereux, 1883, p. 29. (= *Dammarites caudatus*, Lx., 1892.)
Dakota Group; Kansas, U.S.A.
- **distantinervis**, Fontaine, 1889, p. 179, pl. lxxix, fig. 5; pl. lxxxii, fig. 4; pl. lxxxiii, figs. 1, 2, 6, 7; pl. lxxxiv, figs. 1, 2, 8, 10, 14, 15; pl. lxxxv, figs. 1, 2, 16.
Potomac Formation; Virginia, U.S.A.
- **Eichwaldi** (Schimper), Heer, 1877 A, p. 36, pl. vii, fig. 7 e; pl. viii, figs. 1-4; pl. vi, fig. 22 a.
Jurassic; Siberia.
(Recorded Perucer Beds, Bohemia, Velenovsky, 1885.)
- **ellipsoideus**, Saporta, 1894, p. 87, pl. xvi, fig. 31; p. 176, pl. xxxii, figs. 8-9; pl. xxxiii, fig. 5; pl. xxxv, fig. 12.
Valanginian & Albian; Portugal.
- **emarginatus**, Lesquereux, 1883, p. 29.
Dakota Group; Kansas, U.S.A.
- **gracilior**, Saporta, 1894, p. 175, pl. xxxiii, fig. 6.
Albian; Portugal.
- **grandifolius**, Fontaine, 1889, p. 180, pl. lxxxii, fig. 2; pl. lxxxiii, fig. 5.
Potomac Formation; Virginia, U.S.A.
- **Haydenii**, Lesquereux, 1883, p. 27. (= *Pterophyllum Haydenii*, Lesquereux, 1868.)
Dakota Group; Nebraska, U.S.A.
- **Henriquesi**, Saporta, 1894, p. 174, pl. xxxiv, figs. 6-7.
Albian; Portugal.
- **Hoheneggeri**, Schenk, 1871 A, p. 9, pl. ii, figs. 3-6. (= *Glossozamites Hoheneggeri*, Yokoyama, 1906.)
Wernsdorfer Beds; Austrian Silesia.
- **inæquilateralis**, Berry, 1910 c, p. 194. (= *Nageiopsis inæquilateralis*, Fontaine, 1889.)
Patuxent Formation; Virginia, U.S.A.
- **Knowltoni**, Berry, 1909, p. 247 (re-naming *Podozamites angustifolius* (Eichwald), Schimper, 1872, & *Zamites angustifolius*, Eichwald, 1860.)
- **lanceolatus** (Lindley & Hutton), Schimper, 1872, p. 160. Oolitic.
(Recorded, Lesquereux, 1891, Kootanie Formation of America.)
- **latipennis**, Heer, 1882, p. 42, pl. xiv, figs. 1-9; pl. xv, figs. 2 a, 3 b.
Atane Beds; Greenland.
- **linearis**, Saporta, 1894, p. 86, pl. xv, figs. 23-24.
Valanginian; Portugal.
- **lingulatus**, Schimper, 1872, p. 161.
Urgonian; Silesia.
- **longipennis**, Velenovsky, 1885, p. 10, pl. ii, fig. 7.
Perucer Beds; Bohemia.
- **marginatus**, Heer, 1882, p. 43, pl. xvi, fig. 10.
Atane Beds; Greenland.
- **minor**, Heer (*non* (Schenk) Nathorst), 1882, p. 44, pl. xvi, fig. 8.
Ibid.
- **modestior**, Saporta, 1894, p. 174, pl. xxxi, fig. 11; pl. xxxii, fig. 7.
Albian; Portugal.

- Podozamites nervosa**, Newberry (*non* Schenk), 1891, p. 200, pl. xiv, fig. 6 (name antedated by Schimper, 1872, p. 161).
Kootanie Formation; Montana, U.S.A.
- **oblongus**, Lesquereux, 1883, p. 28, pl. i, figs. 10-11.
Dakota Group; Kansas, U.S.A.
- **obovatus**, Schenk, 1871 A, p. 10, pl. ii, figs. 7-10; pl. iii, figs. 1-3.
Urgonian; Austrian Silesia.
- **obtus**, Velenovsky, 1885, p. 9, pl. i, figs. 8, 9.
Perucer Beds; Bohemia.
- **oviformis**, Saporta, 1894, p. 87, pl. xvi, fig. 29.
Valanginian; Portugal.
- **pedicellatus**, Fontaine, 1889, p. 180, pl. lxxvi, fig. 1; pl. lxxviii, fig. 7; pl. lxxxii, fig. 5. Potomac Formation; Virginia, U.S.A.
- **praelongus**, Lesquereux, 1883, p. 29.
Dakota Group; Kansas, U.S.A.
- **proximans**, Conrad, 1869, p. 361, text-fig.
Raritan Clay; New Jersey, U.S.A.
- **pusillus**, Velenovsky, 1885, p. 11, pl. ii, figs. 20, 21, 22, 24 a.
Perucer Beds; Bohemia.
- **Stenopus**, Lesquereux, 1892, p. 27, pl. i, fig. 7.
Dakota Group; Kansas, U.S.A.
- **striatus**, Velenovsky, 1885, p. 10, pl. ii, fig. 8. (= *Dammaphyllum striatum*, Velenovsky, 1889.)
Perucer Beds; Bohemia.
- **subfalcatus**, Fontaine, 1889, p. 179, pl. lxviii, fig. 6; pl. clxx, fig. 9.
Dakota Group; Virginia, U.S.A.
- **tenuinervis**, Heer, 1882, p. 44, pl. xvi, fig. 9.
Atane Beds; Greenland.
- **Zittelii**, Schenk, 1871 A, p. 8, pl. i, fig. 8.
Wernsdorfer Beds; Austrian Silesia.
- **sp.**, Fontaine, 1894, p. 267, pl. xxxvi, fig. 8.
Comanche Formation; Texas, U.S.A.
- ? **sp.**, Hollick, 1894 A, p. 62, pl. clxxx, fig. 4.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- **sp.?**, Hollick, 1902 A, p. 401, pl. xli, figs. 8, 9.
Cretaceous; Massachusetts, U.S.A.
- Polypodites blechnoides**, Debey, 1848 A, p. 116; & 1850, p. 116.
Senonian; Aix, Rhenish Prussia.
- **gracilis**, Marik, 1901, p. 6, pl. i, figs. 14-15.
Cenomanian; Bohemia.
- **Mantelli**, Goeppert, 1836, p. 341. (= *Lonchopteris Mantelli*, Brongniart, 1828.)
- **Schneiderianus**, Goeppert, 1845, p. 214.
Quadersandstein; Löwenberg.
- **zonatus**, Marik, 1901, p. 4, pl. i, figs. 9-12.
Cenomanian; Bohemia.
- Polypodium dentatum**, Fontaine, 1889, p. 105, pl. xxii, figs. 4, 5.
Potomac Formation; Virginia, U.S.A.
- **fadyenioides**, Fontaine, 1889, p. 104, pl. xvi, figs. 4, 5. Ibid.

- Polypodium Graahianum**, Heer, 1888 A, p. 3, pl. xlviii, figs. 4, 4 *b*, 5 *b*, 5 *bb*.
Patoot Beds; Greenland.
- **Hochstetteri**, Unger, 1866, p. 5, pl. ii, figs. 1, 2.
Lower Cretaceous (?); New Zealand.
- Polystichum Hillsianum**, Hollick, 1902, p. 146, pl. iv, fig. 7.
Laramie Formation; Colorado, U.S.A.
- Polytaenia quinquesecta**, Saporta & Marion, 1885, p. 117, text-fig. 125 A.
Turonian; France.
- Populites affinis**, Lesquereux, 1873, p. 423. (= *Platanus affinis*, Lx., 1874, & *Cissites affinis*, Lx., 1876.)
Dakota Group; Kansas, U.S.A.
- **amplus**, Knowlton in Stanton & Hatcher, 1905, p. 140, pl. xviii, fig. 1.
Judith River Beds; Montana, U.S.A.
- **cuneatus**, Lesquereux in Saporta & Marion, 1878, p. 10 [nomen nudum].
Dakota Group; U.S.A.
- **cyclophylla** (Heer), Lesquereux, 1868, p. 93. (= *Populus cyclophylla*, Heer, 1858, & *Cissites cyclophylla*, Lx., 1876.)
Dakota Group; Nebraska, U.S.A.
- **elegans**, Lesquereux, 1868, p. 94; & 1895, p. 10, pl. A, fig. 2; pl. B, fig. 1.
Ibid.
- **fagifolia**, Lesquereux, 1873, p. 422.
Dakota Group; Kansas, U.S.A.
- **flabellata**, Lesquereux, 1868, p. 94. (Re-named *Grewiopsis flabellata* by Knowlton, 1898.)
Dakota Group; Nebraska, U.S.A.
- **lancastriensis**, Lesquereux, 1874, p. 58, pl. iii, fig. 1. (= *Populus lancastriensis*, Lesquereux, 1868.)
Ibid.
- **litigosus** (Heer), Lesquereux, 1892, p. 46, pl. vii, fig. 7; pl. viii, fig. 5; pl. xvi, fig. 6; pl. xlvii, fig. 1. (= *Populus litigiosa*, Heer, 1867.)
Dakota Group; Kansas, U.S.A.
- **microphyllus**, Lesquereux, 1869, p. 430, pl. xxiii, figs. 2, 3.
Dakota Group; Nebraska, U.S.A.
- **ovata**, Lesquereux, 1868, p. 94. (= *Ampelophyllum ovatum*, Lx., 1876.)
Ibid.
- **probalsarnifera**, Dawson, 1894, p. 57, pl. vii, fig. 23.
Upper Cretaceous; Vancouver Island, U.S.A.
- **quadrangularis**, Lesquereux, 1868, p. 94. (= *Hamamelites quadrangularis*, Lesquereux, 1876, and *Ahnites quadrangularis*, Lesquereux, 1874.)
Dakota Group; Kansas, U.S.A.
- **salinæ**, Lesquereux, 1873, p. 423. (Re-named *Menispermites salina* by Knowlton, 1898.)
Ibid.
- **salisburyæfolia**, Lesquereux, 1868, p. 94. (= *Cissites salisburyæfolius*, Lx., 1883.)
Dakota Group; Nebraska, U.S.A.
- **Sternbergii**, Lesquereux, 1892, p. 45, pl. vii, figs. 8, 9.
Dakota Group; Kansas, U.S.A.
- **tenuifolius**, Berry, 1905 E, p. 69, pl. xlix, fig. 7.
Matawan Formation; New Jersey, U.S.A.
- **Winchelli**, Lesquereux, 1895, p. 12, pl. B, fig. 2.
Dakota Group; Minnesota, U.S.A.

- Populocaulis yezoensis*, Stopes & Fujii, 1910, pp. 63-64, pl. viii, fig. 49. Upper Cretaceous; Hokkaido, Japan.
- Populophyllum crassinerve*, Fontaine, 1889, p. 312, pl. clviii, fig. 4. Potomac Formation; Virginia, U.S.A.
- *hederæforme*, Fontaine, 1889, p. 311, pl. clxvi, fig. 3. Ibid.
- *menispermoides*, Ward, 1905, p. 498, pl. ex, figs. 2-4. Ibid.
- *minutum*, Ward, 1905, p. 499, pl. cvii, fig. 9. Ibid.
- *reniforme*, Fontaine, 1889, p. 311, pl. clv, fig. 9; pl. clvi, fig. 3. Ibid.
- Populus acerifolia*, Newberry, 1870, p. 65; & 1878, pl. xiii, figs. 5-8. Laramie Formation. (Recorded Kurtz, 1902, p. 51 from Cenomanian, Argentina.)
- *amissa*, Heer, 1882, p. 65, pl. xxviii, fig. 18. Atane Beds; Greenland.
- ? *apiculata*, Newberry in Hollick, 1893, p. 31, pl. iii, fig. 2; & Newberry, 1895, p. 65, pl. xv, figs. 3, 4. Raritan Formation; Staten Island, U.S.A.
- *aristolochioides*, Lesquereux, 1876 A, p. 393. Dakota Group; Kansas, U.S.A.
- *auriculata*, Ward, 1895 A, p. 356, pl. iv, fig. 4. Potomac Formation; Virginia, U.S.A.
- *Berggreni*, Heer, 1871, p. 1183; & 1874 A, p. 106, pl. xxix, figs. 1-5. Atane Beds; Greenland.
- *cordata*, Heer (*non* Newberry), 1871, p. 1183 [nomen nudum]. Ibid.
- *cordifolia*, Newberry, 1870, p. 18; & 1878, pl. iii, fig. 7. Dakota Group; Nebraska, U.S.A.
- *cretacea*, Knowlton in Stanton & Hatcher, 1905, p. 138, pl. xvii, figs. 1-5. Judith River Beds; Montana, U.S.A.
- *cyclophylla*, Heer in Meek & Hayden, 1859, p. 266. (= *Populites cyclophylla* (Heer), Lesquereux, 1868, & *Cissites cyclophylla*, Lx., 1876.) Dakota Group; Nebraska, U.S.A.
- ? *Debeyana*, Heer in Capellini & Heer, 1867, p. 14, pl. i, fig. 1. (= *Juglans Debeyana* (Heer), Lesquereux, 1868.) Ibid.
- *denticulata*, Heer, 1883 A, p. 20, pl. lv, fig. 5. Patoot Beds; Greenland.
- *elliptica*, Newberry, 1870, p. 16; & 1878, pl. iii, figs. 1, 2. Dakota Group; Nebraska, U.S.A.
- *flabellum*, Newberry, 1863, p. 524; & 1898, p. 44, pl. xx, fig. 4. Cretaceous; Washington, U.S.A.
- *harkeriana*, Lesquereux, 1892, p. 44, pl. xlvi, fig. 4. Dakota Group; Kansas, U.S.A.
- *hyperborea*, Heer, 1871, p. 1183; & 1874 A, p. 106, pl. xxvii, fig. 8 d; pl. xxix, figs. 6, 7 a, 8 a, 9; pl. xxx, fig. 2 b. Atane Beds; Greenland.
- *Kansaseana*, Lesquereux, 1892, p. 42, pl. xvii, figs. 1-7. Dakota Group; Kansas, U.S.A.

- Populus lancastriensis*, Lesquereux, 1868, p. 93. (= *Populites lancastriensis*, Lesquereux, 1874.) Dakota Group; Nebraska, U.S.A.
- *latidentata*, Dawson, 1886, p. 16. Belly River Series; Canada.
- *Leuce* (Rossin.), Unger, 1850 A, p. 417. (Recorded by Heer in Meek & Hayden, 1859, p. 265, from Dakota Cretaceous.)
- *litigiosa*, Heer in Capellini & Heer, 1867, p. 18, pl. i, fig. 2. Dakota Group; Nebraska, U.S.A.
- *litigiosa denticulata*, Lesquereux, 1892, p. 47 [nomen nudum].
- *longior*, Dawson, 1883, p. 27. Upper Cretaceous; Vancouver Island, Canada.
- *melanaria*, Heer in Lesquereux, 1878 B, p. 173, pl. lxiv, fig. 5. Montana Formation; Wyoming, U.S.A.
- *melanarioides*, Lesquereux, 1876 C, p. 379; & 1878 B, p. 174, pl. lxii, fig. 5. Ibid.
- *microphylla*, Newberry, 1870, p. 17; 1878, pl. iii, fig. 5; & 1898, p. 46, pl. iii, fig. 5. Dakota Group; Nebraska, U.S.A.
- *obovata*, Knowlton, 1900 A, p. 34, pl. vii, fig. 4. Montana Formation; Wyoming, U.S.A.
- *orbicularis* (Newberry), Berry, 1909, p. 250 (re-naming *Phyllites orbicularis*, Newberry, 1895). Raritan Formation; New Jersey, U.S.A.
- *potomacensis*, Ward, 1895 A, p. 356, pl. iv, figs. 1-3. Potomac Formation; Virginia, U.S.A.
- *primæva*, Heer, 1871, p. 1182; & 1874 A, p. 88, pl. xxiv, fig. 6. Kome Beds; Greenland.
- ? *problematica*, Knowlton in Weed & Knowlton, 1893, p. 15, pl. vi, figs. 5, 6. Laramie Group; Montana, U.S.A.
- *protczadachii*, Dawson, 1883, p. 26, pl. vii, fig. 25. Upper Cretaceous; British Columbia.
- *rectinervata*, Dawson, 1883, p. 27, pl. vii, fig. 26. Upper Cretaceous; Vancouver Island, Canada.
- *rhomboides*, Lesquereux, 1859, p. 360; & in Newberry, 1898, p. 51, pl. xx, figs. 1, 2. Ibid.
- ? *Ricei*, Fontaine in Ward, 1905, p. 266, pl. lxix, fig. 10. Shasta Formation; Oregon, etc., U.S.A.
- *stygia*, Heer, 1874 A, p. 107, pl. xxix, fig. 10. Atane Beds; Greenland.
- *temulæformis*, Hosius & von der Marck, 1880, p. 152, pl. xxviii, figs. 43, 44 (45?). Upper Senonian; Haldem, Westphalia.
- *trinervis*, Dawson, 1883, p. 26. Upper Cretaceous; Vancouver Island, Canada.
- *Wardii*, Knowlton, 1900 A, p. 36, pl. vi, fig. 7. Montana Formation; Wyoming, U.S.A.
- *sp.*, Knowlton, 1900 A, p. 37, pl. vii, fig. 5. Ibid.
- *sp.*, Knowlton, 1905, p. 140, pl. xix, fig. 1. Judith River Beds; U.S.A.
- *sp.*, Hollick, 1906 A, p. 50, pl. vii, figs. 16-18 (Aments). Cretaceous; Martha's Vineyard, U.S.A.

- Posidonia cretacea*, Hosius & von der Marck, 1880, p. 134, pl. xxiv, figs. 7, 8, 9. Upper Senonian; Westphalia.
- Potamogeton cretaceus*, Heer, 1883 a, p. 19, pl. lv, figs. 23-24. Patoot Beds; Greenland.
- *priscus*, Feistmantel, 1874, p. 274. Perucer Beds; Bohemia.
- Potamogetophyllum vernonense*, Fontaine in Ward, 1905, p. 500, pl. cix, fig. 7. Older Potomac Formation; Virginia, U.S.A.
- Premna exul*, Velenovsky, 1882 a, p. 213 [nomen nudum]. Cretaceous; Bohemia.
- Premnophyllum exulum*, Velenovsky, 1889, p. 24. (= *Cissophyllum exulum*, Velenovsky, on same page.) Cenomanian; Bohemia.
- *trigonum*, Velenovsky, 1886, p. 51, pl. xviii, fig. 2. Ibid.
- Prepinus japonicus*, Stopes & Kershaw, 1910, p. 396, pl. xxvii, figs. 1, 2; pl. xxviii, fig. 1. Upper Cretaceous; Hokkaido, Japan.
- *statenensis*, Jeffrey, 1908, p. 209, pl. xiii. Raritan Formation; Staten Island, U.S.A.
- *viticetensis*, Jeffrey, 1910 c, p. 336, pl. xxxiii. Magothy or Upper Potomac; Staten Island, U.S.A.
- Protea Haidingeri*, Ettingshausen, 1867 a, p. 254, pl. ii, fig. 12. Cenomanian; Niederschoena, Saxony.
- Proteæphyllum californicum*, Fontaine, 1905 a, p. 267, pl. lxi, fig. 11. Shasta Formation; California, U.S.A.
- *dentatum*, Fontaine, 1889, p. 286, pl. clvi, fig. 7; pl. clxxii, figs. 1, 4; pl. clxxiii, figs. 12, 14. Potomac Formation; Maryland, U.S.A.
- *ellipticum*, Fontaine, 1889, p. 285, pl. cxlii, figs. 1, 2. Potomac Formation; Virginia, U.S.A.
- *oblongifolium*, Fontaine, 1889, p. 284, pl. cxxxix, fig. 5; pl. cxl, figs. 1, 2. Ibid.
- *orbiculare*, Fontaine, 1889, p. 283, pl. cxxxix, fig. 4. Ibid.
- *ovatum*, Fontaine, 1889, p. 285, pl. cxli, fig. 1. Ibid.
- *reniforme*, Fontaine, 1889, p. 282, pl. cxxxix, fig. 3; pl. clvi, fig. 4; pl. clx, figs. 1, 2. Ibid.
- *tenuinerve*, Fontaine, 1889, p. 286, pl. cl, fig. 13; pl. clvi, fig. 2. Ibid.
- *Uhleri*, Fontaine in Ward, 1905, p. 564, pl. cxviii, fig. 5. Older Potomac Formation; Maryland, U.S.A.
- *sp. ?*, Fontaine, 1889, p. 284, pl. cxxxix, fig. 2. Potomac Formation; Virginia, U.S.A.
- Proteoides acuta*, Heer in Capellini & Heer, 1867, p. 17, pl. iv, figs. 7, 8. Dakota Group; Kansas, U.S.A.
- *affinis*, Schenk, 1876, p. 169, pl. xxix, fig. 14. Upper Cretaceous; Brandenburg, Tyrol.
- *australiensis*, Ettingshausen, 1893, pp. 140, 149; & 1895, p. 26, pl. iii, fig. 16. Cretaceous; Australia.
- *crassipes*, Heer (*non* Ettingshausen & Gardner), 1871, p. 1183; & 1874 a, p. 110, pl. xxxi, figs. 6-8 a. (= *Ficus crassipes*, Heer, 1882.) Atane Beds; Greenland.

- Proteoides daphnogenoides*, Heer in Capellini & Heer, 1867, p. 17,
pl. iv, figs. 9, 10. Dakota Group; Nebraska, U.S.A.
- *Ettingshauseni*, Schenk, 1876, p. 170, pl. xxix, fig. 13.
Upper Cretaceous; Brandenburg, Tyrol.
- *granulatus*, Heer, 1871, p. 1183; & 1874 A, p. 111, pl. xxxi,
figs. 11, 11 b. Atane Beds; Greenland.
- *grevilleæformis*, Heer in Capellini & Heer, 1867, p. 17, pl. iv,
fig. 11. Dakota Group; Kansas, U.S.A.
- *cf. grevilleæformis*, Heer in Kerner, 1896, p. 54, pl. v, fig. 2.
Cretaceous; Lesina, Dalmatia.
- *ilicoides*, Heer, 1871 A, p. 13, pl. iii, figs. 7, 8.
Senonian; Quedlinburg, Saxony.
- *lancifolius*, Heer, 1871 A, p. 12, pl. iii, figs. 5, 6. Ibid.
- *longus*, Heer, 1871, p. 1183; & 1874 A, p. 110, pl. xxxi, figs. 4, 5;
pl. xxix, fig. 8 b. (= *Myrica longa*, Heer, 1882.)
Atane Beds; Greenland.
- *major*, Dawson, 1893, p. 61, pl. xii, fig. 54.
Upper Cretaceous; Vancouver Island, Canada.
- *Neillii*, Dawson, 1894, p. 61, pl. xii, fig. 53. Ibid.
- *Reussi*, Engelhardt, 1892, p. 105. (= *Salix macrophylla*, Reuss.)
Cenomanian; Bohemia.
- *vexans*, Heer, 1871, p. 1183; & 1874 A, p. 111, pl. xxxi, figs. 9, 10.
Atane Beds; Greenland.
- *sp.*, Dawson, 1894, p. 61, pl. xiii, fig. 55.
Upper Cretaceous; Vancouver Island, Canada.
- Proteophyllum coriaceum*, Velenovsky, 1889, p. 18, pl. iv, fig. 13;
pl. vi, fig. 15. Cenomanian; Bohemia.
- *cornutum*, Velenovsky, 1889, p. 18, pl. iv, fig. 12. Ibid.
- *daphnoides*, Saporta, 1894, p. 203, pl. xxxvi, fig. 3.
Upper Albian; Portugal.
- *decorum*, Velenovsky, 1889, p. 18, pl. v, fig. 13.
Cenomanian; Bohemia.
- *demersum*, Saporta, 1894, p. 203, pl. xxxvii, fig. 15.
Upper Albian; Portugal.
- *dissectum*, Saporta, 1894, p. 156, pl. xxx, figs. 13, 14; pl. xxxi,
figs. 15-16; pl. xxxiv, fig. 17. Ibid.
- *laminarium*, Velenovsky, 1889, p. 18, pl. iv, fig. 7.
Cenomanian; Bohemia.
- *Launayi*, Zeiller, 1905, p. 342, pl. vii, figs. 12, 13, 13 a.
Upper Cretaceous; Balkans.
- *leucospermoides*, Saporta, 1894, p. 185, pl. xxx, fig. 15.
Albian; Portugal.
- *oblongatum*, Saporta, 1894, p. 204, pl. xxxvi, fig. 4. Ibid.
- *oxyanthæmorphum*, Saporta, 1894, p. 185, pl. xxxv, fig. 8.
Ibid.
- *paucidentatum*, Velenovsky, 1889, p. 18, pl. vi, figs. 12, 13.
Cenomanian; Bohemia.
- *productum*, Velenovsky, 1889, p. 18, pl. iv, figs. 10, 11. Ibid.

- Proteophyllum Saportanum**, Velenovsky, 1889, p. 18, pl. v, fig. 14.
Cenomanian; Bohemia.
- **trifidum**, Velenovsky, 1889, p. 18, pl. vi, fig. 14; pl. v, fig. 15.
Ibid.
- **truncatum**, Saporta, 1894, p. 203, pl. xxxvii, fig. 16.
Upper Albian; Portugal.
- Proteopsis Proserpinæ**, Velenovsky, 1889, p. 19, pl. i, figs. 6-9.
Cenomanian; Vyserovic, Bohemia.
- Protocyathea trichinopoliensis**, Feistmantel, 1877, p. 136, pl. i, figs. 1-2.
Neocomian (?); India.
- **Ungeri** (Unger), Feistmantel, 1877, p. 136 [nomen nudum = *Caulopteris cyatheoides*, Unger].
Neocomian; Austria.
- Protodammara speciosa**, Hollick & Jeffrey, 1906, p. 199, pl. i, figs. 5-13; pl. ii, figs. 1-5. (= *Dammara minor*, Hollick.)
Raritan Formation; Staten Island, U.S.A.
- Protophyllocladus lanceolatus** (Knowlton), Berry, 1903 D, p. 441 (re-naming *Thinnfeldia lanceolata*, Knowlton, 1893).
Laramie Formation; Montana, U.S.A.
- **polymorphus** (Lesquereux), Berry, 1903 D, p. 442 (re-naming *Salisburya polymorpha*, Lesquereux, 1859, & *Thinnfeldia polymorpha*, Knowlton, 1898).
- **subintegrifolius** (Lesquereux), Berry, 1903 A, p. 440; & 1904, p. 69, pl. i, fig. 5 (re-naming *Thinnfeldia subintegrifolia* (Lx.), Knowlton, 1898, & *Thinnfeldia Lesquereuxiana*, Heer, & *Phyllocladus subintegrifolius*, Lesquereux).
- Cretaceous; Staten Island, U.S.A.
- Protophyllum boreale**, Dawson, 1883, p. 23, pl. iv, fig. 13.
Upper Cretaceous; North-West Territory, Canada.
- **crassum**, Lesquereux, 1892, p. 193, pl. lxx, fig. 4.
Dakota Group; Kansas, U.S.A.
- **crednerioides**, Lesquereux, 1876 B, p. 363, pl. iii, fig. 1; pl. viii, fig. 4.
Ibid.
- **crenatum**, Knowlton in Lesquereux, 1892, p. 190, pl. lxx, fig. 7.
Ibid.
- **denticulatum**, Lesquereux, 1892, p. 193, pl. xxxvi, fig. 9.
Ibid.
- **dimorphum**, Lesquereux, 1892, p. 190, pl. xli, fig. 1.
Ibid.
- **Haydenii**, Lesquereux, 1874, p. 106, pl. xvii, fig. 3 (= *Pterospermites Haydenii*, Lesquereux, 1872).
Ibid.
- **integerrimum**, Lesquereux, 1892, p. 192, pl. xliii, fig. 3; & 1895, p. 21, pl. B, figs. 10, 11.
Dakota Group; Minnesota, U.S.A.
- **Leconteanum**, Lesquereux, 1874, p. 103, pl. xvii, fig. 4; pl. xxvi, fig. 1. (= *Credneria Le Conteana*, Lesquereux, 1868.)
Dakota Group; Kansas, U.S.A.
- **minus**, Lesquereux, 1874, p. 104, pl. xix, fig. 2; pl. xxvii, fig. 1.
Ibid.
- **Mudgii**, Lesquereux, 1874, p. 106, pl. xviii, fig. 3. (= *Quercus Mudgii*, Lx., 1872.)
Ibid.

- Protophyllum multinerve**, Lesquereux, 1874, p. 105, pl. xviii, fig. 1.
(=*Pterospermites multinervis*, Lesquereux, 1872.)
Dakota Group; Kansas, U.S.A.
- **Nanaimo**, Dawson, 1883, p. 28, pl. viii, fig. 35.
Upper Cretaceous; Vancouver Island, Canada.
- **nebrascense**, Lesquereux, 1874, p. 103, pl. xxvii, fig. 3.
Dakota Group; Nebraska, U.S.A.
- **obovatum**, Newberry, 1895, p. 128, pl. xxxviii, fig. 4. (= *Pterospermites obovatus*, Berry, 1909.) Amboy Clay; Woodbridge, U.S.A.
- **præstans**, Lesquereux, 1892, p. 188, pl. xli, figs. 2, 3; pl. xlii, figs. 3, 4.
Dakota Group; Kansas, U.S.A.
- **pseudospermoides**, Lesquereux, 1892, p. 194, pl. lix, fig. 2. Ibid.
- **ptercspermifolium**, Lesquereux, 1892, p. 195, pl. lix, fig. 1. Ibid.
- **quadratum**, Lesquereux, 1874, p. 104, pl. xix, fig. 1. (= *Pterospermites quadratus*, Lesquereux, 1872.) Ibid.
- **querciforme**, Hollick, 1895, p. 227, pl. cccxxviii, fig. 1. Ibid.
- **rugosum**, Lesquereux, 1874, p. 105, pl. xvii, figs. 1-2; pl. xix, fig. 3. (= *Pterospermites rugosus*, Lesquereux, 1873.) Ibid.
- **Sternbergii**, Lesquereux, 1874, p. 101, pl. xvi; pl. xviii, fig. 2. (= *Pterospermites Sternbergii*, Lesquereux, 1873.) Ibid.
- ? **trilobatum**, Lesquereux, 1876 A, p. 397. Ibid.
- **undulatum**, Lesquereux, 1892, p. 189, pl. xlii, fig. 2. Ibid.
- **sp.**, Dawson, 1894, p. 63, pl. xi, figs. 46, 47.
Upper Cretaceous; Vancouver Island, Canada.
- Protopteris Buvignieri**, Brongniart, 1849 A, p. 111 [nomen nudum].
Lower Cretaceous; France.
- **Cotteana**, Presl in Sternberg, 1838, p. 170, pl. lxx, figs. 4-6; pl. lxxvii; & Stenzel, 1886, p. 25. Turonian; Oppeln.
- **Duplessyana**, Saporta, 1880, p. 646, pl. ii; pl. iii, figs. 1, 2.
Albian; Havre, France.
- **fibrosa**, Stenzel, 1886, p. 25, pl. iii, figs. 30-36. Turonian; Oppeln.
- **punctata**, (Sternberg), Presl in Sternberg, 1838, p. 170, pl. lxx, figs. 1-3. (= *Lepidodendron punctata*, Sternberg, 1820, = *Dicksonia punctata*, Heer.) Greensand; Bohemia.
- **Singeri**, Presl in Sternberg, 1838, p. 171, pl. lxx, fig. 7. (= *Caulopteris Singeri*, Goeppert, 1836, removed from Cretaceous by Debey & Ettingshausen, 1859 B, p. 245.) Cretaceous; Bohemia.
- **Sternbergii**, Corda, 1845, p. 77, pl. xlvi, fig. 1. Ibid.
- **Wohlegemuthi**, Fliche, 1896, p. 132, pl. i, fig. 2.
Albian; Clermont, France.
- Protorhipis cordata**, Heer, 1882, p. 10, pl. iii, fig. 11. (= *Hausmannia cordata* (Heer), Richter, 1906.) Kome Beds; Greenland.
- **Fisheri**, Knowlton, 1907, p. 114, pl. xii, fig. 3.
Kootanie Formation; Montana, U.S.A.
- Protorrhipis Choffati**, Saporta, 1891, p. 252; & 1894, p. 144, pl. xxii, figs. 9-11; pl. xxvi, figs. 17-18; pl. xxvii, figs. 1-5.
Urgonian; Portugal.

- Prunus? acutifolia*, Newberry, 1895, p. 90, pl. xiv, fig. 1.
Amboy Clay; Woodbridge, U.S.A.
- (*Amygdalus*)? *antecedens*, Lesquereux, 1892, p. 144, pl. lv, fig. 4.
Dakota Group; Kansas, U.S.A.
- *cerasiformis*, Velenovsky, 1887, p. 69, pl. xxix, figs. 2, 7.
Cenomanian; Bohemia.
- *cretaceus*, Lesquereux, 1868, p. 102; & 1874, p. 111, pl. xxiii, figs. 8, 9.
Dakota Group; Nebraska, U.S.A.
- *Parlatorii* (Heer), Lesquereux, 1868, p. 102. (= *Andromeda Parlatorii*, Heer, 1867.)
Dakota Group; Kansas, U.S.A.
- Psamopteris knorriæformis*, Eichwald, 1861, p. 304.
Greensand; Russia.
- Pseudo-Araucaria Lamberti*, Fliche, 1896, p. 193, pl. vii, fig. 2, text-fig. 7.
Albian; Grandpré, France.
- *Loppinetti*, Fliche, 1896, p. 189, pl. vi, figs. 3-5.
Albian; Clermont, France.
- *major*, Fliche, 1896, p. 191, pl. vii, fig. 1.
Albian; Islettes, France.
- Pseudoasterophyllites cretaceus* (Feistmantel), Velenovsky, 1887 A, p. 643, on plate figs. 19-25. (= *Asterophyllites cretaceus*, Feistmantel, 1874.)
Cenomanian; Bohemia.
- Pseudocycas Dicksoni*, Nathorst, 1907, p. 8. (= *Cycadites Dicksoni*, Heer, 1874 A, = *Cycas Dicksoni*, Heer, 1882.)
Atane Beds; Greenland.
- *insignis*, Nathorst, 1907, p. 7, pl. i, figs. 1-5; pl. ii, figs. 1-9; pl. iii, fig. 1.
Cenomanian; Greenland.
- *pumilio*, Nathorst, 1907, p. 7, pl. i, figs. 6, 7.
Ibid.
- *Steenstrupi*, Nathorst, 1907, p. 8, pl. ii, figs. 10, 11. (= *Cycas Steenstrupi*, Heer, 1882.)
Ibid.
- Pseudofrenelopsis Felixi*, Nathorst in Felix & Nathorst, 1893, p. 52, text-figs. 6-9.
Neocomian; Mexico.
- Pseudogeinitzia sequoiiformis*, Hollick & Jeffrey, 1909, p. 45, pl. x, fig. 11; pl. xxv, fig. 4.
Raritan Formation; Staten Island, U.S.A.
- Pseudostrobus Guillieri*, Crie, 1879 (?), p. 30 [nomen nudum].
Cretaceous; Sainte-Croix, France.
- Ptenostrobus nebrascensis*, Lesquereux, 1874, p. 114, pl. xxiv, fig. 1.
Dakota Group; Nebraska, U.S.A.
- Pteridolemma aneimiifolium*, Debey & Ettingshausen, 1859 B, p. 230, pl. vii, fig. 1.
Senonian; Aix, Rhenish Prussia.
- *antiquum*, Debey & Ettingshausen, 1859 B, p. 232, pl. vii, figs. 4-5, 10.
Ibid.
- *arborescens*, Debey & Ettingshausen, 1859 B, p. 237, pl. vii, figs. 11-13.
Ibid.
- *Benincasæ*, Debey & Ettingshausen, 1859 B, p. 225, pl. vi, figs. 1-5.
Ibid.
- *deperditum*, Debey & Ettingshausen, 1859 B, p. 233, pl. vii, fig. 8.
Ibid.

- Pteridoleimma dictyoides**, Debey & Ettingshausen, 1859 b, p. 236,
pl. vi, figs. 15-17. Senonian; Aix, Rhenish Prussia.
- **dubium**, Debey & Ettingshausen, 1859 b, p. 231, pl. vii, figs. 2-3.
Ibid.
- **durum**, Bayer, 1896, p. 5, text-figs. 3, 4, & p. 31.
Upper Senonian; Bohemia.
- **Elisabethæ**, Debey & Ettingshausen, 1859 b, p. 222, pl. v, figs. 5-9.
Senonian; Aix, Rhenish Prussia.
- **gymnorhachis**, Debey & Ettingshausen, 1859 b, p. 234, pl. vii,
figs. 21-22. Ibid.
- **Haidingeri**, Debey & Ettingshausen, 1859 b, p. 228, pl. vi, fig. 10.
Ibid.
- **Heissianum**, Debey & Ettingshausen, 1859 b, p. 227, pl. vi, figs. 9,
14. Ibid.
- **Kaltenbachi**, Debey & Ettingshausen, 1859 b, p. 233, pl. vii, fig. 9.
Ibid.
- **Koninckianum**, Debey & Ettingshausen, 1859 b, p. 224, pl. v,
figs. 1-4. Ibid.
- **leptophyllum**, Debey & Ettingshausen, 1859 b, p. 235, pl. vii,
figs. 18-19. Ibid.
- **Michelisi**, Debey & Ettingshausen, 1859 b, p. 229, pl. vi, figs. 11-12.
Ibid.
- **odontopteroides**, Debey & Ettingshausen, 1859 b, p. 234, pl. vii,
fig. 20. Ibid.
- **orthophyllum**, Debey & Ettingshausen, 1859 b, p. 227, pl. vi,
fig. 6. Ibid.
- **pecopteroides**, Debey & Ettingshausen, 1859 b, p. 226, pl. vi,
figs. 7-8. Ibid.
- **phycomorpha**, Saporta, 1894, p. 171, pl. xxviii, figs. 7-9, 12.
Albian; Portugal.
- **pseudadiantum**, Debey & Ettingshausen, 1859 b, p. 235, pl. vii,
figs. 14-17. Senonian; Aix, Rhenish Prussia.
- **Ritzianum**, Debey & Ettingshausen, 1859 b, p. 224, pl. v, figs. 10-
12. Ibid.
- **Serresi**, Debey & Ettingshausen, 1859 b, p. 229, pl. vi, figs. 13,
18-19. Ibid.
- **spoliatum**, Saporta, 1894, p. 85, pl. xvi, fig. 25; pl. xviii, fig. 7.
Valanginian; Portugal.
- **tripartitum**, Saporta, 1894, p. 85, pl. xv, fig. 1. Ibid.
- **Waterkeyni**, Debey & Ettingshausen, 1859 b, p. 231, pl. vii,
figs. 6-7. Senonian; Aix, Rhenish Prussia.
- Pteridophyllum fastigatum**, Schulze, 1888, p. 11 [nomen nudum].
Neocomian; Helmstein.
- Pteris? Albertini** (Dunker), Velenovsky, 1888 b, p. 15, pl. iv, figs. 5-10.
Perucer Beds; Bohemia.
- **Albertsii** (Dunker), Heer, 1882, p. 29, pl. xvi, figs. 5, 6; pl. xxviii,
figs. 1-3; pl. xlvi, figs. 22-24. Atane Beds; Greenland.

- Pteris dakotensis**, Lesquereux, 1892, p. 24, pl. i, figs. 2, 3.
Dakota Group; Kansas, U.S.A.
- **frigida**, Heer, 1882, p. 3, pl. ii, fig. 13; p. 25, pl. vi, fig. 5 b; pl. x, figs. 1-4; pl. xi; pl. xii, fig. 2; pl. xiii, fig. 2; pl. xvi, figs. 1, 2; pl. xviii, fig. 10 b. Kome Beds & Atane Beds; Greenland.
- (**Oleandra**) **glossopteroides**, Dawson, 1883, p. 24, pl. iv, fig. 16.
Upper Cretaceous; Protection Island, Canada.
- **longipennis**, Heer, 1882, p. 28, pl. x, figs. 5-13; pl. xiii, fig. 1.
Atane Beds; Greenland.
- **nebraskana**, Heer in Lesquereux, 1874, p. 46 (?), pl. xxix, fig. 5.
(Named *Pecopteris* (?) *nebraskana* (?), p. 46.)
Dakota Group; Kansas, U.S.A.
- **Reichiana**, Ettingshausen, 1867 A, p. 243.
Cenomanian; Niederschoena, Saxony.
- **Russellii**, Newberry, 1883, p. 503; & Hollick in Newberry, 1898, p. 7, pl. li, figs. 1, 1 a. Laramie Formation; New Mexico.
- **slivnecensis**, Marik, 1901, p. 7, pl. i, fig. 18.
Cenomanian; Bohemia.
- Pterophyllum alaskense**, Fontaine in Ward, 1905, p. 152, pl. xxxviii, figs. 19, 20.
Lower Cretaceous; Alaska.
- **blechniforme**, Hosius & von der Marck, 1880, p. 212, pl. xxiv, fig. 197.
Neocomian; Westphalia.
- **Buchianum**, Ettingshausen, 1852 c, p. 21, pl. i, fig. 1. (= *Dicranites Buchianus*, Bornemann, 1856.)
Wealden; Silesia.
- **californicum**, Fontaine in Diller & Stanton, 1894, p. 450 [nomen nudum].
Horsetown Beds; California, U.S.A.
- **concinnum**, Heer, 1874 A, p. 68, pl. xiv, figs. 15-20; pl. xv, fig. 5 b.
Kome Beds; Greenland.
- **cretosum**, Reich in Cotta, 1836, p. 585; and in Goeppert, 1847, p. 362, pl. xxxviii, fig. 4. (= *Pterozamites* (?) *cretosus*, Miquel, 1861.)
Cenomanian; Niederschoena, Saxony.
- **Ernestinae**, Stiehler, 1857, p. 454; & 1858, p. 76, pl. xv. (= *Pterozamites Ernestinae*, Miquel, 1861.)
Senonian; Quedlinburg, Saxony.
- **Germari**, Otto, 1854, p. 36, pl. v, figs. 3, 4. Quader; Saxony.
- **Haydenii**, Lesquereux, 1868, p. 91; & 1874, p. 49, pl. i, fig. 6.
(= *Podozamites Haydenii*, Lesquereux, 1888, and in part = *Abietites Ernestinae*, Lesquereux, 1874, p. 49, pl. i, fig. 7.)
Dakota Group; Nebraska, U.S.A.
- **lepidum**, Heer, 1874 A, p. 68, pl. xvi, figs. 1-3 b.
Kome Beds; Greenland.
- ? **lowryanum**, Ward, 1905, p. 254, pl. lxvii, fig. 9.
Shasta Formation; California, U.S.A.
- **montanense** (Fontaine), Knowlton, 1907, p. 122, pl. xiv, fig. 3.
(= *Zamites montanensis*, Fontaine, 1893.)
Kootanie Formation; Montana, U.S.A.
- **Reichianum**, Engelhardt, 1892 A, p. 89, pl. ii, figs. 1, 2.
Cenomanian; Niederschoena, Saxony.

- Pterophyllum saxonicum**, Reichenbach in Geinitz, 1842, p. xxii; and in Goeppert, 1847, p. 362, pl. xxxviii, fig. 13. (= *Pterozamites* (?) *saxonicus*, Miquel, 1861.) Cenomanian; Niederschoena, Saxony.
- Pterocpermites auriculatus**, Heer, 1882, p. 95, pl. xxvii, fig. 4.
Atane Beds; Greenland.
- **carolinensis**, Berry, 1907, p. 198, pl. xiv, fig. 2.
Middle Cretaceous; North Carolina, U.S.A.
- **cordifolius**, Heer, 1882, p. 94, pl. xxvii, figs. 2-3.
Atane Beds; Greenland.
- **credneriifolius**, Berry, 1907, p. 199, pl. xiii, fig. 4.
Middle Cretaceous; North Carolina, U.S.A.
- **Haydenii**, Lesquereux, 1872, p. 302. (= *Protrophyllum Haydenii*, Lesquereux, 1874.) Dakota Group; Kansas, U.S.A.
- **longeacuminatus**, Lesquereux, 1892, p. 186, pl. lix, fig. 3. Ibid.
- **modestus**, Lesquereux, 1892, p. 186, pl. lviii, fig. 5. Ibid.
- **multinervis**, Lesquereux, 1872, p. 302. (= *Protrophyllum multinerve*, Lesquereux, 1874.) Ibid.
- **obovatus** (Newberry), Berry, 1909, p. 259. (= *Protrophyllum obovatum*, Newberry, 1895.) Amboy Clay; New Jersey, U.S.A.
- **quadratus**, Lesquereux, 1872, p. 301. (= *Protrophyllum quadratum*, Lesquereux, 1874.) Dakota Group; Kansas, U.S.A.
- **rugosus**, Lesquereux, 1873, p. 426. (= *Protrophyllum rugosum*, Lesquereux, 1874.) Ibid.
- **Sternbergii**, Lesquereux, 1873, p. 425. (= *Protrophyllum Sternbergii*, Lesquereux, 1874.) Ibid.
- **undulatus**, Knowlton, 1900 A, p. 67, pl. xvi, fig. 3; pl. xvii, fig. 2; pl. xviii, fig. 4. Montana Formation; Wyoming, U.S.A.
- **Wardii**, Knowlton, 1900 A, p. 66, pl. xv, fig. 4; pl. xvi, fig. 1. Ibid.
- Pterocpermum cretaceum**, Unger, 1865, p. 376, pl. i, figs. 2-3.
Cretaceous; Transylvania.
- Pterozamites Ernestinae**, Miquel, 1861, p. 29. (= *Pterophyllum Ernestinae*, Stiehler, 1857.) Senonian; Quedlinburg, Saxony.
- ? **saxonicus**, Miquel, 1861, p. 29. (= *Pterophyllum saxonicum*, Reichenbach, 1842.) Cenomanian; Niederschoena, Saxony.
- Puccinites cretaceus**, Velenovsky, 1889, p. 26, pl. iii, fig. 14, & pp. 48 & 52. (= *Uredinites cretaceus*, Velenovsky, same work, same plate, & p. 29.) Perucer Beds; Bohemia.
- Pycnois speciosa**, Stenzel, 1872, p. 71 [nomen nudum]. Cretaceous (?).
- Pyrus cretacea**, Newberry, 1870, p. 12; & 1878, pl. iii, fig. 7; & 1898, p. 110, pl. i, fig. 7. Dakota Group; Kansas, U.S.A.
- Quercophyllum chinkapinense**, Ward, 1905, p. 513, pl. cxii, figs. 3, 4. Older Potomac Formation; Virginia, U.S.A.
- **grossedentatum**, Fontaine, 1889, p. 307, pl. clvi, fig. 9. Ibid.
- **tenuinerve**, Fontaine, 1889, p. 308, pl. cxlix, figs. 6, 7. Ibid.
- **wyomingense**, Fontaine in Ward, 1899 B, p. 688, pl. clxix, fig. 6. Lower Cretaceous; Black Hills, U.S.A.

- Quercus acrodon*, Lesquereux (*non* Massalongo), 1878 b, p. 158, pl. xix, figs. 11-13. (= *Quercus Lesquereuxiana*, Knowlton, 1898.)
Laramie Formation; Wyoming, U.S.A.
- *alnoides*, Lesquereux, 1892, p. 54, pl. vii, fig. 3.
Dakota Group; Kansas, U.S.A.
- *anceps*, Lesquereux, 1868, p. 96.
Dakota Group; Nebraska, U.S.A.
- *antiqua*, Newberry, 1870, p. 26; & 1898, p. 69, pl. xiii, fig. 2.
Dakota Group; Utah, U.S.A.
- *asymetra*, Hosius & von der Marek, 1880, p. 165, pl. xxxi, fig. 82.
Upper Senonian; Haldem, Westphalia.
- *banksiæfolia*, Newberry, 1863, p. 522; & 1898, p. 69, pl. xviii, figs. 2-5.
Cretaceous; Washington, U.S.A.
- *Benzoin*, Lesquereux, 1859, p. 340. (= *Persea Leconteana*, Lesquereux, 1874.)
Cretaceous (?); Vancouver Island, Canada.
- *Beyrichii*, Ettingshausen, 1867 a, p. 248, pl. ii, fig. 2.
Cenomanian; Niederschoena, Saxony.
- *calliprinoides*, Ettingshausen, 1887 a, p. 182, pl. viii, fig. 9.
Upper Cretaceous; New Zealand.
- *castanoides*, Hosius & von der Marek (*non* Goeppert), 1880, p. 163, pl. xxx, figs. 76, 77.
Upper Senonian; Haldem, Westphalia.
- *colpophylla*, Ettingshausen, 1893, p. 134; & 1895, p. 19, pl. i, figs. 22, 23.
Cretaceous; Australia.
- *competens*, Lesquereux, 1876 c, p. 370.
Montana Formation (?); Wyoming, U.S.A.
- *coriacea*, Newberry, 1863, p. 521; & 1898, p. 73, pl. xix, figs. 1-3; pl. xx, fig. 5.
Lower Cretaceous; Washington, U.S.A.
- *cuneata*, Hosius (*non* Newberry, *nec* Saporta), 1870 a, p. 97, pl. xiii, fig. 10.
Senonian; Legden, Westphalia.
- *cuneata*, Newberry, 1870, p. 25.
Dakota Group; Nebraska, U.S.A.
- *cuspidigera*, Heer, 1883 a, p. 25, pl. lvi, fig. 22.
Patoot Beds; Greenland.
- *dakotensis*, Lesquereux, 1883, p. 30; & 1892, p. 53, pl. vii, fig. 4.
Dakota Group; Kansas, U.S.A.
- *denticulata*, Heer, 1883 a, p. 25, pl. lvi, fig. 16.
Patoot Beds; Greenland.
- *dentonoides*, Knowlton, 1900 a, p. 40, pl. vii, fig. 7.
Montana Formation; Wyoming, U.S.A.
- *dryandraefolia*, von der Marek, 1864, p. 79, pl. xiii, figs. 6-7.
Upper Senonian; Westphalia.
- *elliptica*, Newberry, 1863, p. 523; & 1898, p. 74, pl. xviii, fig. 1; pl. xx, fig. 3.
Cretaceous; Washington, U.S.A.
- *ellsworthianus*, Lesquereux, 1868, p. 96; & 1874, p. 65, pl. vi, fig. 7. (= *Laurophyllum ellsworthianum*, Lesquereux, 1892.)
Dakota Group; Nebraska, U.S.A.
- *eoprinoideis*, Berry, 1904 a, p. 74, pl. iv, fig. 11.
Matawan Formation; New Jersey, U.S.A.

- Quercus eucalyptoides*, Ettingshausen, 1893, pp. 140, 148; & 1895, p. 18, pl. i, fig. 26. Cretaceous; Australia.
- *euryphylla*, Hosius & von der Marck, 1880, p. 160, pl. xxviii, figs. 48, 49, 50; pl. xxix, fig. 51. Upper Senonian; Haldem, Westphalia.
- *ferox*, Heer, 1882, p. 68, pl. xix, fig. 7. Atane Beds; Greenland.
- *flexuosa*, Newberry, 1863, p. 521; & 1898, p. 74, pl. xix, figs. 4-6. Cretaceous; Washington, U.S.A.
- *formosa*, Hosius & von der Marck, 1880, p. 164, pl. xxxi, fig. 81. Upper Senonian; Haldem, Westphalia.
- *glascoena*, Lesquereux, 1892, p. 55, pl. vi, fig. 6. Dakota Group; Kansas, U.S.A.
- *gracilis*, Newberry (*non* Saporta), 1883, p. 504; & 1898, p. 75, pl. lxvii, fig. 4. Montana Formation; Wyoming, U.S.A.
- *Hatcheri*, Knowlton, 1906, Proc. Biol. Soc. Washington, vol. xxix, p. 95. (= *Quercus montana*, Knowlton, *non* Willd.) Judith River Beds; Montana, U.S.A.
- *hexagona*, Lesquereux, 1868, p. 95; & 1874, p. 64, pl. v, fig. 8. Dakota Group; Nebraska, U.S.A.
- *hieracifolia* (Debey), Hosius & von der Marck, 1880, p. 163, pl. xxxi, figs. 85-88. (= *Dryandroides hieracifolia*, Debey *in litt.*) Upper Senonian; Westphalia.
- *Hollickii*, Berry, 1905 n, p. 71, pl. li, figs. 1, 2. Matawan Formation; New Jersey, U.S.A.
- *Holmesii*, Lesquereux, 1892, p. 58. (= *Dryophyllum (Quercus) Holmesii*, Lesquereux, 1883.) Upper Cretaceous; Vancouver, Canada.
- (*Dryophyllum*) *Hosiana*, Lesquereux, 1892, p. 57, pl. iii, fig. 14. Dakota Group; Kansas, U.S.A.
- ? *iliciformis*, Hosius & von der Marck (*non* Saporta), 1880, p. 165, pl. xxxi, fig. 84. Upper Senonian; Haldem, Westphalia.
- *Johnstrupi*, Heer, 1883 a, p. 24; pl. lvi, figs. 7-11, 11 b, 12 a. Patoot Beds; Greenland.
- *Judithæ*, Knowlton in Stanton & Hatcher, 1905, p. 143, pl. xviii, fig. 2. Judith River Beds; Montana, U.S.A.
- *kanseana* (Lx.), Knowlton, 1898, p. 194 (re-naming *Hamamelites kansaseana*, Lesquereux, 1876 n, and *Alnus kanseana*, Lx., 1878.) Dakota Group; Kansas, U.S.A.
- *Langeana*, Heer, 1883 a, p. 24, pl. lvi, figs. 13-15. Patoot Beds; Greenland.
- *latifolia*, Lesquereux, 1892, p. 58. (= *Dryophyllum latifolium*, Lesquereux, 1876.) Dakota Group; Kansas, U.S.A.
- *latissima*, Hosius, 1870 a, p. 97, pl. xiii, fig. 11. Senonian; Legden, Westphalia.
- *Legdensis*, Hosius, 1870 a, p. 97, pl. xiii, fig. 7. Ibid.
- *Lesquereuxiana*, Knowlton, 1898, p. 194; & 1900 a, p. 39, pl. vii, fig. 1 (re-naming *Quercus acrodon*, Lesquereux, 1878 n). Montana Formation; Wyoming, U.S.A.

- Quercus longifolia*, Hosius, 1870 A, p. 97, pl. xiii, figs. 8-9.
Senonian; Legden, Westphalia.
- *Marioni*, Heer, 1883 A, p. 23, pl. lvi, figs. 1-6.
Patoot Beds; Greenland.
- *montana*, Knowlton in Stanton & Hatcher, 1905, p. 143, pl. xvii,
fig. 6 (antedated by Willd. = *Quercus Hatcheri*).
Judith River Beds; Montana, U.S.A.
- ? *montanensis*, Knowlton, 1900 A, p. 11, pl. i, fig. 10.
Montana Formation; Missouri R., U.S.A.
- *Morrisoniana*, Lesquereux, 1883, p. 40, pl. xvii, figs. 1-2.
Dakota Group; Colorado, U.S.A.
- *Mudgii*, Lesquereux, 1872, p. 302. (= *Protophyllum Mudgii*,
Lesquereux, 1874.) Dakota Group; Kansas, U.S.A.
- *multinervis*, Lesquereux, 1859, p. 360.
Cretaceous (?); Vancouver Island, Canada.
- *Myrtillus*, Heer, 1883 A, p. 25, pl. lvi, figs. 12 b, 17-20.
Patoot Beds; Greenland.
- *negundoides*, Lesquereux, 1872, p. 292.
Laramie Formation; Wyoming, U.S.A.
- *Nelsonica*, Ettingshausen, 1887 A, p. 182, pl. viii, fig. 10.
Upper Cretaceous; New Zealand.
- ? *Novæ-Cæsareæ*, Hollick, 1898 D, p. 131, pl. xiii, figs. 8, 10.
Upper Cretaceous; New Jersey, U.S.A.
- ? (*Dryophyllum*) *occidentalis* (Dawson), Knowlton, 1898,
p. 195 (re-naming *Dryophyllum occidentale*, Dawson, 1894).
Upper Cretaceous; Vancouver, Canada.
- *pachyphylla*, Ettingshausen, 1887 A, p. 181, pl. viii, figs. 8, 8 a.
Upper Cretaceous; New Zealand.
- *patootensis*, Heer, 1883 A, p. 25, pl. lvi, fig. 21.
Patoot Beds; Greenland.
- *paucinervis*, Hosius (*non* Watelet), 1870 A, p. 98, pl. xiii, fig. 12.
Senonian; Legden, Westphalia.
- *platinervis*, Lesquereux, 1859, p. 361.
Cretaceous (?); Vancouver Island, Canada.
- *poranoides*, Lesquereux, 1874, p. 66, pl. xxx, fig. 9.
Dakota Group; Kansas, U.S.A.
- *Pratti*, Berry, 1907, p. 192, pl. xi, fig. 9.
Middle Cretaceous; North Carolina, U.S.A.
- *primordialis*, Lesquereux, 1868, p. 95; & 1874, p. 64, pl. v, fig. 7.
(= *Dryophyllum primordiale*, Lesquereux, 1883.)
Dakota Group; Nebraska, U.S.A.
- *pseudo-chlorophylla*, Ettingshausen, 1893, p. 148; & 1895,
p. 17, pl. ii, fig. 10. Cretaceous; Australia.
- *pseudo-drymeja*, Velenorsky, 1882 A, p. 213; & 1883, p. 42,
pl. x, figs. 21, 22. Senonian; Bohemia.
- *raritanensis*, Berry, 1900, p. 249 (re-naming *Quercus Johnstrupi*,
Newberry (*non* Heer), 1895, p. 69, pl. xix, fig. 7).
Raritan Formation; New Jersey, U.S.A.

- Quercus reticulata*, Eichwald, 1865, p. 62, pl. iii, fig. 16.
Cretaceous; Russia.
- (*Dryophyllum*) *rhamnoides*, Lesquereux, 1892, p. 57, pl. xlviii, fig. 4.
Dakota Group; Kansas, U.S.A.
- *rhomboidalis*, Hosius & von der Marck, 1880, p. 165, pl. xxxi, fig. 82.
Upper Senonian; Haldem, Westphalia.
- *Rinkiana*, Heer, 1882, p. 67, pl. xix, fig. 5.
Atane Beds; Greenland.
- *robusta*, Schulze, 1888, p. 20 [nomen nudum].
Senonian; Heidelberg.
- *rosmarinifolia*, Ettingshausen, 1893, pp. 134, 148; & 1895, p. 19, pl. i, fig. 25.
Cretaceous; Australia.
- *salicifolia*, Newberry, 1870, p. 24; & 1878, pl. ii, fig. 1; & 1898, p. 77, pl. i, fig. 1.
Dakota Group; Nebraska, U.S.A.
- *semi-alatus*, Lesquereux, 1868, p. 96. (= *Anisophyllum semi-alatum*, Lesquereux, 1874.)
Ibid.
- *severnensis*, Berry, 1910 II, p. 22, pl. viii, fig. 3.
Magothy Formation; Severn River, U.S.A.
- *sinuata*, Newberry, 1870, p. 27; & 1898, p. 78, pl. xiii, fig. 1.
Dakota Group; Utah, U.S.A.
- *spathulata*, Eichwald, 1865, p. 62, pl. iii, figs. 9, 10.
Neocomian; Kursk, Russia.
- *sphenobasis*, Hosius & von der Marck, 1880, p. 164, pl. xxx, figs. 78–80.
Upper Senonian; Haldem, Westphalia.
- *spurio-ilex*, Knowlton in Lesquereux, 1892, p. 53, pl. xlviii, fig. 3.
Dakota Group; Kansas, U.S.A.
- *Stokesii*, Ettingshausen, 1893, pp. 133, 148; & 1895, p. 18, pl. i, fig. 21.
Cretaceous; Australia.
- *suspecta*, Lesquereux, 1892, p. 52, pl. xlvii, fig. 7; pl. xlviii, figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- *thulensis*, Heer, 1882, p. 69, pl. xxvi, fig. 7.
Atane Beds; Greenland.
- *trogodytis*, Heer, 1882, p. 69, pl. xxix, fig. 14.
Ibid.
- *Velenovskyi*, Bayer, 1896, pp. 11, 33, text-figs. 7–9.
Upper Senonian; Kieslingswalda, Bohemia.
- *venulosa*, Eichwald, 1865, p. 63, pl. iii, fig. 11.
Neocomian; Kursk, Russia.
- *viburnifolia*, Lesquereux, 1878, p. 159, pl. xx, figs. 11, 12.
Laramie Formation; Wyoming, U.S.A.
- *Victoriæ*, Dawson, 1883, p. 27, pl. vii, fig. 28.
Upper Cretaceous; Vancouver Island, Canada.
- *Wardiana*, Lesquereux, 1892, p. 53, pl. vii, fig. 1.
Dakota Group; Kansas, U.S.A.
- *Warmingiana*, Heer, 1882, p. 68, pl. xix, fig. 6.
Atane Beds; Greenland.
- *westfalica*, Hosius & von der Marck, 1880, p. 161, pl. xxix, figs. 52–63; pl. xxx, figs. 64–75.
Senonian; Haldem, Westphalia.

- Quercus westfalica latior*, Hosius & von der Marek, 1880, p. 161, pl. xxix, figs. 52-59. Senonian; Haldem, Westphalia.
- *westfalica oblonga*, Hosius & von der Marek, 1880, p. 161, pl. xxx, figs. 69-75. Ibid.
- *westfalica obtusata*, Hosius & von der Marek, 1880, p. 161, pl. xxix, figs. 60-63; pl. xxx, figs. 64-68. Ibid.
- *Wilmsii*, Hosius, 1870 A, p. 95, pl. xii, figs. 3-6. Senonian; Westphalia.
- *wyomingiana*, Lesquereux, 1873, p. 400. (Name subsequently cancelled by author). Laramie Formation; Wyoming, U.S.A.
- *sp.*, Berry, 1905 E, p. 72, pl. xlvii, fig. 6. (= *Quercus sp.*, Hollick, 1906 A.) Middle Cretaceous; New Jersey, U.S.A.
- *sp.*, Cornuel, 1866, p. 673, pl. xii, fig. 1. Neocomian; France.
- *sp.*, Ettingshausen, 1895, p. 20, pl. i, figs. 27-28. Cretaceous; Australia.
- *sp.*, Hollick, 1898 D, p. 131, pl. xiv, fig. 9. Middle Cretaceous; New Jersey, U.S.A.
- *sp.*, Hollick, 1906 A, p. 56, pl. viii, fig. 17. Middle Cretaceous; Martha's Vineyard, U.S.A.
- Raphaelia neuropteroides*, Debey & Ettingshausen, 1859 B, p. 220, pl. iv, figs. 23-28; pl. v, figs. 18-20. Senonian; Aix, Rhenish Prussia.
- *Woldrichi*, Marik, 1901, p. 3, pl. i, figs. 3-8. Cenomanian; Bohemia.
- Raritanian gracilis* (Newberry), Hollick & Jeffrey, 1909, p. 26, pl. vi, figs. 4-7; pl. ix, figs. 1-4; pl. x, figs. 14-17; pl. xix, figs. 3-6; pl. xx, fig. 1. (= *Frenelopsis gracilis*, Newb., 1895.) Raritan Formation; Staten Island, U.S.A.
- Raumeria masseiana*, Capellini & Solms-Laubach in Capellini, 1890, p. 446, pl. ii. (= *Cycadeoidea masseiana*, Capellini & Solms-Laubach, 1892.) Cenomanian (?); Italy.
- Ravenalospermum incertissimum*, Saporta, 1894, p. 200, pl. xxxvi, figs. 13-14. Upper Albian; Portugal.
- Rhacoglossum dentatum*, Debey, 1849, p. 299 [nomen nudum]. Senonian; Aix, Rhenish Prussia.
- *heterophyllum*, Debey, 1850, p. 117 [nomen nudum]. Ibid.
- Rhamnites apiculatus*, Lesquereux, 1892, p. 171, pl. xxxvii, figs. 8-13. Dakota Group; Kansas, U.S.A.
- *minor*, Hollick in Newberry, 1895, p. 106, pl. xlii, fig. 36. Amboy Clay; U.S.A.
- Rhamnus? acuta*, Heer, 1882, p. 78, pl. xli, fig. 6; pl. xlv, fig. 13 e. Atane Beds; Greenland.
- *deformatus*, Lesquereux, 1883, p. 126, pl. xx, fig. 6. Laramie Formation; Colorado, U.S.A.
- *discolor*, Lesquereux, 1873, p. 398; & 1878, p. 280, pl. lii, fig. 17. Laramie Formation; Wyoming, U.S.A.
- *elegans*, Newberry, 1870, p. 49; & 1898, p. 117, pl. i, fig. 2. Laramie Formation; Colorado, U.S.A.

- Rhamnus inæquilateralis*, Lesquereux, 1892, p. 170, pl. xxxvii, figs. 4-7. Dakota Group; Kansas, U.S.A.
- *Mudgei*, Lesquereux, 1892, p. 169, pl. xxxvii, figs. 2, 3. Ibid.
- *Novæ-Cæsareæ*, Berry, 1905 n, p. 85, pl. l, figs. 5, 6. Matawan Formation; New Jersey, U.S.A.
- *obovatus*, Lesquereux, 1868, p. 207; & 1878 n, p. 281, pl. liv, figs. 1, 2. Laramie Formation; New Mexico, U.S.A.
- *Oerstedii*, Heer, 1882, p. 98, pl. xxi, figs. 9, 10. Atane Beds; Greenland.
- *Pfaffiana*, Heer, 1883 a, p. 42, pl. lxiv, fig. 8. Patoot Beds; Greenland.
- *prunifolius* Lesquereux, 1883, p. 85; & 1892, p. 169, pl. xxxv, fig. 14. Dakota Group; Kansas, U.S.A.
- *revoluta*, Lesquereux, 1892, p. 171, pl. lxxv, fig. 5. Ibid.
- *Rossmæssleri*, Unger, 1850 a, p. 464. (Recorded Hollick, 1893, Cretaceous, U.S.A.)
- *salicifolius*, Lesquereux, 1868 a, p. 206; & 1878 n, p. 282, pl. liii, figs. 9, 10. Laramie Formation; Colorado, U.S.A.
- *similis*, Lesquereux, 1892, p. 168, pl. xxxv, figs. 12, 13. Dakota Group; Kansas, U.S.A.
- *tenax*, Lesquereux, 1868, p. 101; & 1874, p. 109, pl. xxi, fig. 4. Dakota Group; Nebraska, U.S.A.
- *sp. ?*, Hosius & von der Marck, 1880, p. 175, pl. xxxvi, fig. 131. Upper Senonian; Haldem, Westphalia.
- Rhizocaulon elongatum*, Saporta, 1894, p. 94, pl. xv, figs. 5-6; & p. 146, pl. xxvi, fig. 6. Valanginian & Urgonian; Portugal.
- *najadinum*, Vater, 1884, p. 833, pl. xxvii, fig. 6. Lower Senonian (?); Helmstedt.
- *vetus*, Saporta, 1890 a, p. 814; 1894, p. 94, pl. xv, fig. 4; pl. xvi, fig. 7. Valanginian; Portugal.
- Rhizodendron oppoliense*, Goeppert, 1865 n, p. 397; & in Stenzel, 1886, p. 5, pl. i, figs. 1, 3, 5-12; pl. ii; pl. iii, figs. 20-29. (= *Rhizopterodendron oppoliense*, Goeppert, 1881, = *Caulopteris oppoliensis*, Stenzel, 1897.) Turonian; Oppeln.
- Rhizopterodendron oppoliense*, Goeppert, 1881, p. 3. (= *Rhizodendron oppoliense*, Goeppert, 1865 n.) Turonian; Silesia.
- Rhodomela diluviana*, Agardh, 1824, p. 201. (= *Fucoides strictus*, Brongniart, 1824.)
- Rhodomelites strictus* (Agardh MS.), Sternberg, 1833, p. 25. (= *Fucoides strictus*, Brongniart, 1824, = *Sphaerococcites strictus*, Brongniart, 1828.) Cretaceous; La Rochelle, France.
- Rhodomelites Mantelli*, Debey in Debey & Ettingshausen, 1859 a, p. 207. (= *Sphaerococcites Mantelli*, Roemer, 1841.) Pläner; Saxony.
- Rhodium Ungerii*, Mercklin, 1855, p. 21, pl. i, figs. 1, 2; pl. ii. Greensand; Russia.
- Rhopala primæva*, Ettingshausen, 1867 a, p. 255, pl. iii, fig. 5. Cenomanian; Niederschoena, Saxony.

- Rhopalophyllum australe**, Ettingshausen, 1893, p. 149; & 1895, p. 28, pl. iii, figs. 12, 13. Cretaceous; Australia.
- **primævum**, Ettingshausen, 1895, p. 2 [nomen nudum]. Ibid.
- Rhus antiqua**, Bozzi, 1891, p. 377, pl. xvi, fig. 4. Cretaceous (?); Italy.
- **cretacea**, Bozzi (*non* Heer, *nec* Velenovsky) in Tommasi, 1892, p. 1119 [nomen nudum]. Senonian (?); Italy.
- **cretacea**, Heer (*non* Velenovsky), 1871 A, p. 14, pl. iii, fig. 11. Senonian; Quedlinburg, Saxony.
- **cretacea**, Velenovsky (*non* Heer), 1887, p. 68, pl. xxvii, figs. 7-12. Cenomanian; Bohemia.
- **dens mortis**, Bayer, 1893, pp. 19, 42, fig. 13; & Bayer in Fritsch, 1893, p. 130, text-fig. 185. Cenomanian; Priesen, Bohemia.
- **evansii**, Lesquereux, 1872, p. 293. Laramie Formation; Wyoming, U.S.A.
- **membranacea**, Lesquereux, 1876 A, p. 369; & 1878 B, p. 292, pl. lxiv, figs. 6, 7. Montana Formation (?); Wyoming, U.S.A.
- **microphylla**, Heer, 1874 A, p. 117, pl. xxxii, fig. 18. (Re-named *Comptonia microphylla* by Berry, 1906 A.) Atane Beds; Greenland
- **Powelliana**, Lesquereux, 1892, p. 155, pl. lvi, figs. 4, 5. Dakota Group; Kansas, U.S.A.
- **pseudo-Meriani**, Lesquereux, 1878 B, p. 293, pl. lviii, fig. 11. Laramie Formation; Wyoming, U.S.A.
- **Uddeni**, Lesquereux, 1892, p. 154, pl. lvii, fig. 2. Dakota Group; Kansas, U.S.A.
- ? **Westii**, Knowlton in Lesquereux, 1892, p. 154, pl. xxxviii, figs. 9, 10. Ibid.
- Rhynchogoniopsis neocomiensis**, Neumann, 1907, p. 87, pl. i, fig. 3. Neocomian; Peru.
- Rhytisma hederæ**, Heer, 1882, p. 20, pl. xxiv, fig. 6 a. Atane Beds; Greenland.
- Rhytismites hederæ**, Meschinelli, 1892, p. 781. (= *Rhytisma hederæ*, Heer, 1882.)
- Roemeria americana**, Unger in Roemer, 1852, p. 95. Cretaceous (?); Texas, U.S.A.
- Rogersia angustifolia**, Fontaine, 1889, p. 288, pl. cxliii, fig. 2; pl. cxliv, figs. 4, 8; pl. cl, figs. 2-7. Potomac Formation; Virginia, U.S.A.
- **angustifolia**, var. **parva**, Fontaine in Ward, 1905, p. 523, pl. cxi, fig. 9. Older Potomac Formation; Washington, U.S.A.
- **longifolia**, Fontaine, 1889, p. 287, pl. cxxxix, fig. 6; pl. cxliv, fig. 2; pl. cl, fig. 1; pl. clx, figs. 1, 2. Potomac Formation; Virginia, U.S.A.
- Rohlfisia celastroides**, Schenk, 1883, p. 9, pl. iv, fig. 12. Upper Cretaceous; Libyan Desert.
- Rosellinites lapideus** (Lesquereux), Knowlton, 1898, p. 204 (re-naming *Sphæria lapidea*, Lesquereux, 1873). Laramie Formation; Colorado, U.S.A.

- Rosthornia carinthiaca*, Unger, 1842, p. 110. Senonian; Carinthia.
- Royena desertorum*, Heer, 1876 b, p. 10, figs. 11-16 (fruits). Upper Cretaceous; Egypt.
- Rubiaephyllum* (*Ericophyllum*) *Gaylussaciae*, Bayer, 1893, pp. 31, 47, text-fig. 21; & Bayer in Fritsch, 1893, p. 131, text-fig. 192. Senonian; Priesen, Bohemia.
- Sabal Campbelli*, Newberry, 1863, p. 515; & 1898, p. 27, pl. xxi, figs. 1, 2. Tertiary & Cretaceous; Washington, U.S.A.
- *grandifolia*, Newberry, 1898, p. 28, pl. xxv; pl. lxiii, fig. 5; pl. lxiv, fig. 2, 2 a. Laramie Formation; Colorado, U.S.A.
- *imperialis*, Dawson, 1883, p. 26, pl. vi, fig. 23. Upper Cretaceous; Vancouver Island, Canada.
- *rigida*, Hatcher, 1901, p. 263, text-fig. 1. Laramie Formation; Wyoming, U.S.A.
- Sabalites andegavensis*, hitherto described as Tertiary, said to be Senonian by Welsch, 1897, p. 667.
- Sabiocaulis Sakuraii*, Stopes & Fujii, 1910, pp. 66-70, pl. viii, fig. pl. ix, figs. 55-57. Upper Cretaceous; Hokkaido, Japan.
- Sagenopteris elliptica*, Fontaine, 1889, p. 149, pl. xxvii, figs. 9, 11-17. Potomac Formation; Virginia, U.S.A.
- *latifolia*, Fontaine, 1889, p. 148, pl. xxvii, fig. 10. Ibid.
- *Mantelli* (Dunker), Schenk, 1871 b, p. 222, pl. xxxi, fig. 5. Recorded Diller & Stanton, 1894, American Lower Cretaceous.
- *neocomiensis*, Hosius & von der Marck, 1880, p. 210, pl. xlv, fig. 194. Neocomian; Westphalia.
- *nervosa*, Fontaine in Ward, 1905, p. 237, pl. lxx, figs. 41-45. Shasta Formation; California, U.S.A.
- *nilssoniana* (Brongniart), Ward, recorded Penhallow, 1902 b, p. 39. Upper Cretaceous; Queen Charlotte Island, Canada.
- *oblongifolia*, Penhallow, 1902 b, p. 40, text-fig. 2. Lower Cretaceous; Queen Charlotte Island, Canada.
- *oregonensis*, Fontaine in Ward, 1905, p. 235, pl. lxx, figs. 36-38 (re-naming *Sagenopteris latifolia*, Fontaine in Diller & Stanton, & *Angiopteridium oregonense*, Fontaine in Stanton.) Shasta Formation; California, U.S.A.
- *variabilis*, Velenovsky, 1889, p. 40. (= *Thinnfeldia variabilis*, Velenovsky, 1885.) Perucer Beds; Bohemia.
- *virginiensis*, Fontaine, 1889, p. 150, pl. cxxxviii, fig. 13; pl. cxxxix, fig. 1. Potomac Formation; Virginia, U.S.A.
- ? *sp.*, Fontaine in Diller & Stanton, 1894, p. 450; & in Ward, 1905, p. 238, pl. lxx, fig. 46. Shasta Formation; California, U.S.A.
- Sagittaria Victor-Masoni*, Ward, 1895 a, p. 354, pl. iii, fig. 5. Potomac Formation; Virginia, U.S.A.
- Salicinium varians*, Hofmann, 1884, p. 191. Danian; Maestricht.
- Salicnoxylon biradiatum*, Lignier, 1907, p. 272, pl. xviii, figs. 18-24; pl. xxiii, figs. 84, 92. Cenomanian; Hève, France.

- Saliciphyllum californicum**, Fontaine in Ward, 1905, p. 266, pl. lxix, fig. 9. Shasta Formation; California, U.S.A.
- **ellipticum**, Fontaine, 1889, p. 303, pl. cxlvi, figs. 2, 4; pl. cl, fig. 8; pl. clxiii, fig. 5; pl. clxvi, fig. 2. Potomac Formation; Virginia, U.S.A.
- **longifolium**, Fontaine, 1889, p. 302, pl. cl, fig. 12. Ibid.
- **pachyphyllum**, Fontaine in Ward, 1905, p. 265, pl. lxix, fig. 8. Shasta Formation; California, U.S.A.
- **parvifolium**, Fontaine, 1889, p. 303, pl. clxxii, fig. 5. Potomac Formation; Virginia, U.S.A.
- **sp.**, Krasser, 1896, p. 129, pl. xiii, fig. 4. Cenomanian; Kunstadt, Moravia.
- Salicites angustus**, Reuss, 1844, p. 169; & 1846, p. 96, pl. li, figs. 7, 8. (= *Grevillea Reussii*, Ettingshausen, 1851.) Cretaceous; Bohemia.
- **fragiliformis**, Zenker (see Brongniart, 1849 A, p. 111). Quadersandstein; Blankenburg, Saxony.
- **Hartigi**, Dunker, 1856, p. 181, pl. xxxiv, fig. 2. Ibid.
- **Petzeldianus**, Goeppert, 1845, p. 220. Cretaceous; Silesia.
- ? **Wahlbergii** (Nilsson), Brongniart, 1849 A, p. 111. Cretaceous; Scania, Sweden.
- Salisburia Baynesiana**, Dawson, 1883, p. 25, pl. v, figs. 21, 21 a. Upper Cretaceous; Vancouver Island, Canada.
- (**Ginkgo**) **lepida**, Dawson, 1886, p. 8, pl. ii, fig. 2. (= *Ginkgo lepida*, Heer, 1876 c.) Kootanie Formation; British Columbia.
- (**Ginkgo**) **nana**, Dawson, 1886, p. 8, pl. ii, fig. 3. (Re-named *Ginkgo nana*, Knowlton, 1898.) Kootanie Formation; Coal Creek, Canada.
- **polymorpha**, Lesquereux, 1859, p. 362; & 1878 B, p. 84, pl. vii, figs. 40, 41. Montana Formation; Missouri R., U.S.A.
- **primordialis** (spelt **Salisburea**), Heer, 1871, p. 1182; & 1874 A, p. 100, pl. xxvii, figs. 1-3. Atane Beds; Greenland.
- **pusilla**, Dawson, 1894, p. 56, pl. vi, figs. 11-14. (Re-named *Ginkgo pusilla*, Knowlton, 1898.) Upper Cretaceous; Vancouver Island, Canada.
- (**Ginkgo**) **sibirica**, Dawson, 1886, p. 8, pl. ii, fig. 1. (= *Ginkgo sibirica*, Heer, 1876 c.) Kootanie Formation; British Columbia.
- **sp.**, Dawson, 1886, p. 18 (nutlets). Lower Laramie Formation; Canada.
- **sp.**, Dawson, 1886, p. 9, pl. ii, fig. 4. Ibid.
- Salix angusta**, Reuss (non A. Braun), 1844, p. 169. Pläner; Bohemia.
- **assimilis**, Saporta, 1894, p. 231, pl. xxxvi, fig. 8 a; pl. xxxvii, figs. 2-3, 6, 13, 19-20. Upper Albian; Portugal.
- **cretacea**, Velenovsky, 1882 A, p. 213 [nomen nudum]. Cretaceous; Bohemia.
- **cuneata**, Newberry, 1870, p. 21; & 1878, pl. ii, fig. 2 (as *Salix Meckii*); & 1898, p. 55, pl. ii, figs. 1, 2. Dakota Group; Nebraska, U.S.A.

- Salix deleta*, Lesquereux, 1892, p. 49, pl. iii, fig. 8.
Dakota Group; Kansas, U.S.A.
- *eutawensis*, Berry, 1910 A, p. 93, pl. xxii, figs. 1-11.
Middle Cretaceous; North Carolina, U.S.A.
- *flexuosa*, Newberry, 1870, p. 21; & 1878, pl. i, fig. 4; & 1898, p. 56, pl. ii, fig. 4; pl. xiii, figs. 3, 4; pl. xiv, fig. 1. (Re-named *S. proteæfolia flexuosa* (Newberry), Lesquereux, 1892.)
Dakota Group; Nebraska, U.S.A.
- *foliosa*, Newberry, 1898, p. 57, pl. xiii, figs. 5, 6.
Cretaceous; New Mexico.
- *fragiliformis*, Zenker, 1833, p. 22, pl. iii, fig. H. (= *Myrica fragiliformis*, Engelhardt, 1892 A, = *Dryandroides Zenkeri*, Ettingshausen, 1867 A.)
Quadersandstein; Blankenburg, Saxony.
- *Goetziana*, Heer, 1871 A, p. 11, pl. iii, figs. 3, 4.
Senonian; Quedlinburg, Saxony.
- *Hayei*, Lesquereux, 1892, p. 48, pl. iii, fig. 7.
Dakota Group; Kansas, U.S.A.
- *inæqualis*, Newberry, 1895, p. 67, pl. xvi, figs. 1, 4, 6; pl. xvii, figs. 2-7.
Amboy Clay; New Jersey, U.S.A.
- *infracretacica*, Saporta, 1894, p. 182, pl. xxxi, fig. 18.
Albian; Portugal.
- *laramiana*, Dawson, 1887, p. 28, pl. i, fig. 10.
Upper Laramie Formation; Canada.
- *Lesquereuxii*, Berry, 1909, p. 252 (re-naming *Salix proteæfolia*, et var. Lesquereux, 1868 & 1896).
Raritan Formation; New Jersey, U.S.A.
- *macrophylla*, Reuss, 1844, p. 169. (= *Proteoides Reussi*, Engelhardt, 1892.)
Cretaceous; Perutz, Bohemia.
- *mattewanensis*, Berry, 1905 B, p. 68, pl. li, fig. 5.
Matawan Formation; New Jersey, U.S.A.
- *Meekii*, Newberry, 1870, p. 19; & 1878, pl. i, fig. 3 (as *S. cuneata*); & 1898, p. 58, pl. ii, fig. 3.
Dakota Group; Nebraska, U.S.A.
- *membranacea*, Newberry, 1870, p. 19; & 1878, pl. i, figs. 5-8 A. (= *S. raritanensis*, Berry, 1909.)
Lower Cretaceous; New Jersey, U.S.A.
- *nervillosa*, Heer in Capellini & Heer, 1867, p. 15, pl. i, fig. 3.
Dakota Group; Nebraska, U.S.A.
- *Newberryana*, Hollick in Newberry, 1895, p. 68, pl. xiv, figs. 2-7.
Baritan Formation; New Jersey, U.S.A.
- *pacifica*, Dawson, 1883, p. 26, pl. vii, fig. 24.
Upper Cretaceous; Vancouver Island, Canada.
- *perucensis*, Velenovsky, 1887, p. 71, pl. xxviii, figs. 1-3.
Perucer Beds; Bohemia.
- *proteæfolia*, Lesquereux, 1868, p. 94; & 1874, p. 60, pl. v, figs. 1-3. (Re-named *Salix Lesquereuxii* by Berry, 1909.)
Dakota Group; Nebraska, U.S.A.
- *proteæfolia flexuosa* (Newberry), Lesquereux, 1892, p. 50, pl. lxiv, figs. 4, 5. (= *S. flexuosa*, Newberry, 1870.)
Ibid.

- Salix proteæfolia lanceolata*, Lesquereux, 1892, p. 50, pl. lxiv, figs. 6-8. Dakota Group; Kansas, U.S.A.
- *proteæfolia linearifolia*, Lesquereux, 1892, p. 49, pl. lxiv, figs. 1-3. Ibid.
- *proteæfolia longifolia*, Lesquereux, 1892, p. 50, pl. lxiv, fig. 9. (Re-named *Salix Lesquereuxii* by Berry, 1909.) Ibid.
- *pseudo-Hayei* (Newberry), Berry, 1909, p. 251 (naming *Salix sp.*, Newberry, 1896, p. 68, pl. xlii, figs. 6-8.) Ibid.
- *purpurcides*, Hollick, 1894 A, p. 50, pl. clxxiv, fig. 9. Raritan Formation; New Jersey, U.S.A. Cretaceous; Long Island, U.S.A.
- *raritanensis* (Newberry), Berry, 1909, p. 250 (re-naming *Salix membranacea*, Newberry, 1868). Raritan Formation; New Jersey, U.S.A.
- *retinenda*, Saporta, 1894, p. 182, pl. xxxi, fig. 19. Albian; Portugal.
- *Schoenæ*, Engelhardt, 1892 A, p. 96, pl. ii, figs. 8, 9. Quader; Freiberg, Saxony.
- *Stantoni*, Knowlton, 1900 A, p. 38, pl. vi, fig. 6. Montana Formation; Utah, U.S.A.
- *Vasseuri*, Marion, 1890, p. 1054 [nomen nudum]. Turonian; Martigues, France.
- ? *Wahlbergii*, Nilsson, 1832, p. 347, pl. i, figs. 5, 6. Greensand; Scania, Sweden.
- *sp.*, Dawson, 1893, p. 57, pl. vii, fig. 22. Upper Cretaceous; Vancouver Island, Canada.
- *sp.*, Hollick, 1893, p. 32, pl. ii, figs. 15, 16. Middle Cretaceous; Staten Island, U.S.A.
- ? *sp.*, Hollick, 1894, p. 63, pl. clxxx, fig. 6. Ibid.
- *sp.*, Knowlton, 1900, p. 39, pl. vii, fig. 2. Montana Formation; Utah, U.S.A.
- *sp.*, Lesquereux, 1892, p. 51, pl. viii, fig. 6 (fruit). Dakota Group; Kansas, U.S.A.
- *sp.*, Roemer, 1889, p. 144, pl. xii, figs. 5-6. Senonian; Silesia.
- Salvertia transylvanica*, Unger, 1865, p. 377, pl. i, fig. 7. Cretaceous; Transylvania.
- Salvinia attenuata*, Lesquereux, 1876 c, p. 377. (= *Marsilea attenuata*, Hollick, 1894 D.) Laramie Formation; Wyoming, U.S.A.
- *elliptica*, Newberry MS. in Hollick, 1894 B, p. 255, pl. ccv, figs. 14, 14 A, 15. Upper Cretaceous (?); Washington, U.S.A.
- Santalum Novæ-Cæsareæ*, Berry, 1906 B, p. 182; & 1906 E, p. 153, pl. xx, fig. 7; pl. xxii, fig. 3. Magothy Formation; New Jersey, U.S.A.
- Sapindophyllum apiculatum*, Velenovsky, 1889, pp. 51, 54, 57. (= *Sapindus apiculatus*, Velenovsky, 1886.)
- *brevior*, Saporta, 1894, p. 205, pl. xxxvii, fig. 11. Upper Albian; Portugal.

- Sapindophyllum coriaceum**, Ettingshausen, 1887 A, p. 187, pl. ix, figs. 22-23. Upper Cretaceous; New Zealand.
- **pelagicum**, Unger in Velenovsky, 1886, p. 7. Cenomanian; Bohemia.
- **subapiculatum**, Saporta, 1894, p. 205, pl. xxxvi, figs. 10-11; pl. xxxvii, figs. 4-5, 10 A, 17. Upper Albion; Portugal.
- Sapindopsis brevifolia**, Fontaine, 1889, p. 300, pl. clvii, fig. 4; pl. clv, figs. 1, 7; pl. clxiii, fig. 3. Potomac Formation; Virginia, U.S.A.
- **cordata**, Fontaine, 1889, p. 296, pl. cxlvii, fig. 1. Ibid.
- **elliptica**, Fontaine, 1889, p. 297, pl. cxlvii, fig. 3. Ibid.
- **magnifolia**, Fontaine, 1889, p. 297, pl. cli, figs. 2, 3; pl. clii, figs. 2, 3; pl. cliii, fig. 2; pl. cliv, figs. 1, 5; pl. clv, fig. 6. Ibid.
- **obtusifolia**, Fontaine, 1889, p. 301, pl. clvi, fig. 13; pl. cliv, figs. 3-6. Ibid.
- **oregonensis**, Fontaine, 1905 A, p. 268, pl. lnx, figs. 15-17. (Renamed by Berry, 1910 D, *Nilssonia oregonensis*.) Shasta Formation; Oregon, U.S.A.
- **parvifolia**, Fontaine, 1889, p. 300, pl. cliv, fig. 6. Potomac Formation; Virginia, U.S.A.
- **tenuinervis**, Fontaine, 1889, p. 301, pl. cliii, fig. 1. Ibid.
- **variabilis**, Fontaine, 1889, p. 298, pl. cli, fig. 1; pl. clii, figs. 1, 4; pl. cliii, fig. 3; pl. cliv, figs. 2-4; pl. clv, figs. 2-5. Ibid.
- Sapindus apiculatus**, Velenovsky, 1886, p. 53, pl. xxii, figs. 1-8. (= *Sapindophyllum apiculatum*, Velenovsky, 1889.) Cenomanian; Bohemia.
- **diversifolius**, Lesquereux, 1892, p. 158, pl. lxiv, fig. 18. Dakota Group; Kansas, U.S.A.
- **imperfectus**, Hollick, 1905 C, p. 415, pl. lxxviii, fig. 4. Cretaceous; Long Island, U.S.A.
- **inexpectans**, Knowlton in Stanton & Hatcher, 1905, p. 144, pl. xvii, fig. 7. Judith River Beds; Montana, U.S.A.
- **Morrisoni**, Lesquereux MS. in Heer, 1882, p. 95, pl. xl, fig. 1; pl. xli, figs. 1 A, 1 B; pl. xliii, figs. 1 A, 1 B; pl. xlv, figs. 7-8; & in Lesquereux, 1883, p. 83, pl. xvi, figs. 1, 2. Dakota Group; Colorado, U.S.A.
- **prodromus**, Heer, 1871, p. 1184; & 1874 A, p. 117, pl. xxxiv, fig. 5. Atane Beds; Greenland.
- **saxonicus**, Engelhardt, 1892 A, p. 101, pl. ii, fig. 14. Quader; Freiberg, Saxony.
- Sapotacites Haydenii**, Heer in Meek & Hayden, 1859, p. 265. Dakota Group; Nebraska, U.S.A.
- **hyperboreus**, Heer, 1883 A, p. 32, pl. lxi, figs. 7-9. Patoot Beds; Greenland.
- **Knowltoni**, Berry, 1906 B, p. 181, pl. viii, fig. 1. (= *Sapotacites* sp. (?), Lesquereux, 1892.) Magothly Formation; Delaware, U.S.A.
- **nervillosus**, Heer, 1883 A, p. 32, pl. lxi, fig. 11. Patoot Beds; Greenland.

- Sapotacites obovata*, Velenovsky, 1882 A, p. 213; & 1886, p. 50, pl. xviii, fig. 6. (= *Sapotophyllum obovatum*, Velenovsky, 1889.)
Cenomanian; Bohemia.
- *retusus*, Heer (*non* Massalongo), 1883 A, p. 32, pl. lxi, fig. 10.
(= *Liriodendropsis retusa* (Heer), Hollick, 1906 A.)
Patoot Beds; Greenland.
- *Stelzneri*, Engelhardt, 1892 A, p. 98, pl. ii, fig. 12.
Quader; Freiberg, Saxony.
- *sp.?*, Lesquereux, 1892, p. 114, pl. lxxv, fig. 3.
Dakota Group; Kansas, U.S.A.
- Sapotophyllum obovatum*, Velenovsky, 1889, pp. 50, 54, 58.
(= *Sapotacites obovata*, Velenovsky, 1882 A.)
Cenomanian; Bohemia.
- Sarcostrobilus Paulini*, Fliche, 1900, pp. 19-23, pl. i, figs. 2-5.
Lower Cretaceous; Haute-Marne, France.
- Sargassites Lyngbianus* (Brongniart), Sternberg, 1833, p. 36.
(= *Fucoides Lyngbianus*, Brongniart, 1828.) Cretaceous; Arnager.
- *Rosthorni*, Sternberg, 1833, p. 36, pl. xxv, fig. 6.
Lower Cretaceous; Carinthia.
- Sassafras acutilobum*, Lesquereux, 1874, p. 79, pl. xiv, figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- *angustilobum*, Hollick, 1906 A, p. 77, pl. xxix, figs. 1-3.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- *arctica*, Heer, 1871, p. 1183; & 1874 A, p. 109, pl. xxxi, figs. 3 a, 3 b.
Atane Beds; Greenland.
- *bilobatum*, Fontaine, 1889, p. 290, pl. clvi, fig. 12; pl. clxiv, fig. 4.
Potomac Formation; Virginia, U.S.A.
- *cretaceum*, Newberry, 1870, p. 14; & 1878, pl. vi, figs. 1-4; & 1898, p. 98, pl. vi, figs. 1-4; pl. vii, figs. 1-3; pl. viii, figs. 1, 2.
Dakota Group; Nebraska, U.S.A.
- *cretaceum dentatum*, Lesquereux, 1876, p. 344; & 1874, pl. xi, figs. 1, 2.
Dakota Group; Kansas, U.S.A.
- *cretaceum grossedentatum*, Lesquereux, 1892, p. 101, pl. li, fig. 5.
Ibid.
- *cretaceum heterolobum*, Fontaine, 1889, p. 289, pl. clii, fig. 5; pl. elix, fig. 8; pl. clxiv, fig. 5.
Potomac Formation; Virginia, U.S.A.
- *cretaceum obtusum*, Lesquereux, 1874, p. 80, pl. xii, fig. 3; pl. xiii, fig. 1.
Dakota Group; Kansas, U.S.A.
- *cretaceum recurvatum* (Lesquereux), Newberry, 1898, p. 99, pl. ix, fig. 2. (= *S. recurvatum*, Lesquereux, 1873.)
Ibid.
- (*Araliopsis*) *dissectum*, Lesquereux, 1883, p. 57; & 1892, p. 101, pl. xiv, fig. 1.
Ibid.
- (*Araliopsis*) *dissectum symmetricum*, Hollick, 1895, p. 226, pl. ccxxxvi.
Ibid.
- *harkeriana* (*sic*), Lesquereux, 1873, p. 425; & 1874, p. 81, pl. xi, figs. 3-4; pl. xxvii, fig. 2.
Ibid.

- Sassafras hastatum*, Newberry, 1895, p. 88, pl. xxvii, figs. 4-6;
pl. xxviii, figs. 1, 2; pl. xl, fig. 4. Amboy Clay; Woodbridge, U.S.A.
- *Krejci*, Velenovsky, 1882 A, p. 213 [nomen nudum].
Cretaceous; Bohemia.
- *Leconteanum*, Lesquereux, 1869, p. 431, pl. xxiii, fig. 1. (= *Persea*
Leconteana, Lesquereux, 1874.) Dakota Group; Nebraska, U.S.A.
- (*Araliopsis*) *mirabilis*, Lesquereux, 1873, p. 424; & 1874, p. 80,
pl. xii, fig. 1. Dakota Group; Kansas, U.S.A.
- *Mudgii*, Lesquereux, 1868, p. 99; & 1874, p. 78, pl. xiv, figs. 3, 4;
pl. xxx, fig. 7. Dakota Group; Nebraska, U.S.A.
- *obtusum*, Lesquereux, 1872, p. 303; & 1874, p. 81, pl. xiii,
figs. 2-4. Ibid.
- (*Araliopsis*) *papillosum*, Lesquereux, 1892, p. 102, pl. vi, fig. 7,
Dakota Group; Kansas, U.S.A.
- *parvifolium*, Fontaine, 1889, p. 289, pl. cxxxix, fig. 7.
Potomac Formation; Maryland, U.S.A.
- *Paulini*, Fliche, 1900, p. 15, pl. i, figs. 2-5. Neocomian; France.
- *Pfaffiana*, Heer, 1883 A, p. 29, pl. lv, fig. 18.
Patoot Beds; Greenland.
- (*Araliopsis*) *platanoides*, Lesquereux, 1883, p. 58, pl. vii, fig. 1.
Dakota Group; Kansas, U.S.A.
- ? *primordiale*, Lesquereux, 1892, p. 100, pl. xvi, fig. 10. Ibid.
- *progenitor*, Newberry MS. (?) in Hollick, 1894 A, p. 53, pl. clxxiv,
fig. 1; & Newberry, 1895, p. 88, pl. xxvii, figs. 1-3.
Cretaceous; Long Island, U.S.A.
- *protophyllum*, Saporta, 1894, p. 182, pl. xxxi, fig. 17.
Albian; Portugal.
- (*Araliopsis*) *recurvatum*, Lesquereux, 1873, p. 424. (= *Platanus*
recurvata, Lesquereux, 1874, and *S. cretaceum recurvatum* (Les-
quereux), Newberry, 1898.) Dakota Group; Kansas, U.S.A.
- (*Araliopsis*) *rotundilobum*, Newberry in Ward, 1895 A, p. 380.
Potomac Formation; Virginia, U.S.A.
- *subintegrifolium*, Lesquereux, 1868, p. 99; & 1874, p. 82, pl. iii,
fig. 5. Dakota Group; Nebraska, U.S.A.
- *sp.*, Dawson, 1883, p. 27, pl. vii, figs. 30, 30 a.
Upper Cretaceous; Vancouver Island, Canada.
- Saururopsis niponensis*, Stopes & Fujii, 1910, pp. 58-62, pl. vii,
figs. 42-47; text-fig. 19. Upper Cretaceous; Hokkaido, Japan.
- Schizaepteris mesozoica*, Stopes & Fujii, 1910, pp. 6-10, text-figs.
1-3, pl. ii, phot. 1. Ibid.
- Schizoneuropsis posthuma*, Richter, 1906, p. 13, pl. vi, fig. 10.
Senonian; Quedlinburg, Saxony.
- Sclerophyllina cretosa* (Schenk), Heer, 1874 A, p. 59, pl. xiii, figs. 13,
14; pl. xvii, fig. 12; & p. 124, pl. xxxv, figs. 8-10. (= *Baiera*
cretosa, Schenk, 1871 A.) Kome Beds; Greenland.
- *cretosa major*, Heer, 1874 A, p. 124, pl. xxxv, figs. 9-10.
Cretaceous; Spitzbergen.

- Sclerophyllina dichotoma*, Heer, 1868, p. 82, pl. xlv, fig. 6.
Kome Beds; Greenland.
- Scleropteris bellidula*, Heer, 1874 A, p. 35, pl. ii, figs. 17 c, 17 d, 18;
pl. xi, fig. 8. Ibid.
- *callosa*, Schulze, 1888, p. 15 [nomen nudum]. Senonian; Salzberg.
- *debilior*, Saporta, 1894, p. 107, pl. xix, fig. 14.
Neocomian (?); Portugal.
- *dentata*, Fontaine, 1889, p. 153, pl. lxiii, figs. 3, 4.
Potomac Formation; Virginia, U.S.A.
- *distantifolia*, Fontaine in Ward, 1899 B, p. 662, pl. clxii, figs. 2, 3.
Lower Cretaceous; Black Hills, U.S.A.
- *elliptica*, Fontaine, 1889, p. 151, pl. xxviii, figs. 2, 4, 6; pl. xxix,
fig. 1. Potomac Formation; Virginia, U.S.A.
- *elliptica*, var. *longifolia*, Fontaine, 1889, p. 152, pl. xxviii, fig. 7.
Ibid.
- *rotundifolia*, Fontaine in Ward, 1899 B, p. 663, pl. clxii, figs. 4, 5.
Lower Cretaceous; Black Hills, U.S.A.
- *vernonensis*, Ward, 1895 A, p. 349, pl. ii, figs. 1-3.
Potomac Formation; Virginia, U.S.A.
- *virginica*, Fontaine, 1889, p. 152, pl. xxviii, figs. 3, 5. Ibid.
- Sclerotites* sp., Geinitz, 1842, p. 99, pl. xxiv, figs. 1-3.
Cenomanian; Saxony.
- sp. (Lx.), Meschinelli, 1895, p. 659. (= *Sclerotium* (?) sp., Les-
quereux.)
- Sclerotium* ? sp., Lesquereux, 1892, p. 23, pl. lix, figs. 4, 4 a.
Dakota Group; Kansas, U.S.A.
- Scolopendrium officinarum dædaleum*, Cotta, 1836, p. 586.
Cenomanian; Niederschoena, Saxony.
- Sedites* (?) *Rabenhorstii*, Geinitz, 1842, p. 97, pl. xxiv, fig. 5.
Pläner; Strehlen, Dresden.
- Selaginella arctica*, Heer, 1882, p. 39, pl. xiii, fig. 5.
Atane Beds; Greenland.
- *dichotoma*, Velenovsky, 1888 B, p. 29, pl. vi, figs. 8-11.
Perucer Beds; Vyserovic, Bohemia.
- ? *falcata*, Lesquereux, 1876 c, p. 365; & 1878 B, p. 48, pl. lxi,
figs. 12-15. Montana Formation; Wyoming, U.S.A.
- *laciniata*, Lesquereux, 1876 c, p. 378; & 1878 B, p. 47, pl. lxiv,
figs. 12, 12 a. Ibid.
- *marylandica*, Fontaine in Ward, 1905, p. 553, pl. cxv, figs. 9, 10.
Older Potomac Formation; Maryland, U.S.A.
- Sequoia acuminata*, Lesquereux, 1876 c, p. 384.
Laramie Formation; Wyoming, U.S.A.
- *acutifolia*, Newberry, 1891, p. 200, pl. xiv, figs. 7, 7 a.
Kootanie Formation; Montana, U.S.A.
- *albertensis*, Penhallow, 1908, p. 83, figs. 1-6.
Judith River Series; Alberta, Canada.
- *ambigua*, Heer, 1874 A, p. 78, pl. xxi, figs. 1, 2 a, 3-8, 9 a, 10 a,
b, c, 11. Kome Beds; Greenland.

- Sequoia biformis**, Lesquereux, 1876 c, p. 366; & 1878 b, p. 80, pl. lxii, figs. 15-18. (Re-named *Geinitzia biformis*, Knowlton, 1900 a.)
Montana Formation; Wyoming, U.S.A.
- **brevifolia**, Heer, 1868, p. 93, pl. ii, fig. 23. Ibid.
- **concinna**, Heer, 1883 a, p. 13, pl. xlix, figs. 8 b, 8 c; pl. l, fig. 1 b; pl. li, figs. 2-10; pl. lii, figs. 1-3; pl. liii, fig. 1 b.
Patoot Beds; Greenland.
- **condita**, Lesquereux, 1876 a, p. 391; & 1876 b, p. 335, pl. iv, figs. 5-7. Dakota Group; Kansas, U.S.A.
- **Couttsiae**, Heer, 1863, p. 1051, pls. lix, lx, lxi. Tertiary.
(Recorded American Cretaceous, Hollick, 1892, p. 30, pl. i, fig. 5.)
- **crispa**, Velenovsky, 1885, p. 22, pl. x, figs. 5-7, 9, 14, 16.
Perucer Beds; Bohemia.
- **cuneata**, Newberry, 1898, p. 18, pl. xiv, figs. 3-4 a.
Upper Cretaceous; Canada.
- **cycadopsis**, Fontaine, 1889, p. 243, pl. cxii, figs. 9-11; pl. cxiii, figs. 1-3. Potomac Formation; Virginia, U.S.A.
- **delicatula**, Fontaine, 1889, p. 247, pl. cxxi, fig. 3. Ibid.
- **densifolia**, Fontaine, 1889, p. 246, pl. cxxi, fig. 4. Ibid.
- **Fairbanksi**, Fontaine, 1905, p. 178, pl. xlvi, figs. 9-11.
Lower Cretaceous; California, U.S.A.
- **fastigiata** (Sternberg), Heer, 1869 a, p. 11, pl. i, figs. 10-13.
(= *Caulerpites fastigiatus*, Sternberg, 1833.)
Cenomanian; Moleteia, Moravia.
- **formosa**, Lesquereux, 1868, p. 92; & 1874, p. 50, pl. i, fig. 9.
Dakota Group; Nebraska, U.S.A.
- **Goepperti** (Dunker), Schulze, 1888, p. 19 [nomen nudum].
Senonian; Altenburg.
- **gracilis**, Heer, 1871, p. 1181; & 1874 a, p. 80, pl. xviii, fig. 1 c; pl. xxii, figs. 1-10. Kome Beds; Greenland.
- **gracilis laxa**, Heer, 1874 a, p. 82, pl. xxii, fig. 10. Ibid.
- **gracillima** (Lx.), Newberry, 1895, p. 50, pl. ix, figs. 1-3.
Amboy Clay; New Jersey, U.S.A.
- **heterophylla**, Velenovsky, 1885, p. 22, pl. xiii, figs. 2-4, 6-9; pl. xiii, fig. 12. Perucer Beds; Bohemia.
- **? inferna**, Ward, 1905, p. 507 (naming *Sequoia* sp. (?), Fontaine, 1889, p. 248, pl. cxvi, fig. 7; pl. cxxxii, figs. 2, 5, 6).
Lower Potomac Formation; Virginia, U.S.A.
- **intermedia**, Richter, 1904, p. 15, pl. i, fig. 8; & 1905, p. 5, pl. i, fig. 8. Senonian; Saxony.
- **Langsdorffii** (Brongn.), Heer, 1883 a, p. 15, pl. liii, fig. 8.
Patoot Beds; Greenland.
- **legdensis**, Hosijs & von der Marck, 1880, p. 180, pl. xxxvii, fig. 147. Lower Senonian; Westphalia.
- **lepidota**, Bayer in Fritsch, 1893, p. 129, text-fig. 180; & Bayer, 1893, pp. 8, 37, figs. 6, 7, 8. Senonian; Priesen, Bohemia.

- Sequoia longifolia*, Lesquereux, 1876 c, p. 365; & 1878 b, p. 79, pl. vii, figs. 14, 14 a; pl. lxi, figs. 28, 29. (Re-named *Geinitzia longifolia*, Knowlton, 1900 A.) Montana Formation; Wyoming, U.S.A.
- *lusitanica*, Heer, 1881, p. 18, pl. xvii, figs. 7 b, c, 7 bb, 9. Cretaceous; Portugal.
- *macrolepis*, Heer, 1883 A, p. 16, pl. li, figs. 1, 12 b. Patoot Beds; Greenland.
- *major*, Velenovsky, 1888 A, p. 594, figs. 4-6 on unnumb. pl. Cenomanian; Bohemia.
- *microcarpa*, Velenovsky, 1885, p. 24, pl. x, figs. 2, 3. Ibid.
- *minor*, Velenovsky, 1887 A, p. 638, figs. 11, 12 on plate. Ibid.
- *moravica*, Krasser & Kubart, 1906 b, p. 46 [nomen nudum]. Cenomanian; Moletein, Moravia.
- *oblonga*, Marik, 1901, p. 8, pl. i, fig. 21. Cenomanian; Bohemia.
- *pectinata*, Heer, 1871 A, p. 8, pl. i, fig. 8. Senonian; Quedlinburg, Saxony.
- *pogiophylloides*, Fontaine, 1894, p. 276, pl. xlii, figs. 1-3 a. Trinity Division; Texas, U.S.A.
- *Reichenbachi* (Geinitz), Heer, 1868, p. 83, pl. xliii, figs. 1 d, 2 b, 5 a, 5 d, 5 dd, 8, 8 b; & Geinitz, 1875 A, p. 306, pl. lxvii, fig. 6. (= *Araucarites Reichenbachi*, Geinitz, 1842, = *Geinitzia Reichenbachi*, Hollick & Jeffrey, 1909.) Kome Beds; Greenland.
- *Reichenbachi longifolia*, Fontaine, 1889, p. 244, pl. cxvii, fig. 8. Potomac Formation; Virginia, U.S.A.
- *Reichenbachi Rabenhorsti* (Geinitz), Heer, 1871, p. 1181 [nomen nudum]. Kome Beds; Greenland.
- *rigida*, Heer, 1871, p. 1182; & 1874 A, p. 80, pl. xxii, figs. 5 g, 11 a; pl. xxv, fig. 6. Atane Beds & Kome Beds; Greenland.
- *Smittiana*, Heer, 1871, p. 1181; & 1874 A, p. 82, pl. xii, fig. 10 b; pl. xvii, figs. 3, 4; pl. xviii, fig. 1 b; pl. xx, figs. 5 b, 7 c; pl. xxiii, figs. 1-6. Ibid.
- *subulata*, Heer, 1871, p. 1182; & 1874 A, p. 102, pl. xxvii, figs. 3 b, 7, 8 b, 15 a; pl. xxviii, figs. 3-6; pl. xxix, figs. 2 c, 7 b; pl. xxxiv, fig. 1 a. Atane Beds; Greenland.
- *subulata lusitanica*, Saporta, 1894, p. 177, pl. xxxiii, figs. 7-12. Albian; Portugal.
- *Winchellii*, Lesquereux in Winchell, 1885, p. 76; & Lesquereux, 1895, p. 10, pl. A, fig. 1. Dakota Group; Minnesota, U.S.A.
- *Woodwardii* (Carr.), Schimper, 1872, p. 316. (= *Sequoiites Woodwardii*, Carruthers, 1866 B.) Upper Greensand; Dorset.
- *sp.*, Fontaine, 1889, p. 247, pl. cxx, fig. 9. Potomac Formation; Virginia, U.S.A.
- *sp.*, Fontaine in Ward, 1899 b, p. 676, pl. clxvi, figs. 3, 4. Lowest Cretaceous; Wyoming, U.S.A.
- *sp.*, Fontaine, 1889, p. 248, pl. cxvi, fig. 7; pl. cxxii, figs. 2, 5, 6. (= *Sequoia? inferna*, Ward, 1905.) Potomac Formation; Virginia, U.S.A.

- Sequoia* sp., Hollick, 1905 c, p. 410, pl. lxxii, fig. 2.
Middle Cretaceous; Long Island, U.S.A.
- sp., Hollick, 1906 A, p. 44, pl. ii, fig. 42 (cone).¹
Middle Cretaceous; Martha's Vineyard, U.S.A.
- sp.?, Knowlton in Stanton & Hatcher, 1905, p. 131, pl. xiv, fig. 2
(cone). Judith River Beds; Alberta, Canada.
- sp.?, Newberry, 1895, p. 49, pl. ix, figs. 4, 4 a.
Amboy Clay; New Jersey, U.S.A.
- Sequoiites Gardneri*, Carruthers, 1869, p. 7, pl. i, figs. 7, 8.
Gault; Folkestone.
- *Holsti*, Nathorst in Conwentz, 1892, p. 28, pl. iii, figs. 4-5; pl. iv,
figs. 1-4; pl. viii, figs. 2-7. Senonian; Sweden.
- *ovalis*, Carruthers, 1871, p. 541, text-fig. Gault; Folkestone.
- *polyanthes*, Marik, 1901, p. 9, pl. ii, fig. 1.
Cenomanian; Bohemia.
- *Woodwardii*, Carruthers, 1866 B, p. 544, pl. xxi, figs. 11-16.
Upper Greensand; Blackdown.
- Sequoiopsis speciosa*, Richter, 1899 B, p. 44.
Senonian; Quedlinburg, Saxony.
- Serenopsis Kempii*, Hollick, 1893 B, p. 169, pl. cxlix. (= *Nelumbo*
Kempii, Hollick, 1905 c.) Formation (?); Long Island, U.S.A.
- Sillmania texana*, Unger, 1850 A, p. 524; & in Roemer, 1852, p. 95.
Cretaceous (?); Texas, U.S.A.
- Simaba ? saxonica*, Engelhardt, 1892 A, p. 102, pl. ii, fig. 6.
Quader; Freiberg, Saxony.
- Smilax grandifolia-cretacea*, Lesquereux, 1892, p. 40, pl. xlvi, fig. 3.
Dakota Group; Kansas, U.S.A.
- *panartia*, Bayer, 1896, pp. 8, 32, fig. 5.
Upper Senonian; Kieslingswalda, Bohemia.
- *raritanensis*, Berry, 1909, p. 248. Raritan Formation; U.S.A.
- *undulata*, Lesquereux, 1892, p. 39, pl. xlvi, fig. 2.
Dakota Group; Kansas, U.S.A.
- Solenostelopteris japonica*, Kershaw, 1910, p. 689, pl. lviii & text-
fig. Upper Cretaceous; Hokkaido, Japan.
- Sparganium cretaceum*, Heer, 1874 A, p. 105, pl. xxviii, fig. 12.
Atane Beds; Greenland.
- Spathites* sp., Knowlton, 1897, p. 140.
Montana Formation; Wyoming, U.S.A.
- Sphaeria cretacea*, Heer, 1883 A, p. 1, pl. ix, fig. 2. (= *Spherites*
cretaceus, Meschinelli, 1892.) Patoot Beds; Greenland.
- *lapidea*, Lesquereux, 1873, p. 373. (= *Spherites lapideus*, Meschi-
nelli, 1892, & *Rosellinites lapideus*, Knowlton, 1898.)
Laramie Formation; New Mexico, U.S.A.
- *myricæ*, Lesquereux, 1873, p. 390. (= *Spherites myricæ*, Meschi-
nelli, 1892.) Laramie Formation; Wyoming, U.S.A.
- *phyllostichoides*, Saporta, 1894, p. 156, pl. xxviii, fig. 1.
Albian; Portugal.

- Sphæria problematica**, Knowlton in Lesquereux, 1892, p. 23, pl. xxxi, figs. 2, 2 a. (= *Sphærites problematicus*, Meschinelli, 1895.)
Dakota Group; Kansas, U.S.A.
- **rhytismoides**, Lesquereux, 1876 A, p. 382.
Laramie Formation; Wyoming, U.S.A.
- Sphærites cretaceus** (Heer), Meschinelli, 1892, p. 753. (= *Sphæria cretacea*, Heer, 1883 A.) Patoot Beds; Greenland.
- **lapideus** (Lesquereux), Meschinelli, 1892, p. 766. (= *Sphæria lapidea*, Lesquereux, 1873.)
- **myricæ** (Lesquereux), Meschinelli, 1892, p. 759. (= *Sphæria myricæ*, Lesquereux, 1873.) Laramie Formation; Wyoming, U.S.A.
- **primævus**, Goepfert, 1836, p. 87 [nomen nudum].
- **problematicus**, Meschinelli, 1895, p. 657. (= *Sphæria problematica*, Knowlton, 1892.) Dakota Group; Kansas, U.S.A.
- **solitarius**, Debey & Ettingshausen, 1859 A, p. 213, pl. iii, figs. 4 e-g. Senonian; Aix, Rhenish Prussia.
- Sphærococcites centralis**, Goepfert, 1851, p. 46; & 1854, p. 229, pl. iii. Quader; Westphalia.
- **cornutus**, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **Laubei**, Engelhardt, 1892, p. 91, pl. i, fig. 1.
Perucer Beds; Bohemia.
- **lesinensis**, Unger, 1869, p. 56, pl. v, fig. 1.
Cretaceous; Dalmatia.
- **lichenoides**, Goepfert, 1865, p. 642 [nomen nudum].
Quader; Germany.
- **Mantelli**, Roemer, 1841, p. 1, pl. i, fig. 2. (= *Rhodomenites Mantelli*, Debey, = *Chondrites Mantelli*, Geinitz, 1850 A.)
Pläner; Saxony.
- **Meyrati**, Fischer-Ooster, 1858, p. 56, pl. iv, fig. 4.
Neocomian; Switzerland.
- **Mohli**, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **pinnatifidus**, Unger, 1850 A, p. 27; & 1853, p. 80, pl. xxiv, fig. 7.
Cretaceous; Bavaria.
- **strictus**, Agardh MS. in Brongniart, 1828, p. 52. (= *Rhodomenites strictus*, Sternberg.) Cretaceous; La Rochele, France.
- **striolatus**, Presl in Sternberg, 1825, p. 105, pl. xxvii, fig. 3; pl. lxxv, figs. 32, 33. Tertiary. (Recorded Otto, 1852 ?, p. 14, pl. iv, fig. 1.)
Lower Quader; Dippoldiswalde
- Sphenaspis statenensis**, Hollick & Jeffrey, 1909, p. 51, pl. x, figs. 22, 23; pl. xxvi, figs. 2-4. Raritan Formation; Staten Island, U.S.A.
- Sphenolepidium debile**, Heer, 1881, p. 20, pl. iii, figs. 20, 21, 20 b.
Cretaceous; Portugal.
- **dentifolium**, Fontaine, 1889, p. 258, pl. cxxvii, figs. 3, 4; pl. cxxviii, figs. 2-6; pl. cxxix, fig. 5; pl. cxxx, figs. 4-6, 10.
Potomac Formation; Virginia, U.S.A.

- Sphenolepidium Kurrianum** (Dunker), Heer, 1881, p. 19, pl. xiii, figs. 1*b*, 8*b*; pl. xviii, figs. 1-8. (= *Thuites Kurrianus*, Dunker, 1846.)
Cretaceous; Almagem, Portugal.
- **pachyphyllum**, Fontaine, 1889, p. 259, pl. cxxi, figs. 6, 7.
Potomac Formation; Virginia, U.S.A.
- **parceramosum**, Fontaine, 1889, p. 257, pl. cxxix, fig. 7; pl. cxxx, fig. 8; pl. cxxvi, fig. 2. Ibid.
- **recurvifolium**, Fontaine, 1889, p. 258, pl. cxxvii, fig. 2; pl. cxxx, figs. 2, 7. Ibid.
- **Sternbergianum** (Dunker), Heer, 1881, p. 19, pl. xiii, figs. 2-8, 1*a*; pl. xiv. (= *Sphenolepis Sternbergiana* (Dunker), Schenk, 1871.)
Lower Cretaceous; Portugal.
- **Sternbergianum**, var. **densifolium**, Fontaine, 1889, p. 261, pl. cxviii, fig. 7; pl. cxxi, figs. 5, 7, 9; pl. cxxv, fig. 2; pl. cxxix, fig. 3; pl. cxxx, fig. 1; pl. cxxxi, figs. 1-3; pl. cxxxii, fig. 4.
Potomac Formation; Virginia, U.S.A.
- **virginicum**, Fontaine, 1889, p. 259, pl. cxxv, fig. 4; pl. clxvi, fig. 6. Ibid.
- **sp.**, Dawson, 1893, p. 90, text-fig. 13.
Kootanie Formation; North-West Territory, Canada.
- Sphenolepis Kurriana**, Schenk, 1871*b*, p. 243, pl. xxvii, figs. 5-8; pl. xxxviii, fig. 1. (= *Sphenolepidium Kurrianum* (Dunker), Heer, 1881.)
Wernsdorfer Beds; Carpathians.
- **Sternbergiana** (Dunker), Schenk, 1871*b*, p. 243, pl. xxvii, figs. 3, 4; pl. xxxviii, figs. 3-13. (= *Sphenolepidium Sternbergianum* (Dunker), Heer, 1881.)
Wealden; Germany.
- Sphenopteridium tenerum**, Marik, 1901, p. 7, pl. i, fig. 17.
Cenomanian; Bohemia.
- Sphenopteris acrodentata**, Fontaine, 1889, p. 90, pl. xxxiv, fig. 4.
Potomac Formation; Virginia, U.S.A.
- **acutidens**, Saporta, 1894, p. 128, pl. xxiii, fig. 4.
Urgonian; Portugal.
- **aneimiaeformis**, Saporta, 1894, p. 128, pl. xxii, fig. 3*a*. Ibid.
- **angustiloba**, Heer, 1881, p. 14, pl. xvi, figs. 1, 2, 3.
Cretaceous; Portugal.
- **asplenifolia**, Feistmantel, 1874, p. 267.
Perucer Beds; Bohemia.
- **capillaris**, Saporta, 1894, p. 74, pl. xvii, figs. 19-21.
Valanginian; Portugal.
- **cercalensis**, Saporta, 1894, p. 126, pl. xxiv, figs. 4-6; pl. xxv, fig. 20.
Urgonian; Portugal.
- **Choffatiana infracretacica**, Saporta, 1894, p. 67, pl. xvii, figs. 11-13.
Neocomian; Portugal.
- **corrugata**, Newberry, 1870, p. 10; & 1878, pl. ii, fig. 6.
Dakota Group; Nebraska, U.S.A.
- **crenularis** Saporta 1894, p. 160, pl. xxix, fig. 13; pl. xxx, fig. 12.
Albian; Portugal.

- Sphenopteris cuneifida**, Saporta in Choffat, 1889, p. 200; & Saporta, 1894, p. 69, pl. xvi, fig. 11, & p. 127, pl. xxiii, fig. 5.
Valanginian & Urgonian; Portugal.
- **debiliformis**, Saporta, 1894, p. 67, pl. xviii, fig. 15.
Neocomian; Portugal.
- **debilior**, Saporta, 1894, p. 161, pl. xxviii, fig. 5. Albian; Portugal.
- **debilis**, Heer (*non* (Sternberg), Unger), 1871, p. 1181 [nomen nudum].
Kome Beds; Greenland.
- **dissectiformis**, Saporta, 1894, p. 68, pl. xv, fig. 18; pl. xvi, figs. 22-23.
Valanginian; Portugal.
- **Drygalskii**, Engelhardt in Vauhöffen, 1897, pp. 363, 371, text-fig. 27.
Kome Beds; Greenland.
- **elongata**, Newberry, 1863, p. 511.
Laramie Formation (?); Washington, U.S.A.
- **flabellina**, Saporta, 1894, p. 160, pl. xxviii, figs. 3, 6; pl. xxix, fig. 16.
Albian; Portugal.
- **flabellinervia**, Saporta, 1894, p. 70, pl. xv, fig. 20.
Valanginian; Portugal.
- **flabellisecta**, Saporta, 1894, p. 69, pl. xv, figs. 14-15. Ibid.
- **fragilis**, Heer (*non* (Schlotheim), Brongniart), 1874 A, p. 34, pl. ii, figs. 20, 20 b.
Kome Beds; Greenland.
- **ginkgoides**, Saporta, 1894, p. 68, pl. xv, fig. 13.
Valanginian; Portugal.
- **Goepperti**, Dunker. Recorded Saporta, 1894, p. 157.
Albian; Portugal.
- **gomesiana**, Heer, 1881, p. 13, pl. xi, fig. 7. Cretaceous; Portugal.
- **grevillioides**, Heer, 1874 A, p. 34, pl. xi, figs. 10, 11. (= *Thyrsopteris grevillioides*, Hollick, 1906 A.)
Kome Beds; Greenland.
- **hyperborea**, Heer, 1874 A, p. 123, pl. xxxviii, figs. 1 b, 2 b, 9 b, c.
Cretaceous; Spitzbergen.
- **invovens**, Saporta, 1894, p. 159, pl. xxviii, fig. 10; pl. xxiv, figs. 2-5, 7; pl. xxv, figs. 10-11; pl. xxxi, figs. 3, 5; pl. xxxv, fig. 6.
Albian; Portugal.
- **Johnstrupi**, Heer, 1868, p. 78, pl. xliii, figs. 7, 7 b. (= *Asplenium Johnstrupi*, Heer, 1874 A.)
Kome Beds; Greenland.
- **latiloba**, Fontaine, 1889, p. 90, pl. xxxv, figs. 3-5; pl. xxxvi, figs. 4-9; pl. xxxvii, fig. 1. Potomac Formation; Virginia, U.S.A.
- (*Asplenium*) **lepida**, Heer, 1871, p. 1180 (= *Jeandaulia lepida*, Heer, 1874 A.)
Kome Beds; Greenland.
- **lesinensis**, Kerner, 1896, p. 37, pl. i, figs. 1-5; pl. v, fig. 9.
Cenomanian; Lesina, Dalmatia.
- **linearisecta**, Saporta, 1894, p. 125, pl. xxiii, figs. 10, 11; pl. xxv, fig. 19.
Urgonian; Portugal.
- **lobulifera**, Saporta, 1894, p. 71, pl. xv, fig. 16; & p. 124, pl. xxiii, figs. 6-7.
Valanginian & Urgonian; Portugal.
- **longifolia**, Feistmantel (*non* Dunker), 1874, p. 267.
Perucer Beds; Bohemia.

- Sphenopteris lupulina**, Heer, 1881, p. 15, pl. xi, figs. 8, 8*b-c*.
Cretaceous; Portugal.
- **Mantelli**, Brongniart, 1828, p. 170, pl. xlv, figs. 3-7. (= *Onychiopsis Mantelli* (Brongniart), Seward, 1894, = *Hymenopteris psilotoides*, Stokes & Webb, 1824.)
Hastings Sands; Sussex.
- **Morrisiana**, Johnston, 1896, p. 58, figs. 14, 15.
Cretaceous (?); Tasmania.
- **Nordenskiöldi**, Heer, 1871, p. 1181 [nomen nudum].
Kome Beds; Greenland.
- **pachyphylla**, Fontaine, 1889, p. 93, pl. I, fig. 5.
Potomac Formation; Virginia, U.S.A.
- **Pichleri**, Schenk, 1876, p. 166, pl. xxix, figs. 2-5.
Upper Cretaceous; Tyrol.
- **plurinerva**, Heer, 1881, p. 13, pl. xi, figs. 6 & 6*b*; pl. xv, figs. 8, 8*b-c*.
Aptian-Neocomian; Portugal.
- **polyclada**, Saporta, 1894, p. 127, pl. xxiii, fig. 12.
Urgonian; Portugal.
- **pseudo-Cordai**, Saporta, 1894, p. 163, pl. xxviii, fig. 11; pl. xxx, figs. 3-8; pl. xxxiii, fig. 4*a*.
Albian; Portugal.
- **pseudolepida**, Saporta, 1894, p. 73, pl. xv, fig. 32.
Valanginian; Portugal.
- **pygmæa**, Saporta, 1894, p. 127, pl. xxiv, fig. 7*a*.
Urgonian; Portugal.
- **recurrens**, Saporta, 1894, p. 162, pl. xxix, fig. 15.
Albian; Portugal.
- **spatulata**, Fontaine, 1889, p. 93, pl. I, fig. 4.
Potomac Formation; Virginia, U.S.A.
- **subadnata**, Feistmantel, 1874, p. 267. Peruczer Beds; Bohemia.
- **subtilinervis**, Saporta, 1894, p. 19, pl. iii, fig. 32; pl. xiv, figs. 12, 17; p. 68, pl. xv, fig. 22; pl. xvi, fig. 4. Valanginian; Portugal.
- **tasmanica**, Johnston, 1896, p. 60, figs. 10-13.
Cretaceous (?); Tasmania.
- **tenuicula**, Yokoyama, 1894, p. 217, pl. xx, fig. 11; pl. xxi, figs. 2, 2*a*; pl. xxviii, fig. 6. Neocomian; Japan.
- **tenuifissa**, Saporta, 1894, p. 161, pl. xxviii, fig. 4.
Albian; Portugal.
- **thrysopteroides**, Fontaine, 1889, p. 89, pl. xxv, fig. 3; pl. lviii, fig. 5.
Potomac Formation; Virginia, U.S.A.
- **valdensis**, Heer, 1881, p. 14, pl. xv, figs. 9-14; pl. xvi, fig. 5*b*.
Lower Cretaceous; Portugal.
- Sphenozamites** sp., Dawson, 1886, p. 7.
Kootanie Formation; Martin Creek, Canada.
- Spiropteris** sp., Nathorst, 1891, p. 34, pl. i, figs. 15-17.
Upper Cretaceous (?); Königs-Wusterhausen.
- Spongia Saxonica**, Geinitz, 1850 A, p. 264. (= *Spongites Saxonicus*, Geinitz, 1842.)
- Spongites Saxonicus**, Geinitz, 1842, p. 96, pl. xxiii, figs. 1, 2.
Quadersandstein; Saxony.

- Steinhauera globosa*, } Presl in Sternberg, 1838, p. 202.
 — *minuta*, }
 — *oblonga*, } Perucur Beds; Bohemia.
- Stenopteris? cretacea*, Hollick, 1902, p. 148, pl. iii, fig. 2.
 Laramie Formation; Colorado, U.S.A.
- *virginica*, Fontaine, 1889, p. 112, pl. xxi, fig. 8.
 Potomac Formation; Virginia, U.S.A.
- Sterculia aperta*, Lesquereux, 1883, p. 82, pl. x, figs. 2-3.
 Dakota Group; Kansas, U.S.A.
- *cliffwoodensis*, Berry, 1905 E, p. 88, pl. xliii, fig. 5.
 Matawan Formation; Cliffwood, U.S.A.
- *Drakei*, Cummins, 1892, p. 210, text-fig. 8.
 Dakota Group; New Mexico, U.S.A.
- *elegans*, Fontaine, 1889, p. 314, pl. clvii, fig. 2; pl. clviii, figs. 2, 3.
 Potomac Formation; Virginia, U.S.A.
- *Geinitzi*, Engelhardt, 1892 A, p. 101, pl. ii, figs. 3, 4, 7.
 Quader; Freiberg, Saxony.
- *Krejci*, Velenovsky, 1883, p. 47, pl. xiii, fig. 1.
 Cenomanian; Bohemia.
- *Labrusca*, Unger, 1851, p. 175, pl. xlix, figs. 1-11. Eocene; Stotzka.
 (Recorded American Cretaceous, Hollick, 1896.)
- *limbata*, Velenovsky, 1883, p. 46, pl. xiii, figs. 2-5; pl. xiv, fig. 1.
 Cenomanian; Bohemia.
- *lineariloba*, Lesquereux, 1876 B, p. 358.
 Dakota Group; Kansas, U.S.A.
- *lugubris*, Lesquereux, 1883, p. 81, pl. vi, figs. 1-3.
 Dakota Group (?); Colorado, U.S.A.
- *minima*, Berry, 1906 B, p. 177.
 Magothy Formation; New Jersey, U.S.A.
- *mucronata*, Lesquereux, 1892, p. 182, pl. xxx, figs. 1-4.
 Dakota Group; Kansas, U.S.A.
- *obtusiloba*, Lesquereux, 1883, p. 82, pl. viii, fig. 3. (Re-named
Aralia tripartita (Lx.), Knowlton, 1898.) Ibid.
- *pre-labrusca*, Hollick, 1906 A, p. 24, pl. xxxiv, figs. 21, 22.
 Middle Cretaceous; Martha's Vineyard, U.S.A.
- *reticulata*, Lesquereux, 1892, p. 185, pl. xxxiv, fig. 10.
 Dakota Group; Kansas, U.S.A.
- *Snowii*, Lesquereux, 1892, p. 183, pl. xxx, fig. 5; pl. xxxi, figs. 2, 3;
 pl. xxxii; pl. xxxiii, figs. 1-4. Ibid.
- *Snowii bilobata*, Berry, 1905 E, p. 89, pl. xliii, fig. 7.
 Matawan Formation; New Jersey, U.S.A.
- *Snowii disjuncta*, Lesquereux, 1892, p. 184, pl. lviii, fig. 6.
 Dakota Group; Kansas, U.S.A.
- *tripartita* (Lx.), Knowlton, 1898, p. 224. (= *Sterculia obtusiloba*,
 Lx., 1883, & *Aralia tripartita*, Lx., 1876.) Ibid.
- *variabilis*, Saporta, 1868, p. 400, pl. xii, figs. 6, 7; Heer, 1883 A,
 p. 88, pl. lvii, fig. 7. Patoot Beds; Greenland.

- Sterculia vetustula*, Dawson, 1886, p. 10, pl. iii, fig. 2.
Middle Cretaceous; Canada.
- *sp.*?, Hollick, 1898 b, p. 133, pl. xiv, figs. 4-7.
Upper Cretaceous; New Jersey, U.S.A.
- *sp.*?, Hollick, 1898 b, p. 422, pl. xxxvii, fig. 5.
Middle Cretaceous; Staten Island, U.S.A.
- *sp.*?, Hollick, 1906 a, p. 95, pl. xxxiv, figs. 18, 19.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- Stichus mermiscides*, Etheridge, 1904, p. 255, pls. xxx, xxxi.
Cretaceous; Australia.
- Stigmaria flexuosa*, Debey, 1848 a, p. 117 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- Strezeleckia gangamopteroides*, Johnston, 1896, p. 58, figs. 5-7.
Cretaceous (?); Tasmania.
- *tenuifolia*, Johnston, 1896, p. 58, fig. 8.
Ibid.
- Strobilites Bucklandi*, Lindley & Hutton, 1835, pl. cxxix; Gardner, 1886 a, p. 200.
Upper Greensand (?); Wiltshire.
- *Davisii*, Hollick & Jeffrey, 1909, p. 68, pl. iii, fig. 10.
Raritan Formation; Staten Island, U.S.A.
- *inquirendus*, Hollick, 1898 a, p. 130, pl. xi, fig. 1.
Upper Cretaceous; New Jersey, U.S.A.
- *microsporophorus*, Hollick & Jeffrey, 1909, p. 66, pl. x, figs. 18-21; pl. xxiv, figs. 2-6.
Raritan Formation; Staten Island, U.S.A.
- *perplexus*, Hollick, 1906 a, p. 107, pl. ii, fig. 43.
Middle Cretaceous; Martha's Vineyard, U.S.A.
- *sp.*, Hollick & Jeffrey, 1909, p. 69, pl. iii, fig. 9.
Raritan Formation; Staten Island, U.S.A.
- Sycophyllum dentatum*, Schulze, 1888, p. 464.
Perucér Beds; Bohemia.
- Symphocarpophyllum Albertum*, Dawson, 1887, p. 30, pl. ii, fig. 17.
Upper Laramie Formation; Canada.
- *Linnaeiforme*, Dawson, 1887, p. 30, pl. ii, fig. 18.
Ibid.
- Syringodendron*? *sp.*, Otto, 1854, p. 26, pl. iv, figs. 3, 4.
Lower Quader; Dippoldiswalde, Saxony.
- Tænidium alysioides*, Hosius & von der Marck, 1880, p. 131, pl. xxiv, fig. 5.
Upper Senonian; Westphalia.
- *lusitanicum*, Heer, 1881, p. 12, pl. xx. (= *Volubiliites lusitanicus*, Lorenz, 1901.)
Cretaceous; Portugal.
- *pinnatisectum*, Saporta, 1880, p. 642, pl. i, fig. 2.
Upper Neocomian; Havre, France.
- Tæniopteris arctica* (spelt *Tanæopteris*), Heer, 1871, p. 1181 [nomen nudum].
Kome Beds; Greenland.
- *auriculatum* (Fontaine), Berry, 1910 b, p. 634. (= *Angiopteridium auriculatum*, Fontaine, 1889.)
Patuxent Formation; Virginia, U.S.A.
- *deperdita*, Heer, 1883 a, p. 8, pl. xlviii, fig. 14.
Patoot Beds; Greenland.

- Tæniopteris Gibbsii**, Newberry, 1863, p. 512. (= *Nilssonia Gibbsii*, Hollick in Newberry, 1898.) Cretaceous (?); Washington, U.S.A.
- **kuchelbadensis**, Feistmantel, 1874, p. 269. Perucur Beds; Bohemia.
- **nervosum** (Fontaine), Berry, 1910 B, p. 634. (= *Angiopteridium nervosum*, and others, Fontaine, 1889). Fatuxent Formation; Virginia, U.S.A.
- **orovillensis**, Fontaine in Penhallow, 1902 B, p. 37. Upper Cretaceous; Vancouver Island.
- **plumosa**, Dawson, 1883, p. 24, pl. iv, fig. 15. Upper Cretaceous; Vancouver Island.
- **sp.**, Heer, 1878, p. 30, pl. viii, fig. 19. Lower Cretaceous (?); Siberia.
- Tænioxylon varians**, Felix (**cretaceum**), Vater, 1884, p. 852. Lower Senonian; Harzburg.
- Tacnurus incertus**, Dawson, 1886, p. 10. Kootanie Formation; British Columbia, Canada.
- **Saportai**, Dewalque, 1880, p. 43, pl. i, figs. 1 a, 1 b, 2 a, 2 b. Senonian; Anzin, France.
- **tenuestriatus**, Heer, 1877, p. 145, pl. lvii, figs. 7-10. Upper Cretaceous; Switzerland.
- Taxites pecten**, Heer, 1883 A, p. 9, pl. liii, figs. 9, 9 b. Patoot Beds; Greenland.
- Taxodium (Glyptostrobus) brookense**, Fontaine, 1889, p. 254, pl. cxvii, fig. 1; pl. cxviii, figs. 3-9; pl. cxxxi, fig. 5; pl. clxv, figs. 1-3; pl. clxvi, figs. 4, 7; pl. clxvii, fig. 3. Potomac Formation; Virginia, U.S.A.
- (**Glyptostrobus**) **brookense**, var. **angustifolium**, Fontaine, 1889, p. 256, pl. clxvii, fig. 1. Ibid.
- **cuneatum**, Newberry, 1863, p. 517. Kootanie Formation; British Columbia, Canada.
- (**Glyptostrobus**) **denticulatum**, Fontaine, 1889, p. 253, pl. cxviii, fig. 1. Potomac Formation; Virginia, U.S.A.
- (**Glyptostrobus**) **expansum**, Fontaine, 1889, p. 252, pl. cxxiii, fig. 1. Ibid.
- (**Glyptostrobus**) **fastigiatum**, Fontaine, 1889, p. 253, pl. cxv, figs. 1, 3. Ibid.
- (**Glyptostrobus**) **ramosum**, Fontaine, 1889, p. 251, pl. cxxiii, figs. 2, 3; pl. cxviii, fig. 2; pl. cxxvii, fig. 1; pl. cxxviii, fig. 1; pl. clxvi, fig. 1. Ibid.
- (**Glyptostrobus**) **virginicum**, Fontaine, 1889, p. 252, pl. cxxi, fig. 6. Ibid.
- **sp.**, Dawson, 1894, p. 56, pl. vi, fig. 10. Upper Cretaceous; Vancouver Island, Canada.
- Taxo-Torreya trinervia**, Ettingshausen, 1887 A, p. 176, pl. vii, figs. 16, 16 a. Upper Cretaceous; New Zealand.
- Taxoxylon cretaceum**, Unger, 1859, p. 231, pl. iii, figs. 12-14. (= *Cedroxylon cretaceum*, Kr. in Schimper, 1870.) Quadersandstein; Amberg, Germany.

- Taxoxylon halternianum**, Hosius & von der Marck, 1880, p. 194,
pl. xli, figs. 166-168. Lower Senonian; Westphalia.
- Tempskya cretacea**, Hosius & von der Marck, 1880, p. 192, pl. xxxix,
figs. 161-163. Ibid.
- **varians** (Corda), Velenovsky, 1888 B, p. 23, pl. vi, figs. 1-7; pl. v,
fig. 5, & text-fig. Perucer Beds; Bohemia.
- Terminalia rectinervis**, Velenovsky, 1886, p. 52, pl. xx, figs. 1, 2.
(= *Terminaliphyllum rectinerve*, Velenovsky, 1889.)
Cenomanian; Kaunic, Bohemia.
- Terminaliphyllum rectinerve**, Velenovsky, 1889. (= *Terminalia
rectinervis*, Velenovsky, 1886.)
- Ternstroemia crassipes**, Velenovsky, 1886, p. 54, pl. xviii, figs. 3, 4.
(= *Ternstræmiphyllum crassipes*, Velenovsky, 1889.)
Cenomanian; Vyserovic, Bohemia.
- Ternstræmiphyllum crassipes**, Velenovsky, 1889. (= *Ternstroemia
crassipes*, Velenovsky, 1886.)
- Tetraphyllum dubium**, Hosius & von der Marck, 1880, p. 137,
pl. xxv, fig. 14. Upper Senonian; Westphalia.
- **oblongum**, Heer, 1882, p. 105, pl. xxvi, figs. 5 b, 6.
Atane Beds; Greenland.
- Thalassocharis Binkhorsti**, Debey, 1865, p. 57.
Cretaceous; Prussia.
- **Bosqueti**, Debey, 1851, p. 568; & in Hosius & von der Marck, 1880,
p. 146. Senonian; Aix, Rhenish Prussia.
- **Bosqueti, forma brevi-articulata**, Miquel, 1853, p. 51, pl. vi,
fig. 2. Cretaceous; Belgium.
- **Bosqueti, forma lata**, Miquel, 1853, p. 51, pl. vi, fig. 3.
Upper Senonian; Belgium.
- **Muelleri**, Debey, 1848 A, p. 119; & 1851, p. 568.
Senonian; Aix, Rhenish Prussia.
- **westfalica**, Hosius & von der Marck, 1880, p. 147, pl. xxvi, figs. 25-
28, 30-34; pl. xxvii, figs. 29, 35-39; pl. xxviii, figs. 40-42.
Upper Senonian; Westphalia.
- Thinnfeldia arctica**, Heer, 1874 A, p. 123, pl. xxxv, figs. 11-16;
pl. xxxvi, fig. 10 b. Cretaceous; Spitzbergen.
- **Buftoni**, Johnston, 1896, p. 61, fig. 18.
Cretaceous (?); Tasmania.
- **Fontainei**, Berry, 1903 D, p. 443 (re-naming *Thinnfeldia variabilis*,
Fontaine, 1889). Potomac Formation; Virginia, U.S.A.
- **granulata**, Fontaine, 1889, p. 111, pl. xxvi, figs. 10-12; pl. xxvii,
figs. 1-5, 8; pl. clxix, fig. 1. Ibid.
- **lanceolata**, Knowlton in Weed & Knowlton, 1893, p. 49, pl. v,
fig. 5. (Re-named *Protophyllocladus lanceolatus*, Berry, 1903 D.)
Laramie Formation; Montana, U.S.A.
- **Lesquereuxiana**, Heer, 1882, p. 37, pl. xliv, figs. 9-10; pl. xlvii,
figs. 1-12. Atane Beds; Greenland.
- **marylandica**, Fontaine in Ward, 1905, p. 541, pl. cxiv, figs. 8, 9.
Older Potomac Formation; Maryland, U.S.A.

- Thinnfeldia montana**, Knowlton, 1898, p. 227 (re-naming *Thinnfeldia polymorpha* (Lesquereux), Knowlton, = *Salisburia polymorpha*, Lesquereux). Montana Formation; Coal Banks, U.S.A.
- **montanense**, Fontaine in Weed & Pirsson, 1898, p. 481. Lower Cretaceous; Montana, U.S.A.
- **polymorpha** (Lesquereux), Knowlton in Weed & Knowlton, 1893. (= *T. montana*, Knowlton, re-named *Protophyllocladus polymorphus* by Berry, 1903 D.)
- **polymorpha**, Johnston (*non* Schenk *nec* Knowlton), 1896, p. 62, fig. 16. Cretaceous (?); Tasmania.
- **rotundiloba**, Fontaine, 1889, p. 111, pl. xxvii, figs. 6, 7. Potomac Formation; Virginia, U.S.A.
- **subintegrifolia** (Lx.), Knowlton, 1898, p. 228 (re-naming *Phyllocladus subintegrifolius*, Lesquereux, 1868 & 1874). (= *Protophyllocladus subintegrifolius*, Berry, 1903 D.) Dakota Group; Nebraska, U.S.A.
- **variabilis**, Fontaine (*non* Velenovsky), 1889, p. 110, pl. xvii, figs. 3-7; pl. xviii, figs. 1-6. (= *Thinnfeldia Fontainei*, re-named by Berry, 1903 D.) Potomac Formation; Virginia, U.S.A.
- **variabilis**, Velenovsky, 1885, p. 6, pl. ii, figs. 1-5. (Re-named *Sagenopteris variabilis*, Velenovsky, 1889.) Perucur Beds; Bohemia.
- Thuites alienus**, Sternberg, 1825, p. xxxviii, pl. xlv, fig. 1. Cretaceous; Bohemia.
- **crassus**, Lesquereux, 1883, p. 32. Dakota Group; Kansas, U.S.A.
- **gramineus**, Sternberg, 1825, p. 31, p. xxxviii, pl. xxxv, fig. 4. Perucur Beds; Bohemia.
- **Hoheneggeri**, Ettingshausen, 1852 c, p. 26, pl. i, figs. 6, 7. (= *Frenelopsis Hoheneggeri*, Schenk, 1871 A.) Wernsdorfer Beds (see Krasser, 1896, pp. 146, 147.)
- **Kurrianus**, Dunker, 1846, p. 20, pl. vii, fig. 8. (= *Sphenolepidium Kurrianum* (Dunk.), Heer, 1881.) Wealden; Germany.
- **Meriani**, Heer, 1874 A, p. 73, pl. xvi, figs. 17, 18. Kome Beds; Greenland.
- **Pfaffii**, Heer, 1874 A, p. 100, pl. xxxi, figs. 8 b, c, d, e. Atane Beds; Greenland.
- **Wilkinsoni**, Ettingshausen, 1893, p. 147; & 1895, p. 11, pl. i, figs. 7-9. Cretaceous; Australia.
- **sp.**, Hollick & Jeffrey, 1909, p. 31, pl. viii, figs. 12-18; pl. xxvii, figs. 4-6; pl. xxviii, figs. 1-4. Raritan Formation; Staten Island, U.S.A.
- Thuja cretacea** (Heer), Newberry, 1895, p. 53, pl. x, figs. 1, 1 a. Amboy Clay; New Jersey, U.S.A.
- Thuya sp.**, Cornuel, 1866, p. 673, pl. xii, fig. 23. Neocomian; France.
- **sp.**, Jasche, 1858, p. 97, pl. iv, figs. 8, 9. Cretaceous; Saxony.
- Thuyites debilis**, Saporta, 1894, p. 115, pl. xxii, fig. 6. Aptian; Portugal.
- **densior**, Saporta, 1894, p. 109, pl. xx, fig. 6. Ibid.

- Thuyites pulchelliformis*, Saporta, 1894, p. 109, pl. xix, figs. 21-24;
pl. xx, figs. 7-10; p. 115, pl. xxi, fig. 7. Aptian; Portugal.
- *sp.*, Nathorst, 1891, p. 27, pl. i, figs. 8, 9, 10. Upper Cretaceous (?); Rostock.
- Thuyoxylon americanum*, Unger in Roemer, 1852, p. 95. Cretaceous (?); Texas, U.S.A.
- Thyrsopteris alata*, Fontaine, 1889, p. 124, pl. xxxvi, fig. 5. Potomac Formation; Virginia, U.S.A.
- *angustifolia*, Fontaine, 1889, p. 131, pl. xlv, fig. 4; pl. xlv, fig. 3;
pl. xlviii, fig. 2; pl. xlix, figs. 3, 4; pl. lv, fig. 2; pl. lviii, fig. 8. Ibid.
- *angustiloba*, Fontaine, 1889, p. 134, pl. xlviii, figs. 3-5; pl. lv, fig. 3. Ibid.
- *bella*, Fontaine, 1889, p. 139, pl. liii, fig. 5; pl. lv, figs. 6, 7; pl. lvi, figs. 2, 5; pl. lvii, figs. 1, 5; pl. lviii, fig. 4. Ibid.
- *brevifolia*, Fontaine, 1889, p. 121, pl. xxiv, figs. 5, 10. Ibid.
- *brevipennis*, Fontaine, 1889, p. 124, pl. xxxiv, fig. 3; pl. xxxvi, fig. 2; pl. xxxvii, figs. 3, 9; pl. xxxviii, fig. 1; pl. xli, fig. 4. Ibid.
- *capsulifera*, Velenovsky, 1888 B, p. 10, pl. i, figs. 6-12. (= *Ongchiopsis capsulifera*, Nathorst, 1890.) Perucer Beds; Bohemia.
- *crassinervis*, Fontaine, 1889, p. 130, pl. xli, figs. 1-3. Potomac Formation; Virginia, U.S.A.
- *crenata*, Fontaine, 1889, p. 127, pl. xxxix, figs. 1, 2. Ibid.
- *decurrens*, Fontaine, 1889, p. 130, pl. xliii, fig. 7; pl. xlv, figs. 2, 4; pl. xlix, figs. 5-7. Ibid.
- *densifolia*, Fontaine, 1889, p. 129, pl. xxxix, fig. 3; pl. xl, figs. 2-5; pl. li, fig. 5. Ibid.
- *dentata*, Fontaine, 1889, p. 121, pl. xxiv, figs. 4, 6, 7, 9; pl. xxv, figs. 1, 2. Ibid.
- *dentifolia*, Fontaine in Ward, 1899 B, p. 660, pl. clxi, figs. 6-9. Lower Cretaceous; Black Hills, U.S.A.
- *distans*, Fontaine, 1889, p. 124, pl. xlvii, fig. 3; pl. liv, fig. 8. Potomac Formation; Virginia, U.S.A.
- *divaricata*, Fontaine, 1889, p. 125, pl. xxxvii, figs. 5-8; pl. clxx, fig. 1. Ibid.
- *elliptica*, Fontaine, 1889, p. 133, pl. xxiv, fig. 3; pl. xlv, fig. 1; pl. l, figs. 6, 9; pl. li, figs. 4, 6, 7; pl. liv, fig. 6; pl. lv, fig. 4; pl. lvi, figs. 6, 7; pl. lvii, fig. 6; pl. lviii, fig. 2. Ibid.
- *grevillioides* (Heer), Hollick, 1906 A, p. 31, pl. i, figs. 10-13. (= *Sphenopteris grevillioides*, Heer, 1874 A.) Middle Cretaceous; Martha's Vineyard, U.S.A.
- *heteroloba*, Fontaine, 1889, p. 139, pl. liii, fig. 4. Potomac Formation; Virginia, U.S.A.
- *heteromorpha*, Fontaine, 1889, p. 136, pl. lii, fig. 1. Ibid.
- *heterophylla*, Fontaine, 1889, p. 142, pl. lviii, fig. 3. Ibid.
- *inaequipinnata*, Fontaine, 1889, p. 142, pl. lvii, figs. 3, 8. Ibid.
- *insignis*, Fontaine, 1889, p. 127, pl. xxxix, fig. 4; pl. xl, fig. 1; pl. xli, fig. 6; pl. xlii, figs. 1, 2, 4; pl. xliii, figs. 1, 3. Ibid.

- Thyrsopteris insignis angustipennis*, Fontaine, 1889, p. 128, pl. xlii, fig. 3; pl. xliii, fig. 2. Potomac Formation; Virginia, U.S.A.
- *Meekiana*, Fontaine, 1889, p. 125, pl. xxxviii, figs. 2-4, 8; pl. 1, figs. 7, 8; pl. li, fig. 3. Potomac Formation; Maryland, U.S.A.
- *Meekiana angustiloba*, Fontaine, 1889, p. 126, pl. xxxviii, figs. 5-7, 9; pl. xliii, fig. 8; pl. xlv, fig. 3; pl. xlvii, fig. 4; pl. xlviii, fig. 1; pl. liv, figs. 2, 11; pl. lv, fig. 1; pl. lvi, figs. 1, 3. Ibid.
- *microloba*, Fontaine, 1889, p. 140, pl. lvii, fig. 4. Potomac Formation; Virginia, U.S.A.
- *microloba alata*, Fontaine, 1889, p. 140, pl. lv, fig. 5; pl. lviii, fig. 1. Ibid.
- *microphylla*, Fontaine, 1889, p. 131, pl. xlv, figs. 1, 2, 4, 5. Ibid.
- *nana*, Fontaine, 1889, p. 141, pl. lvi, figs. 4, 8. Ibid.
- *nervosa*, Fontaine, 1889, p. 122, pl. xxv, figs. 4, 5, 16; pl. xxxvii, figs. 2, 4; pl. xxxix, fig. 5; pl. xl, fig. 6. Ibid.
- *obtusiloba*, Fontaine, 1889, p. 143, pl. lviii, figs. 7, 10. Ibid.
- *pachyphylla*, Fontaine, 1889, p. 135, pl. 1, fig. 3. Ibid.
- *pachyrachis*, Fontaine, 1889, p. 132, pl. xlvi, figs. 3, 5; pl. xlvii, figs. 1, 2; pl. xlix, fig. 1. Ibid.
- *pecopteroides*, Fontaine, 1889, p. 135, pl. li, fig. 1. Ibid.
- *pinnatifida*, Fontaine, 1889, p. 136, pl. li, fig. 2; pl. liv, figs. 4, 5, 7; pl. lvii, fig. 7. Ibid.
- *rarinervis*, Fontaine, 1889, p. 123, pl. xxvi, figs. 6, 7; pl. xliii, figs. 4-6; pl. xlv, figs. 1, 2, 5; pl. xlix, fig. 2; pl. clxix, figs. 6, 7. Ibid.
- *retusa*, Fontaine, 1889, p. 144, pl. lix, fig. 10. Ibid.
- *rhombifolia*, Fontaine, 1889, p. 138, pl. lii, fig. 5; pl. liv, fig. 1. Ibid.
- *rhombiloba*, Fontaine, 1889, p. 144, pl. lix, figs. 6, 7; pl. lx, fig. 8. Ibid.
- *sphenopteroides*, Fontaine, 1889, p. 143, pl. lviii, fig. 6. Ibid.
- *squarrosa*, Fontaine, 1889, p. 143, pl. lix, fig. 3. Ibid.
- *varians*, Fontaine, 1889, pl. lii, figs. 2-4; pl. liii, figs. 1-3; pl. liv, fig. 10; pl. lvii, fig. 2. Ibid.
- *virginica*, Fontaine, 1889, p. 120, pl. xxiv, fig. 1. Ibid.
- Tillæphyllum dubium*, Newberry, 1895, p. 109, pl. xv, fig. 5. Amboy Clay; New Jersey, U.S.A.
- Todea? saportanea*, Lesquereux, 1874, p. 48, pl. xxix, figs. 1-4. (= *Lomatia saportanea*, Lesquereux, 1876 B.) Dakota Group; Kansas, U.S.A.
- Torreyia densifolia* (spelt *Torreia*), Dawson, 1883, p. 25, pl. v, figs. 20, 20 a. (Re-named *Tumion densifolium* by Knowlton, 1898.) Upper Cretaceous; Protection Island, Canada.
- *Dicksoniana*, Heer, 1871, p. 1182; & 1874 A, p. 70, pl. xviii, figs. 1 a, 2 a, 3, 4. Kome Beds; Greenland.

- Torreya dicksonioides** (spelt **Torreia**), Dawson, 1883, p. 21, pl. ii, fig. 4. (Re-named *Tumion dicksonioides* by Knowlton, 1898.)
Upper Cretaceous; North-West Territory, Canada.
- **falcata**, Fontaine, 1889, p. 235, pl. cxiii, fig. 4. (Re-named *Tumion falcatum* by Knowlton, 1898.)
Potomac Formation; Virginia, U.S.A.
- **oblanceolata**, Lesquereux, 1883, p. 30, pl. i, fig. 2. (Re-named *Tumion oblanceolatum* by Knowlton, 1898.)
Dakota Group (?); Colorado, U.S.A.
- **parvifolia**, Heer, 1874 A, p. 71, pl. xvii, figs. 1, 2.
Kome Beds; Greenland.
- **venusta**, Yokoyama, 1894, p. 230, pl. xxii, figs. 11, 12, 12 a.
Neocomian; Japan.
- **virginica**, Fontaine, 1889, p. 234, pl. cix, fig. 8. (Re-named *Tumion virginicum* by Knowlton, 1898.)
Potomac Formation; Virginia, U.S.A.
- Trametes Pini**, Conwentz, 1892, pp. 13, 21, pl. vii, fig. 1. (= *Trametites Pini*, Meschinelli, 1892.)
Senonian; Sweden.
- Trametites Pini**, Meschinelli, 1892, p. 747. (= *Trametes Pini*, Conwentz, 1892.)
- Trapa ? cuneata**, Knowlton, 1900 A, p. 64, pl. v, fig. 6.
Montana Formation; Wyoming, U.S.A.
- **? microphylla**, Lesquereux, 1876 c, p. 369; & 1878, p. 295, pl. lxi, figs. 16-17 a.
Ibid.
- Tricalycites major**, Hollick, 1905 c, p. 416, pl. lxxii, figs. 3-7.
Cretaceous; Long Island, U.S.A.
- **papyraceus**, Newberry MS. in Hollick, 1894, p. 63, pl. clxxx, fig. 8; & Newberry, 1895, p. 132, pl. xlvi, figs. 30-38.
Amboy Clay; Woodbridge, U.S.A.
- Tricarpellites striatus**, Newberry, 1895, p. 132, pl. xlvi, figs. 9-13.
Ibid.
- Trichomanes Riccioides**, Heer, 1871, p. 1182 [nomen nudum].
Atane Beds; Greenland.
- Trichopitys sp.**, Nathorst, 1885, p. 286, text-fig. 4.
Ibid.
- **sp. ?**, Kerner, 1896, p. 52, pl. iv, fig. 5.
Cretaceous; Lesina, Dalmatia.
- Trichosporites Conwentzi**, Felix, 1894, p. 273.
Upper Cretaceous; Sweden.
- Triplaris cenomanica**, Engelhardt, 1892 A, p. 96, pl. ii, fig. 5.
Quader; Freiberg, Saxony.
- Triploporella Fraasi**, Steinmann, 1880, p. 136, pl. v, figs. 1-8.
Turonian; Lebanon.
- **Fraasi minor**, Steinmann, 1899, p. 193, text-figs. 38, 39.
Cenomanian; Mexico.
- Tsugites magnus**, Fliche, 1896, p. 211, pl. ix, fig. 2.
Albian; Clermont, France.
- Tumion carolinianum**, Berry, 1908 n, p. 383, text-figs. 1-3.
Middle Cretaceous; North Carolina, U.S.A.

- Tumion densifolium** (Dawson), Knowlton, 1898, p. 234 (re-naming *Torreyia densifolia*, Dawson, 1883).
- **dicksonioides** (Dawson), Knowlton, 1898, p. 234 (re-naming *Torreyia dicksonioides*, Dawson, 1883).
- **falcatum** (Fontaine), Knowlton, 1898, p. 234 (re-naming *Torreyia falcata*, Fontaine, 1889).
- **oblanceolatum** (Lx.), Knowlton, 1898, p. 234 (re-naming *Torreyia oblanceolata*, Lesquereux, 1883).
- **virginicum** (Fontaine), Knowlton, 1898, p. 234 (re-naming *Torreyia virginica*, Fontaine, 1889).
- Typha?** sp., Hollick, 1894, p. 63, pl. clxxx, fig. 9.
Upper Cretaceous; Long Island, U.S.A.
- **gigantea**, Unger, 1870 A, pl. iii, fig. 6.
(Includes supposed *Algæ* from Quadersandstein.)
- Typhacites lævis**, Saporta, 1890, p. 3, pl. xiii, figs. 5, 5 a.
Uppermost Cretaceous; France.
- **rugosus**, Saporta, 1890, p. 3, pl. xiii, figs. 4, 4 a. Ibid.
- Typhæloipum cretaceum**, Krasser, 1896, p. 127, pl. xii, fig. 4.
Cenomanian; Kunstadt, Moravia.
- Tysonia marylandica**, Fontaine, 1889, p. 193, pls. clxxiv-clxxx.
(= *Cycadeoidea marylandica* [(Font.), Capellini & Solms-Laubach, 1892.])
Potomac Formation; Maryland, U.S.A.
- Ulmiphyllum Brookense**, Fontaine, 1889, p. 312, pl. clv, fig. 8;
pl. clxiii, fig. 7. Potomac Formation; Virginia, U.S.A.
- **crassinerve**, Fontaine, 1889, p. 313, pl. clviii, figs. 6, 7. Ibid.
- **densinerve**, Fontaine in Ward, 1899 B, p. 689, pl. clxix, fig. 7.
Lower Cretaceous; Black Hills, U.S.A.
- **tenuinerve**, Fontaine, 1889, p. 313, pl. clviii, fig. 1.
Potomac Formation; Virginia, U.S.A.
- Ulmophyllum latifolium**, Ettingshausen, 1887 A, p. 184, pl. ix,
figs. 6-8. Upper Cretaceous; New Zealand.
- **planaræfolium**, Ettingshausen, 1887 A, p. 184, pl. ix, figs. 2-5, 4 a.
Ibid.
- **priscum**, Dawson, 1894, p. 59, pl. viii, fig. 28.
Upper Cretaceous; Vancouver Island, Canada.
- Ulmus dubia**, Dawson, 1883, p. 27, pl. vii, fig. 29. Ibid.
- **præcursor**, Dawson, 1887, p. 28, pl. ii, fig. 11.
Laramie Formation; Canada.
- Uredinites cretaceus**, Velenovsky, 1889, p. 29, pl. iii, fig. 14.
(= *Puccinities cretaceus*, Velenovsky, 1889, same figure.)
Perucer Beds; Bohemia.
- Vaccinium** sp.?, Kerner, 1896, p. 55, pl. v, fig. 5.
Cretaceous; Lesina, Dalmatia.
- Vesquia Tournaisii**, Bertrand, 1883, p. 1382; & 1883 A, p. 294.
Cenomanian; Tournai, France.

- Viburnites crassus**, Knowlton in Lesquereux, 1892, p. 124, pl. xlv, figs. 1-4. Dakota Group; Kansas, U.S.A.
- **Evansanus**, Ward, 1894, p. 262; & 1899 B, p. 709, pl. clxxii, figs. 3, 4. Dakota Group; S. Dakota, U.S.A.
- **Masoni**, Lesquereux, 1892, p. 125, pl. xlv, fig. 5. Dakota Group; Kansas, U.S.A.
- Viburnum anomalum**, Knowlton, 1900 A, p. 72. Montana Formation; Wyoming, U.S.A.
- **attenuatum**, Heer, 1883 A, p. 34, pl. lxiii, fig. 5. Patoot Beds; Greenland.
- **contortum**, Lesquereux, 1873, p. 396. Laramie Formation; Wyoming, U.S.A.
- **dichotomum**, Lesquereux, 1873, p. 399; & 1878, p. 225, pl. xxxviii, fig. 6. Ibid.
- **ellsworthianum**, Lesquereux, 1892, p. 121, pl. xxi, fig. 6. Dakota Group; Kansas, U.S.A.
- **grewiopsideum**, Lesquereux, 1892, p. 120, pl. xxi, fig. 4. Ibid.
- **Hollickii**, Berry, 1903 C, p. 683, text-figs. 5, 6. Matawan Formation; New Jersey, U.S.A.
- **inæquilaterale**, Lesquereux, 1892, p. 119, pl. xxi, figs. 2, 3. Dakota Group; Kansas, U.S.A.
- **integrifolium**, Newberry MS. in Hollick, 1894 A, p. 54, pl. clxxvii, fig. 7; & Newberry, 1895, p. 125, pl. xli, fig. 1. Cretaceous; Long Island, U.S.A.
- **Lesquereuxii**, Ward in Lesquereux, 1892, p. 121. Dakota Group; Dakota, U.S.A.
- **Lesquereuxii commune**, Lesquereux, 1892, p. 122, pl. liii, fig. 2. Dakota Group; Kansas, U.S.A.
- **Lesquereuxii cordifolium**, Lesquereux, 1892, p. 122, pl. lii, fig. 9. Ibid.
- **Lesquereuxii lanceolatum**, Lesquereux, 1892, p. 123, pl. liii, fig. 3. Ibid.
- **Lesquereuxii latius**, Lesquereux, 1892, p. 123, pl. liii, fig. 10. Ibid.
- **Lesquereuxii longifolium**, Lesquereux, 1892, p. 122, pl. liii, fig. 1. Ibid.
- **Lesquereuxii rotundifolium**, Lesquereux, 1892, p. 122, pl. lii, fig. 8. Ibid.
- ? **Lesquereuxii tenuifolium**, Lesquereux, 1892, p. 123, pl. lxiv, fig. 13. Ibid.
- **Mattewanense**, Berry, 1904 A, p. 80, pl. iv, fig. 13. Matawan Formation; New Jersey, U.S.A.
- **montanum**, Knowlton, 1900 A, p. 73, pl. xix, figs. 1, 2. Montana Formation; Wyoming, U.S.A.
- **multinerve**, Heer, 1883 A, p. 33, pl. lxiii, figs. 1-4. Patoot Beds; Greenland.
- **multinerve**, var. **b**, Heer, 1883 A, p. 34, pl. lxiii, fig. 4. Ibid.

- Viburnum oxycoccoides*, Dawson, 1886, p. 17,
Upper Cretaceous (?); Canada.
- *platanoides*, Lesquereux, 1878 B, p. 224, pl. xxxviii, fig. 8. (Re-named *Platanus platanoides* (Lesquereux), Knowlton, 1898.)
Laramie Formation; Wyoming, U.S.A.
- ? *problematicum*, Knowlton, 1900 A, p. 71, pl. xix, fig. 4.
Montana Formation; Utah, U.S.A.
- *robustum*, Lesquereux, 1892, p. 120, pl. xx, figs. 4-6.
Dakota Group; Kansas, U.S.A.
- *rotundifolium*, Lesquereux, 1876 C, p. 368; & 1878 B, p. 225,
pl. xxxvii, fig. 12; pl. xxxviii, fig. 10; pl. lxi, fig. 22.
Montana Formation; Wyoming, U.S.A.
- *Saskatchuense*, Dawson, 1888, p. 35.
Belly River Series; Canada.
- *sphenophyllum*, Knowlton in Lesquereux, 1892, p. 123, pl. liii,
fig. 4. Dakota Group; Kansas, U.S.A.
- *subrepandum*, Hosius & von der Marck, 1880, p. 190, pl. xl,
fig. 158. (= *Phyllites curvinervis*, Hosius, 1870 A.)
Lower Senonian; Westphalia.
- *vetus*, Saporta, 1894, p. 204, pl. xxxvi, fig. 9.
Upper Albian; Portugal.
- *zyziphoides*, Heer, 1883 A, p. 34, pl. lx, fig. 2.
Patoot Beds; Greenland.
- Vitex pentadactyla*, Velenovsky, 1882 A, p. 213 [nomen nudum].
Cretaceous; Bohemia.
- Vitiphyllum crassifolium*, Fontaine, 1889, p. 308, pl. cl, figs. 9, 10.
Potomac Formation; Virginia, U.S.A.
- *multifidum*, Fontaine, 1889, p. 309, pl. clxxiii, figs. 1-9.
Potomac Formation; Maryland, U.S.A.
- *parvifolium*, Fontaine, 1889, p. 309, pl. clxxii, figs. 11, 12. Ibid.
- Vitis Bruneri*, Ward, 1887, p. 69, pl. xxxii, figs. 1, 2.
Laramie Formation; Wyoming, U.S.A.
- *Carbonensis*, Ward, 1887, p. 70, pl. xxxii, fig. 3. Ibid.
- *sparsa*, Lesquereux, 1878 B, p. 241, pl. lx, fig. 24. Ibid.
- Volubilites lusitanicus*, Lorenz, 1901, p. 566. (= *Tenidium lusi-tanicum*, Heer, 1881.)
Cretaceous; Portugal.
- Weichselia erratica*, Nathorst, 1891, p. 24, pl. i, figs. 1-4.
Lower Cretaceous; Mecklenburg, nr. Rostock.
- *Ludovicæ*, Stiehler, 1857, p. 453; & 1858, p. 73, pl. xii; pl. xiii.
Senonian; Quedlinburg, Saxony.
- *reticulata* (Stokes & Webb), Ward, 1899 B, p. 651, pl. clx, figs. 2-4.
(= *Pecopteris reticulata*, Stokes & Webb, 1824.)
Lower Cretaceous; Black Hills, U.S.A.

- Widdringtonia complanata**, Lesquereux, 1876 B, p. 336; & 1878 B, p. 72, pl. lxii, figs. 13, 14. Montana Formation; Wyoming, U.S.A.
- **parvivalvis**, Bayer, 1893, pp. 14 & 40, fig. 11; & Bayer in Fritsch, 1893, p. 129, text-fig. 182. Senonian; Priesen, Bohemia.
- **Reichii** (Ettingshausen), Velenovsky, 1887 A, p. 639, figs. 14-16 on plate. Cenomanian; Bohemia.
- **sarthacensis**, Crié, 1879 (?), p. 30 [nomen nudum]. Cretaceous; Sainte-Croix, France.
- Widdringtonites debilis**, Saporta, 1894, p. 93, pl. xvii, fig. 14; p. 105, pl. v, fig. 19; pl. xvii, fig. 14; pl. xix, figs. 6-8. Valanginian & Neocomian; Portugal.
- **fasciculatus**, Hollick, 1906 A, p. 45, pl. iv, fig. 1. Middle Cretaceous: Martha's Vineyard, U.S.A.
- **fastigiatus**, Endlicher, 1847 B, p. 272. (= *Caulerpites fastigiatus*, Sternberg, 1838.) Middle Cretaceous; Bohemia.
- **gracilis**, Heer, 1868, p. 83, pl. xliii, figs. 1 e, f, g, 3 c. Kome Beds; Greenland.
- **pygmæus**, Saporta, 1894, p. 93, pl. xviii, fig. 10. Valanginian; Portugal.
- **Reichii** (Ett.), Heer, 1882, p. 51, pl. xxviii, fig. 5. (= *Frenelites Reichii*, Ettingshausen, 1867 A.) Atane Beds; Greenland.
- **subtilis**, Heer, 1874 A, p. 101, pl. xxviii, figs. 1, 1 b. Atane Beds; Greenland.
- Williamsonia? Bibbinsi**, Ward, 1905, p. 554, pl. cxv, fig. 11. Older Potomac Formation; Maryland, U.S.A.
- **cretacea**, Heer, 1882, p. 59, pl. xii, fig. 1; pl. xiii, fig. 9. Atane Beds; Greenland.
- **delawarensis**, Berry, 1907 A, p. 678, text-fig. 4. Magothy Formation; Delaware, U.S.A.
- **docata**, Lesquereux, 1892, p. 87, pl. ii, figs. 9, 9 a. Dakota Group; Kansas, U.S.A.
- ? **gallinacea**, Ward, 1905, p. 485, pl. cvii, fig. 4. Older Potomac Formation; Virginia, U.S.A.
- **minima**, Saporta, 1894, p. 105, pl. xix, fig. 9. Neocomian (?); Portugal.
- ? **phoenicopsoides**, Ward, 1899 B, p. 668, pl. elxii, fig. 20. Lower Cretaceous; Black Hills, U.S.A.
- **problematica** (Newberry), Ward, 1895, p. 382. (= *Palaeanthus (Williamsonia) problematicus*, Newberry, 1895.) Amboy Clay; New Jersey, U.S.A.
- **recentior**, Dawson, 1886, p. 12, pl. iv, fig. 1. Mill Creek Series; Canada.
- ? **Reisii**, Hollick, 1893, p. 37, pl. i, figs. 2, 3. Raritan Formation; Staten Island, U.S.A.
- **Smockii**, Newberry, 1895, p. 127, pl. xxxvi, figs. 1-8. Amboy Clay; New Jersey, U.S.A.
- **texana**, Fontaine, 1894, p. 278, pl. xliii, figs. 1, 2. Trinity Division; Texas, U.S.A.

- Williamsonia virginiensis**, Fontaine, 1889, p. 273, pl. cxxiii, figs. 5-7; pl. clxv, fig. 5. Potomac Formation; Virginia, U.S.A.
- Winchellia triphylla**, Lesquereux, 1893, p. 209, pl. viii, fig. 1. Upper Cretaceous; Montana, U.S.A.
- Woodwardia crenata**, Knowlton, 1900 A, p. 22, pl. iii, fig. 3. Montana Formation; Wyoming, U.S.A.
- *sp.*, Knowlton, 1900 A, p. 23, pl. iii, fig. 9. Ibid.
- Xylolithes**, Debey, 1848. (Cancelled by Debey, 1850, p. 117.)
- Xylomites aggregatus**, Heer, 1882, p. 21, pl. xxix, figs. 11, 11 b. Afane Beds; Greenland.
- **ellipticus**, Ettingshausen (*non* Feistmantel), 1867 A, p. 243, pl. i, fig. 7. Cenomanian; Niederschoena, Saxony.
- Yatesia Guillaumoti**, Fliche, 1896, p. 143, pl. i, fig. 4. Albian; Clermont, France.
- **Morrisii**, Carruthers, 1870, p. 688, pl. lv, figs. 3-6. (= *Cycadeoidea Yatesii* (Morris), Carruthers, 1867.) Lower Greensand; Potton.
- Yezonia vulgaris**, Stopes & Fuijii, 1910, pp. 23-32, pl. ii, figs. 5-8; pl. iii, fig. 9; pl. iv, fig. 19; text-figs. 8-11. Upper Cretaceous; Hokkaido, Japan.
- Yezostrobus Oliveri**, Stopes & Fuijii, 1910, pp. 33-41, pl. i, figs. 8-9; pl. iii, figs. 10-13; text-figs. 12-14. Ibid.
- Yuccites fractifolius**, Saporta, 1894, p. 110, pl. xix, fig. 20 a. Lower Cretaceous(?); Portugal.
- ? *sp.*, Miquel, 1853, p. 53, pl. i, fig. 3. Cretaceous; Belgium.
- Zamia macrocephala**, Lindley & Hutton, 1835, p. 117, pl. cxxv. (= *Zamiostrobus macrocephalus*, Endlicher, 1841.) Greensand; Kent.
- **ovata**, Lindley & Hutton, 1837, p. 189, pl. ccxxvi A. (= *Zamiostrobus ovatus*, Schimper, 1870.) Ibid.
- **sussexiensis**, Mantell, 1843, p. 34; & 1846, p. 51, pl. ii, fig. 1. (= *Pinites sussexiensis*, Carruthers, 1866 B.) Lower Greensand; Sussex.
- **Washingtoniana**, Ward, 1895 A, p. 350, pl. ii, fig. 6. Potomac Formation; Virginia, U.S.A.
- Zamiophyllum Buchianum** (Ettingshausen), Nathorst, 1890, p. 46, pl. ii, figs. 1-2; pl. iii; pl. iv, fig. 2; & Yokoyama, 1894, p. 223, pl. xx, fig. 1; pl. xxii, figs. 1, 2; pl. xxiii, fig. 6; pl. xxvii, figs. 5 a, b; pl. xxviii, figs. 1, 2. (= *Pterophyllum Buchianum*, Ettingshausen, 1852 c, & *Dioonites Buchianus*, Bornemann, 1856.) Neocomian; Japan.
- **Naumanni**, Nathorst, 1890, p. 47, pl. v, fig. 1; & Yokoyama, 1894, p. 225 pl. xxi, fig. 3; pl. xxvi. Ibid.

- Zamiopsis brevipennis**, Richter, 1904, p. 14, pl. i, fig. 9; & 1905, p. 5, pl. i, fig. 9. Senonian; Quedlinburg, Saxony.
- **insignis**, Fontaine, 1889, p. 162, pl. lxii, fig. 3; pl. lxiv, figs. 1, 3; pl. lxx, figs. 4-6; pl. lxvi, fig. 2; pl. lxvii, fig. 7. Potomac Formation; Virginia, U.S.A.
- **laciniata**, Fontaine, 1889, p. 164, pl. lxvi, figs. 1, 5-8. Ibid.
- **longipennis**, Fontaine, 1889, p. 164, pl. lxi, fig. 8. Ibid.
- **petiolata**, Fontaine, 1882, p. 166, pl. lxvi, fig. 3. Ibid.
- **pinnatifida**, Fontaine, 1889, p. 161, pl. lxi, fig. 7; pl. lxii, fig. 5; pl. lxiv, fig. 2; pl. lxvii, fig. 2. Ibid.
- Zamiostrobus elongatus**, Ettingshausen in Reuss, 1854, p. 740 [nomen nudum]. Cenomanian; Moleteín, Moravia.
- **familiaris**, Bronn in Roemer, 1852, p. 50. (= *Conites familiaris*, Sternberg, 1825.)
- **Guerangeri**, Brongniart, 1849 A, p. 111 [nomen nudum]. Cretaceous; France.
- **Loppineti**, Fliche, 1896, p. 139, pl. i, fig. 3, text-fig. 2. Albian; Clermont, France.
- **macrocephalus**, Endlicher, 1841, p. 72. (= *Zamia macrocephala*, Lindley & Hutton, 1835.) Greensand; Kent.
- **ovatus**, Schimper, 1870, p. 203. (= *Zamia ovata*, Lindley & Hutton, 1837.) Greensand; Kent.
- **pippingfordensis**, Unger, 1850 A, p. 300 (described by Fitton, 1836, p. 181, pl. xxii, fig. 10.) Lower Greensand; Sussex.
- **sp.**, Otto, 1854, p. 38, pl. v, fig. 5. Dippoldiswalde, Saxony.
- Zamites acutipennis**, Heer, 1871, p. 1181; & 1874 A, p. 66, pl. xv, figs. 3, 4, 5 a; pl. xvi, fig. 10. Kome Beds; Greenland.
- **affinis**, Schenk, 1871 A, p. 13, pl. iii, fig. 8. Wernsdorfer Beds; Silesia.
- **alaskana**, Lesquereux, 1889, p. 32, pl. x, fig. 4. Neocomian; Alaska.
- **apertus**, Newberry, 1891, p. 199, pl. xiv, figs. 4, 5. Kootanie Formation; Montana, U.S.A.
- **arcticus**, Goeppert, 1866, p. 134, pl. ii, figs. 9, 10. Recorded by Heer from Kome Beds, Greenland.
- **bohemicus**, Velenovsky, 1889, p. 6, pl. iii, figs. 7, 8. (= *Cycas sp.*, Velenovsky, 1887 A.) Cenomanian; Bohemia.
- **borealis**, Heer, 1874 A, p. 66, pl. xiv, figs. 13, 14; pl. xv, figs. 1, 2. Kome Beds; Greenland.
- **brevipennis**, Heer, 1871, p. 1181; & 1874 A, p. 67, pl. xv, figs. 8, 9, 10. Ibid.
- **concinus**, Heer, 1871, p. 1181 [nomen nudum]. Ibid.
- **crassinervis**, Fontaine (*non* Germar), 1889, p. 172, pl. lxix, fig. 4; pl. lxxxiii, fig. 3. Potomac Formation; Virginia, U.S.A.
- **distantinervis**, Fontaine, 1889, p. 172, pl. lxxxiii, fig. 4. Ibid.


- Zamites familiaris**, Corda in Reuss, 1846, p. 86, pl. xlix, figs. 10, 11.
(= *Conites familiaris*, Sternberg, 1825.) Pläner; Bohemia.
- **globuliferus**, Heer, 1882, p. 12, pl. iv, figs. 1-7.
Kome Beds; Greenland.
- **Goepperti**, Schenk (*non* Zigno), 1871 A, p. 11, pl. iii, figs. 6, 6 a.
Wernsdorfer Beds; Silesia.
- **Heeri**, Nathorst MS., figured as *Zamites*, n. sp., Nathorst, 1885,
p. 341, fig. in text. Patoot Beds; Greenland.
- **iburgensis**, Hosius & von der Marek, 1880, p. 214, pl. xlv, fig. 202.
Neocomian; Westphalia.
- (**Dionites**) **Kaufmanni**, Heer, 1877, p. 146, pl. lviii, figs. 21, 22.
Neocomian; Switzerland.
- **montana**, Dawson, 1883, p. 7, pl. i, figs. 6, 6 A.
Kootanie Formation; Canada.
- **montanensis**, Fontaine, 1893, p. 494, pl. lxxxiv, fig. 4. (Re-named
Pterophyllum montanense by Knowlton, 1907.)
Kootanie Formation; Montana, U.S.A.
- **ovalis**, Fontaine, 1889, p. 173, pl. lxxxv, fig. 4; pl. clxx, fig. 3.
Potomac Formation; Virginia, U.S.A.
- **ovatus**, Schenk (*non* (Lindley & Hutton), Morris), 1871 A, p. 12,
pl. iii, figs. 7, 7 a. Wernsdorfer Beds; Austrian Silesia.
- **pachyneurus**, Schenk, 1871 A, p. 12, pl. iv, fig. 1. Ibid.
- **Schlotheimii**, Presl in Sternberg, 1838, p. 200. (= *Cycadites*
zamiaefolius, Sternberg, 1825.) Quadersandstein; Hör.
- **speciosus**, Heer, 1874 A, p. 64, pl. xiv, figs. 1-12; pl. xvi, fig. 4.
Kome Beds; Greenland.
- **subfalcatus**, Fontaine, 1889, p. 173, pl. lxxxiv, fig. 13; pl. lxxxv,
fig. 3. Potomac Formation; Virginia, U.S.A.
- **tenuinervis**, Fontaine, 1889, p. 171, pl. lxvii, fig. 1; pl. lxix, fig. 2;
pl. lxx, fig. 1; pl. lxxv, fig. 3; pl. lxxvi, fig. 7; pl. lxxxviii, fig. 6;
pl. lxxxiv, fig. 7. Potomac Formation; Virginia, U.S.A.
- **Vanhöffeni**, Engelhardt in Vanhöffen, 1897, p. 371, text-fig. 28.
Kome Beds; Greenland.
- **Weedii**, Fontaine in Weed & Pirsson, 1898, p. 481.
Lower Cretaceous; Montana, U.S.A.
- **sp.**, Dawson, 1886, p. 7, pl. i, fig. 4. Kootanie Formation; Canada.
- ? **sp.**, Fontaine in Ward, 1899 B, p. 666, pl. clxii, fig. 15.
Lower Cretaceous; Wyoming, U.S.A.
- **sp.**, Fontaine, 1889, p. 173, pl. lxxxiv, fig. 12.
Potomac Formation; Virginia, U.S.A.
- **sp.**, Lesquereux, 1892, p. 26, pl. i, fig. 8.
Dakota Group; Kansas, U.S.A.
- **sp.**, Schenk, 1871 A, p. 13, pl. iii, fig. 11.
Wernsdorfer Beds; Austrian Silesia.
- Zingiberites pulchellus**, Heer, 1871, p. 1183; & 1874 A, p. 105,
pl. xxvii, fig. 12 b. Atane Beds; Greenland.

- Zizyphus Beckwithii**, Lesquereux, 1883, p. 125, pl. xix, fig. 5.
Laramie Formation; Colorado, U.S.A.
- **cliffwoodensis**, Berry, 1907 A, p. 676, text-fig. 5.
Magothy Formation; New Jersey, U.S.A.
- **dakotensis**, Lesquereux, 1892, p. 167, pl. xxxvi, figs. 4-7.
Dakota Group; Kansas, U.S.A.
- **elegans**, Hollick, 1894 A, p. 58, pl. clxxvii, figs. 9, 10.
Cretaceous; Long Island, U.S.A.
- **groenlandicus**, Heer, 1883 A, p. 42, pl. lxii, fig. 20.
Patoot Beds; Greenland.
- **Lewisiana**, Hollick, 1894 A, p. 58, pl. clxxx, fig. 13.
Cretaceous; Long Island, U.S.A.
- **oblongus**, Hollick, 1906 A, p. 92, pl. xxiv, figs. 9, 10.
Middle Cretaceous; Long Island, U.S.A.
- Zonarites digitatus** (Brongniart), Geinitz. (= *Fucoides digitatus*, Brongniart.) Recorded from American Cretaceous by Lesquereux, 1873, p. 421; & 1874, p. 44, pl. i, fig. 1.
- Zonopteris comptoniaefolia**, Debey, 1848, p. 117 [nomen nudum].
(= *Didymosorus comptonifolius*, Debey & Ettingshausen, 1859 B.)
Senonian; Aix, Rhenish Prussia.
- **digitatus** (Brongn.), Geinitz. Recorded from Dakota Group, Kansas, by Lesquereux, 1873, p. 421.
- **Goepperti**, Debey, 1848, p. 117; & Debey & Ettingshausen, 1859 B, p. 213, pl. iv, figs. 11-20. Senonian; Rhenish Prussia.
- **Goepperti heteropleura**, Debey, 1848, p. 117 [nomen nudum].
Ibid.
- Zoophycos Brianteus**, Massalongo, 1855, p. 51, pl. iii, figs. 1, 2.
Upper Cretaceous.
- **emarginatus**, Cocchi. See Sacco, 1888, p. 186.
- **Targionii**, Savi & Meneghini. See Sacco, 1888, p. 186.
- **tenuestriatus** (Heer). See Sacco, 1888, p. 186.
- **Villæ**, Massalongo, 1855, p. 49, pl. ii, figs. 1, 2. Upper Cretaceous.
- Zosterites Agardhianus**, Brongniart, 1828, p. 115.
Cretaceous; Sweden.
- **æquinervis**, Debey, 1849, p. 299 [nomen nudum].
Senonian; Aix, Rhenish Prussia.
- **angustifolius**, Ettingshausen, 1893, p. 137; & 1895, p. 13, pl. i, fig. 2. Cretaceous; Australia.
- **Bellovisana**, Brongniart, 1824, p. 317, pl. xxi, fig. 7.
Greensand; Isle of Aix, France.
- **cauliniaefolia**, Brongniart, 1828 A, p. 204. Ibid.
- **elongata**, Brongniart, 1824, p. 317, pl. xxi, fig. 6. Ibid.
- **lineata**, Brongniart, 1824, p. 318, pl. xxi, fig. 8. Ibid.
- **Loryi**, Fliche, 1902, p. 122, pl. ii, one text-fig.
Senonian; Dévoluy, France.
- **Miqueli**, Debey MS. in Mourlon, 1881, p. 133 [nomen nudum].
Senonian; Limburg, Belgium.

- Zosterites multinervis**, Debey, 1848 A, p. 119 [nomen nudum].
 Senonian; Aix, Rhenish Prussia.
- **Orbigniana**, Brongniart, 1824, p. 317, pl. xxi, fig. 5.
 Greensand; Isle of Aix, France.
- **Orbignyanus**, Bronn in Bronn & Roemer, 1852, p. 49. (= *Zosterites Orbigniana*, Brongniart, 1824, & *Z. cauliniæfolia*, Brongniart, 1828 A.)
- **vittata**, Debey, 1848, p. 119 [nomen nudum].
 Senonian; Aix, Rhenish Prussia.

ADDENDA ET CORRIGENDA.

- Page 50. Under *Abietites cretacea*, for "Dakota" read "New Mexico."
50. Under *Abietites Glueckii*, add Richter, 1904, p. 19, pl. i, fig. 14.
 Senonian; Quedlinburg, Saxony.
66. Add *Asplenium distans*, Heer, recorded Dawson, 1886, p. 5,
 pl. iii, fig. 7.
 Kootanie Formation; Canmore, Rocky Mts., Canada.
71. Add *Dignonia Westerhausiana*, Richter, 1904, p. 20, pl. ii,
 figs. 1-5.
 Senonian; Quedlinburg, Saxony.
78. For "*Cassia præ-mennonnia*" read "*Cassia præ-memnonia*."
80. Under *Cedroxylon manehildense* for "xc" read "xv."
113. Add *Dryopteris parvifolia* (Fontaine), Knowlton, 1898, p. 92; &
 in Ward, 1905, p. 541, pl. cxiv, fig. 7. (= *Aspidium parvifolium*,
 Fontaine, 1889.)
 Potomac Formation; Virginia, U.S.A.
115. Under *Equisetum Zeilleri*, add Richter, 1904, p. 18, pl. i, figs. 2
 & 12.
 Senonian; Quedlinburg, Saxony.
125. Under *Geinitzia microcarpa*, for "1905" read "1904."
138. Under *Laurus crassinervis*, add Dawson, 1886, p. 10, pl. iii,
 figs. 3, 3A.



DESCRIPTIVE CATALOGUE OF CRETACEOUS PLANTS.

Group THALLOPHYTA.

Plants with great variety of external morphology, but without differentiation into true root, stem and leaf. The plant body may consist of a single cell, a small number of cells, or may be a large, complex organism many feet in length. The higher members of the group show both internal and external structures suggestive of some of the features of the Vascular plants, but they are distinguished from them by the absence of true roots and the lack of differentiated vascular elements. Reproduction may be by simple, unisexual spores, but in the majority of forms there is some trace of sexuality, and in some the reproductive processes are exceedingly complex.

As no member of the group produces woody or sclerenchymatous elements, the parts are all soft and are very liable to decay. This perhaps accounts for their scarcity among the fossils, though many purely physical appearances have been mistaken for them. Few reliable determinations of members of this group have been made, except among the Calcareous Algæ, which secrete for themselves a coating of carbonate of lime.

Class ALGÆ.

Plants with vegetative body varying from a single cell to a complex, multicellular structure. All forms are provided with chlorophyll, which is masked by other pigments in some of the groups.

Among living Algæ the majority of the larger forms are marine and the group preponderates largely in the sea-water flora as a whole. It is, therefore, all the more surprising that so few good fossil examples are recorded. That Algæ existed from Palæozoic times is rendered certain by the fact that one of the earliest known plants with its internal structure preserved is an alga, *Nematophycus* (Dawson), Carruthers. The theory that all terrestrial vegetation originated from aquatic forms in the earliest times, is also generally accepted. The mere delicacy of the tissues forming the Algæ is not a sufficient explanation for the remarkable poverty of their petrification, for equally delicate parts of the higher plants are preserved; and it is probably due to the chemical nature of their cell-walls that they are so ill represented in fossil floras.

Although large numbers of appearances in the rocks have been described which are not Algæ but are of purely physical origin, many writers have hastily discarded specimens which merit retention because of the unsatisfactory nature of most fossil Algæ. Seward (1894) has discarded all the numerous generic names given to algal impressions and proposed the comprehensive generic name *Algites* for everything. For the Cretaceous plants this course does not seem advisable, because a number of the described species are at least as good as many of the fossil fern-species which are retained by all palæobotanists, while at the same time it is recognised that their affinities are very imperfectly known.

The literature dealing with fossil Algæ is extensive, and much of it is highly controversial. In a Swedish paper in 1874 Nathorst showed how many of the fossil "algæ" were simply physical markings, and he continued his work in 1881. To this Saporta replied in 1882 with a large monograph on the fossil Algæ. Nathorst published a more exhaustive paper in 1886 in reply both to Saporta and the others who continued to maintain the algal nature of the remains. In 1895 Fuchs contributed a large paper on "Fucoiden und Hieroglyphen", and stated that after an exhaustive examination of all the "fucoids" in most museums, he did not discover a single carbonised specimen, and he denied the statements made by other workers that such specimens are frequent. His opinion was that all the so-called

fucoids of the Flysch were of physical origin or due to animal tracks. Rothpletz more recently (1896) has taken a moderate view, and maintains the algal nature of many of the species. These papers are not confined to a consideration of the Cretaceous Algæ, but deal with the whole range of forms.

The Calcareous Algæ are naturally, by reason of their hard encrustations, of most importance geologically, and they do really play a part as "rock builders" (see Seward, 1894 B). Several undoubted genera and species of this group have been described from Cretaceous rocks.

Order DIATOMACEÆ.

[Living family.]

Minute unicellular Algæ sometimes living in colonies, but generally free swimming. The shape of the cell is exceedingly various, and particularly characterised by the thickened cell-wall which is in two parts, fitting together like a pill-box and its lid. This shell is generally silicified and often flattened and ornamented. In the living cell the chloroplast has a yellowish-brown colour.

Fossil diatoms are recorded in enormous masses in several Tertiary and a few Cretaceous deposits. In some cases, indeed, a fine whitish earth consists almost entirely of myriads of the minute silicified tests.

Ehrenberg (1841) pointed out long ago that many species of diatoms existed in the Upper Cretaceous, several apparently identical with living species, and all belonging to still living genera. He gives a list (p. 119) of the species and their localities which is too long for quotation, but a few names selected from it will indicate the distribution and character of the forms he records:—

Amphitetras antediluviana, from the Chalk Marl of Oran,
Coscinodiscus Argus, from the Chalk Marl of Caltanissetta,
Coscinodiscus eccentricus, from the Chalk Marl of Oran,
Coscinodiscus Patina, from the Chalk Marl of Zante,
Fragilaria rhabdosoma, from the White Chalk of Gravesend,

Fragilaria striolata, from the White Chalk of Gravesend,
Gallionella aurichalcea, from the White Chalk of Rügen,
Gallionella sulcata, from the Chalk Marl of Caltanissetta,
Navicula ventricosa, from the Chalk Marl of Oran.

Cayeux (1892, 1897) has recorded numerous diatoms from the Upper Cretaceous zones of *Ammonites mammillaris* and *A. inflatus* in the Paris basin. In his specimens the silicified tests are sometimes replaced by calcareous matter. Hitherto there are no authentic records of diatoms earlier than the Upper Cretaceous.

Order SIPHONEACEÆ.

[Living family.]

Unicellular Algæ of very various size and external form. The thallus may be much branched and is often of considerable size, but it is not separated by cell-walls into distinct cells. Many of the genera are encrusted with a calcareous coating, and the majority of them are marine.

The Siphoneaceæ are the most important group of fossil Algæ, and a number of well preserved and microscopically studied forms have been described by Rothpletz, Steinmann, and others from the calcareous members of the order. The soft-celled, microscopic forms belonging to this affinity have not been recorded among fossils.

Suborder CODIACEÆ.

Genus **BOUEÏNA**, Toula.

[Toula, Sitzb. k. Akad. Wiss. Wien, math.-naturwiss. Cl. 1884, vol. 88, p. 1319. For the recognition of its plant-nature see Steinmann, Bericht. naturf. Ges. Freiburg, vol. 11, pp. 1-6, text-figs. 1-7.]

The genus is monotypic and cannot therefore be diagnosed apart from the species. It was described by Gümbel, Zittel and others as a sponge, and the reference of the fossil to the Algæ is due to Steinmann.

This genus, which shows such similarity to the living *Halimeda*, is of special interest because previous to its recognition no member of the Codiaceæ had been determined on the basis of its structure, prior to the Tertiary.

***Boueina Hochstetteri*, Toul.**

1884. *Boueina Hochstetteri*, Toul., Sitzb. k. Akad. Wiss. Wien, math.-naturwiss. Cl., vol. 88, pp. 1319-1324, pls. vii-ix.

1901. *Boueina Hochstetteri*, Steinmann, Bericht. naturf. Ges. Freiburg, vol. 11, pp. 1-6, text-figs. 1-7.

In this alga the simple, unbranched body of the calcareous thallus is 10-20 mm. long, 2-3.5 mm. thick, and is penetrated by a system of branching canals which are connected without any visible segmentation. The cross-section is circular or oval, and in the central region the little branched canals of the axis run more or less nearly parallel, in the periphery they are very much smaller and stand at right angles to the surface and are very freely dichotomously branched. The ultimate branches are arranged very close together and form a kind of cortical zone. In large specimens the central canal-system is replaced by a simple cavity. Between the canals is an undifferentiated calcareous skeleton.

HORIZON.—Upper Neocomian.

LOCALITY.—Servia.


The general habitat of this Neocomian member of the Codiaceæ appears to have been similar to that of the modern *Halimeda*.

Steinmann's figures (Steinmann, 1901, text-figs. 2 and 3), reproduced here as text-fig. 1, show the general character of the cross-section of these algæ and illustrate clearly the remarkable likeness to the living *Halimeda*, a similar section of which is shown in text-fig. 2.


Fructifications for this fossil are not recorded, but this is a point which tends further to strengthen its supposed affinity to *Halimeda*, for in the living genus the spores are not developed on the parts already penetrated by calcareous matter.

It may be remarked that though, as Steinmann points out, there is a great similarity between the fossil and the living genus, the parts compared are only the "stalk" region of the plants and this is similar in several of the Codiaceæ.

A.




B.



Text-fig. 1.—*Bouëina Hochstetteri*, Toulou. A. Transverse section showing the character of the central and the peripheral canals. $\times 28$. B. Cross-section of a specimen with a central cavity, and showing clearly the dichotomous branching of the peripheral canals. $\times 28$. After Steinmann.

Other Codiaceæ.—To the Codiaceæ should perhaps be added the impression described by Lorenz v. Liburnau as *Halimeda Fuggeri* (Lorenz, 1897) from the Flysch. Further specimens of this fossil were described by Lorenz in 1902, and the generic



Text-fig. 2.—Cross-section through a living *Halimeda* for comparison with the fossil *Bouëina Hochstetteri*. $\times 24$. After Steinmann.

name *Halimedes* suggested for them. The specimens certainly have the external appearance of a *Halimeda*, but they exhibit no internal structure, and their plant origin does not appear to be entirely beyond doubt.

Suborder *DASYCLADACEÆ*.

Genus **NEOMERIS**, Lamouroux.

[Living genus.]

A member of the Dasycladaceæ with unbranched, cylindrical stem, thickly encrusted. It consists of a thick-walled stem-cell with no cross walls or constrictions, with densely packed whorls of 32–80 side “branches.” These carry usually one terminal and short-stalked sporangium and two sterile paraphysis-like segments with swollen ends.

Neomeris cretacea, Steinmann.


1899. *Neomeris cretacea*, Steinmann in Felix & Lenk, Beitr. Geol. Paläont. Mexico, p. 200, text-figs. 42-46.

The largest portions of the calcareous cylinder of the fossil are 3-4 mm. long and 2 mm. thick, but they were probably 10 mm. long when alive. They are perfectly cylindrical or slightly compressed. The tubes are penetrated by a central cavity which varies from 1.2-1.4 mm. in diameter. The wall is often as much as 0.5 mm. thick, and is penetrated by numerous radially arranged branches of two kinds, the one simple and uniform, about 0.05-0.06 mm. in diameter, and the other in fewer numbers and of flask-like shape with a blind ending.

HORIZON.—Upper Cenomanian.

LOCALITY.—Mexico.

The form much resembles the Eocene representatives of the genus.



Text-fig. 3.—*Munieria baconica*, v. Hantken. External appearance.

A. Nat. size; B. enlarged. After Deecke.

Genus **MUNIERIA**, v. Hantken *in litt.* (see Deecke).

[Neues Jahrb. f. Min., 1883, vol. 1, p. 9.]

The genus contains the single species *M. baconica*, and is not diagnosed by the original author.


Munieria baconica, v. Hantken in Deecke.

1883. *Munieria baconica*, v. Hantken *in litt.*, Deecke, Neues Jahrb. f. Min., vol. 1, pp. 9-11, pl. i, figs. 4-10.

1897. *Munieria baconica*, Hauptfleisch in Engler & Prantl, Pflanzenfamil., p. 56.

1906. *Munieria cf. baconica*, Oswald, Geol. Armenia, plate to face p. 234.

This alga is cylindrical, from 5-7 mm. long, with a blunt apex. It is composed of numerous symmetrical, similar zones,



Text-fig. 4.—*Munieria baconica*, v. Hantken. A. Cross-section showing *k*, chalky skeleton, *v*, verticillate canals. B. Longitudinal section, *c*, central canal. C. Restoration of a small part of the alga: *c*, axial canal; *v*, verticillate canals which contained the living portion of the alga. The perforated outline represents the calcareous skeleton, *w*, with spaces, *h*, in it. After Deecke.

each about 0.5 mm. thick, so that there are 10-14 of these in a specimen. The diameter is about 0.75 mm., and each zone is perforated by about a dozen round pores. In good specimens

there is no external ornamentation, and, except for the pores, no external sign of zones.

The external appearance of the fossil is seen in numerous cases where it has weathered out, so that hand-specimens are covered with individuals of *Munieria*. The internal structure can be studied by means of sections. In section it is seen that there is an axial canal in the centre of each segment, from which the lateral canals radiate. In longitudinal section they are seen more clearly, and the arrangement of the living tissues is evident (see text-fig. 4).

HORIZON.—Cretaceous.

LOCALITY.—Bakony.

Deecke gives no concise definition of the genus, but describes it at some length and places it near *Gyroporella*. In the neighbourhood of Bakony the alga is so abundant it is said practically to compose one of the beds of the Cretaceous. Hovelacque (1900) gives a figure of a section of fossiliferous calcite from the Urgonian of the Sub-Alps in which are fragments of what he considers may be *Munieria* of Deecke, but his published figures are not conclusive.

Details of the reproductive branches are unknown. In the same matrix Deecke noticed several segments differing somewhat from the normal ones, which he took to be the reproductive branches of the same species.

V. 11063. A small hand-specimen, and a microscopic slide of the same, described as *Munieria* by Oswald (1906). The sketch which he gives (see plate to face p. 234) is unsatisfactory, the actual specimen, while showing much more definite structure than does the figure, being still of such an obscure nature that it is highly doubtful if it is an alga. In the matrix are numerous other fragments (not noted by Oswald) which bear a closer resemblance to a *Lithothamnium* than does his specimen to *Munieria*. No accurate determination seems possible at present.

(?) Urgonian; Armenia. Presented by Dr. F. Oswald
and H. F. B. Lynch, Esq., 1904.

Genus **DIPLOPORA**, Schafhäütl.

[Süd-Bayerns Lethæa Geogn. Petrefacten, 1863, pp. 324, 327.]

The generic name was first given to forms which Schafhäütl included among the Bryozoa. Later by Gümbel and others it was superseded by *Gyroporella*, but the older name was revived for some of the algal forms which are characteristic of the Trias; and the name is also used by Lorenz and Arbenz for the Cretaceous species which they describe as follows:—

Diplopore Mühlbergii, T. Lorenz.

1902. *Diplopore Mühlbergii*, T. Lorenz, Bericht Naturf. Ges. Zürich, vol. 12, pp. 19 [52]–20[53], figs. 3–7.

1908. *Diplopore Mühlbergii*, Lorenz, Arbenz, Vierteljahrsschrift Naturf. Ges. Zürich, vol. 53, pp. 387–392, figs. 1–5.

Slender, somewhat bent tube-like forms, average length about 2 mm. (?) From the hollow axial space open whorls of branch canals. The branches vary, but are principally simple. Branch whorls are separated some distance from each other and the branches are not superimposed in vertical series. In a variety described by Arbenz (1908) the average diameter is 0.3–0.5 mm., with a thickness of 0.1–0.15 mm. for the wall.

HORIZON.—Barremian and Aptian.

LOCALITY.—Wildkirchli and elsewhere, locally very abundant.


Lorenz notes that the discovery of this form in the Cretaceous is of interest, because hitherto algæ of this generic type have been known in numbers only from the Trias, with isolated specimens from the Jura. According to Arbenz (1908) the alga is extremely abundant in the Lower "Schrattenkalk" (Up. Barremian) and Aptian of some districts, and may even form one third of all the organisms in a deposit. He says (p. 392) "Die Diploporen der untern Kreide scheinen somit auf das Barrémien und Aptien in neritischer Facies beschränkt zu sein," where they are a typical facies fossil.

The name *D. Mühlbergii* probably covers several species which can only be separated when more detailed work is available.

Genus **TRIPLOPORELLA**, Steinmann.

[Neues Jahrb. f. Min., 1880, vol. 2, p. 130.]

Steinmann gave no generic diagnosis of this genus, but described it fully, and his original description was much extended



Text-fig. 5.—*Triploporella Fraasi*, Steinmann. A. A portion of the Lebanon Cretaceous rock with specimens of *Triploporella*, natural size. B. A fragment of *Triploporella*, $\times 2$. C. An enlarged fragment showing *a*, the primary branching, *b*, the secondary branch system, *c*, the axis broken across. After Steinmann.

by the examination of further specimens from Mexico (see Steinmann, 1899). According to him *Triploporella* is a member of the Dasycladaceæ of Turonian and Cenomanian age, which had a wide geographical distribution, as is evidenced by the two localities in which it has been found, viz. Syria and Mexico. It has some of the characters both of the Acetabulariæ and the


Dasycladaceæ, and may be considered as a form intermediate between the two groups.

Triploporella Fraasi, Steinmann.

1880. *Triploporella Fraasi*, Steinmann, Neues Jahrb. f. Min., vol. 2, p. 136, pl. v, figs. 1-8.

1899. *Triploporella Fraasi*, Steinmann in Felix & Lenk, Beitr. Geol. Palæont. Mexico, pp. 190-195, text-figs. 29-40.

A form with closely packed series of whorled branches bearing spores, and with finely divided branches below and beyond these. The whole structure probably about 30 mm. long, and



Text-fig. 6. — *Triploporella Fraasi*, Steinmann. Portion of a radial section of some of the whorled branches. s, the axial stem. In the cavities of the "branches" are the "mother spores" dividing into four. After Steinmann.


of a diameter of about 5 mm. at the distal, and 1.5-2 mm. at the proximal end.

From the calcareous specimens of Mexico further details were obtainable, and the verticillate series of sporangial branches contained "spores."

The restoration of the plant given in Steinmann's second paper, reproduced here in text-fig. 7 (p. 246), shows the vertical axis and series of whorled branches, both spore-bearing and divided.

HORIZON.—Turonian and Cenomanian.

LOCALITY.—Lebanon, Syria, and Mexico



Text-fig. 7.—*Triploporella Fraasi*, Steinmann. Reconstruction. $\times 4$.
After Steinmann.

Associated with the Siphonæ is the rather doubtful genus *Goniolina*, first described from the Jurassic, and later recorded in Cretaceous deposits. Saporta originally named it as an inflorescence of a proangiosperm, and its external appearance is suggestive of the spines of an Echinoderm. It is now generally accepted among the Siphonæ, but as it is primarily a Jurassic plant it will not be considered here.

Gyroporella (Gümbel) is another genus recorded in the Cretaceous, but it was first described from Permian rocks. It is outwardly similar to *Neomeris*, but with less complex structure.

Tribe ACETABULARIÆ?

The extremely problematic fossils which have been placed in the genus *Gyrophyllites*, founded by Glocker in 1841 for some Jurassic specimens, have a number of representatives in true Cretaceous deposits, and also in the Flysch. Heer (1877) founds and figures three species (pl. lviii), *G. obtusifolius*, *G. Oosteri*, and *G. pentamerus*, from the Swiss Cretaceous. His genus *Discophorites*, with the species *D. angustilobus* and *D. Fischeri*, shows no characters really to separate it from the previously named "genus," and must be included in "*Gyrophyllites*." Lorenz (1901) considers the "genus" at some length and definitely places it in the Acetabulariæ, although Solms-Laubach, in his classical monograph on the Acetabulariæ (Solms-Laubach, 1901), does not discuss these fossils, but notes their extremely problematic nature in his text-book. Lorenz (*l. c.*) determines several new species from the Flysch deposits.

Order PHÆOPHYCEÆ (?)

[Living family.]

Algæ, almost entirely marine, with a large, branched and complex thallus. Multicellular, with some cell-differentiation. Chromatophores golden brown in living cells.

Genus **CHONDRITES**, Sternberg.

[Flora d. Vorwelt, 1838, p. 25.]

Algæ, or fossils which suggest algæ, with a much-branched dichotomous or sympodial thallus which is *cylindrical*.

The essential feature of the fossils described under this generic name is the cylindrical form of the branches. These are not flattened in the rock like most plant-remains, and they often lack all traces of carbonaceous material. They are frequently

distinguished from the surrounding matrix by their different colour—black in a light rock or white in a dark one.

Brongniart (1828 A) described the genus under the name *Fucoides* in his § *Gigartinites* as follows (p. 20):—"Fronde rameuse, à branches presque-cylindriques, charnues, jamais membraneuses." He divided the group into nine species, to the number of which many later writers have added. Much that was said about the fossil Algæ as a whole (p. 234, *ante*) applies particularly to this section, and it does not appear necessary to enter into an exhaustive recapitulation of all the "species" described, even among the Cretaceous specimens of *Chondrites*. In contrast to most of the earlier writers who tended to multiply genera and species of fossil Algæ, Ettingshausen (1863) pointed out that a number of the so-called Algæ, placed in this genus by Fischer-Ooster and others, cannot be accepted. He also noted the difficulty of distinguishing between many of the forms to which distinct specific names have been given, because of the intermediate stages which occur to connect them.

In the controversy on the algal nature of fossil Algæ, the genus *Chondrites* is one of the most disputed. Nathorst (1886 A) maintains that the forms are not algæ and supports this view with weighty arguments. For example, he points out how the very nature of the matrix, particularly in the Flysch deposits, shows that the beds must have accumulated rapidly, while Algæ of the nature of the supposed "*Chondrites*" do not inhabit troubled waters. He also makes much of the argument that the specimens are not carbonised and show no trace of vegetable substance even when they occur in rocks which contain other true vegetable fossils which are carbonised. This objection, however, seems to be sufficiently answered by the fact that even among living seaweeds the decay takes place in a different manner from that of the higher plants, owing to the different chemical composition of the plant-body in the two types of vegetation.

After having read most of the extensive and highly controversial literature on the subject, and having seen a large number of specimens in most of the leading museums, my conclusion is that the "genus" *Chondrites* does contain "species" which are truly the remains of algæ, though many of those described for the genus are probably purely physical phenomena.

And further, that if this conclusion should ultimately prove incorrect and all of them prove to be other than algæ, it is still useful to retain the genus for the present, and wise to keep it in the established position which is as good as any other on *a priori* grounds, and has the advantage of being long recognised.

In his exhaustive monograph on the Algæ of the Flysch Rothpletz (1896) proposes the generic name *Phycopsis* to replace the old-established *Chondrites*. His objection to the latter is very just, namely, that it suggests affinities which are not proven. His own name, however, is not free from the same objection. As he brings forward no new facts which throw light on the affinity of the plant, by the code of nomenclature the older name should be retained. According to my suggestion regarding the names of doubtful species, the name is printed in gothic type*.

The Flysch deposits which are so peculiarly rich in "Algæ" have been the subject of a monograph by Rothpletz (1896) to which reference should be made. They will not be specially treated in the present volume, and indeed most of them are Tertiary, or doubtfully Cretaceous in age.

Chondrites Targionii (Brongniart), Sternberg.

1828. *Fucoides Targionii*, Brongniart, Hist. Végét. Foss., p. 56, pl. iv, figs. 4-6.

1828. *Fucoides Targionii*, Brongniart, Prodrôme, p. 20.

1835 (read 1828). *Fucoides Targionii*, Mantell, Trans. Geol. Soc. [2] vol. 3, p. 210.

* In a paper in the Annals of Botany for October 1911, I brought forward the arguments that form the foundation of my suggestion—which is, that all fossil plants for which there is no good, scientific reason for association with given families and genera, and to which nevertheless names indicative of such affinities have been given, should be printed henceforth in Gothic character. This would instantly indicate the doubtful nature of the fossil without altering and multiplying the nomenclature, and would avoid the use of a trinomial system or any of the other more cumbrous devices which have been suggested. It would also immediately appeal to those not specially trained in palæobotany when looking through lists of names, and set them on their guard against drawing unwarranted conclusions.

1838. *Chondrites Targionii* et varr., Sternberg, Versuch Flora Vorwelt, p. 25, pl. ix, fig. 3.
1853. *Chondrites Targionii*, Unger, Denkschr. k. Akad. Wiss. Wien, vol. 4, p. 79, pl. xxv, fig. 5.
1853. *Chondrites Targionii* (excl. some of Sternberg's varr.), Fischer-Ooster, Fossilen Fucoiden, p. 46, pl. viii, figs. 8 a, 8 b.
1863. *Chondrites Fındobonensis*, varr., Ettingshausen, Fossilen Algen Wiener Karpathen-Sandsteines, p. 457.
1869. *Chondrites Targionii*, Schimper, Traité Paléont. Végét., vol. 1, p. 170, pl. iii, fig. 7.
1877. *Chondrites Targionii* (et varr.), Heer, Flora Foss. Helvetiæ, p. 155; pl. lx, fig. 5; pl. lxi, fig. 9; pl. lxii, figs. 1-10; pl. lxiii, figs. 6 a, 12-17.
1880. *Chondrites Targionii*, Hosius & v. d. Marck, Palæontogr., vol. 26, p. 130, pl. xxiv, figs. 1, 2.
1896. *Phycopsis Targioni*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 48, p. 887.
1900. *Chondrites Targioni*, Zeiller, Élémt. Palæobot., p. 33, fig. 6.


Heer's diagnosis is perhaps the most concise:—" *Chondrites* fronde irregulariter dichotome et pinnatim ramosa, ramis alternis et oppositis, hinc inde valde approximatis .75-1.5 mm. latis, longitudine valde inaequalibus, ramulis nonnullis praelongis, angulo acuto egredientibus, erectis, strictis."

HORIZON.—Lower Greensand—Flysch.

LOCALITY.—Widely distributed over Europe.

Many of the specimens which have been referred to this species are certainly worthless, and are probably worm-burrows or animal-tracks of the kind demonstrated by Nathorst. Further, a number of specimens have been placed in subspecies or varieties which show no real features distinctive enough to separate them from the "species" which is recognised of little scientific value in any case. There are, nevertheless, a number of fossils which can at once be recognised as characteristic and which it is useful to place together under one well-established name.

The specimens in the Museum are principally from the Lower Greensand of Bignor Park, one of the earliest known localities for the fossil, which Mantell described and figured in several of his works. The thallus is whitish and highly calcareous, and the specimens are less suggestive of plants than many of the others included in the species, though they come within the terms of the diagnosis.



Text-fig. 8.—*Chondrites Targionii* (Brongn.), Sternberg. A characteristic specimen of the "species." Nat. size. After Brongniart.

V. 88, V. 88 a. Numerous ramifications of the branched "thallus" penetrating the sandy matrix in various directions. The branches are white in colour and cylindrical. In section they appear as circles scattered in the matrix. Lower Greensand; Bignor Park, Sussex. *Egerton Coll.*, 1882.

5570, 5573, 5574, 5575, 5576, 5577, 5579, 5579 a. Similar specimens from the Greensand. *Mantell Coll.*

10780, 9424, 9426, V. 11507, V. 11508, V. 11509, V. 11510, V. 11511, V. 11512. Similar specimens from the Greensand; Bignor (?). *Mantell Coll. (?)*

Chondrites intricatus (Brongniart), Sternberg.

[Plate I, fig. 1 b.]

1824. *Fucoides intricatus*, Brongniart, Observations sur *Fucoides*, p. 311, pl. xix, fig. 8.
 1828. *Fucoides intricatus*, Brongniart, Hist. Végét. Foss., p. 59, pl. v, figs. 6, 7, & 8.
 1828. *Fucoides intricatus*, Brongniart, Prodrôme, p. 20.
 1838. *Chondrites intricatus*, Sternberg, Versuch Flora Vorwelt, p. 26, pl. vii (non vi), fig. 3.
 1858. *Chondrites intricatus*, Fischer-Ooster, Fossilen *Fucoiden*, p. 44, pl. viii, fig. 1.
 1863. *Chondrites Vindobonensis*, varr., Ettingshausen, Fossilen Algen Wiener Karpathen-Sandsteines, p. 457.
 1869. *Chondrites intricatus*, Schimper, Traité Paléont. Végét., vol. 1, p. 172, pl. iii, fig. 9.
 1877. *Chondrites intricatus* (et varr.), Heer, Flora Foss. Helvetiæ, p. 157, pl. lxiii, figs. 1-10.
 1896. *Phycopsis intricata*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 48, p. 888, pl. xxii, fig. 7. (*Chondrites æqualis* should probably be included here.)

Heer's diagnosis may be quoted:—" *Chondrites fronde subtili, caespitosa, tripinnatim ramosa, ramulis .25-.5 mm. latis, longitudine inaequalibus, setaceis, angulo acuto egredientibus, approximatis.*"

HORIZON.—Cretaceous—Flysch.

LOCALITY.—Widely distributed in the Swiss Alps and elsewhere.

Ettingshausen (1863) concluded that this species is really a younger form of *Chondrites Targionii*, but Rothpletz's more recent and more detailed work does not support this view. The species was one of the earliest to be recognised among fossil algæ, and its characteristic size and appearance make it easily distinguished from most of the other "species" of supposed algæ.


In our Pl. I, fig. 1 a slab of Cretaceous Flysch shale is illustrated, showing numerous fragments (b) of the plant with other algæ.

53020. A number of fragments of the tufted, branching thallus, together with an *Algites*, figured in Pl. I, fig. 1 b. Cretaceous Flysch ; Sievering.

Ettingshausen Coll., 1879.

- V. 11513. Several fragments of the branching thallus, one showing the tufted form very well. Cretaceous Flysch ; Sievering.

Ettingshausen Coll., 1881.



Text-fig. 9.—*Chondrites intricatus* (Brongn.), Sternberg.

A characteristic specimen. Nat. size. After Heer.

- V. 11514, V. 11515. Numerous branches and fragments of thallus lying in various directions and penetrating the slab of matrix. Cretaceous Flysch ; Sievering.

Ettingshausen Coll., 1881.

Chondrites patulus, Fischer-Ooster.

[Plate I, fig. 2.]

1858. *Chondrites patulus*, Fischer-Ooster, Foss. Fucoiden, p. 48, pl. viii, figs. 6, 7.

1877. *Chondrites patulus*, Heer, Flora Foss. Helvetiæ, p. 156, pl. lxiii, figs. 18-24.

Heer's diagnosis may be quoted:—" *Chondrites fronde erecta, pinnatim vel bipinnatim ramosa, ramis approximatis, sub angulo fere recto egredientibus, inaequilongis, .75-1 mm. latis.*"

HORIZON.—Cretaceous—Flysch.

LOCALITY.—Fähnern, etc., Switzerland.

This "species" is readily separated from the others by the arrangement of the lateral branches at right angles to the larger branches.

V. 11516. In Pl. I, fig. 2, a small fragment from the Upper Cretaceous is illustrated to show the branched thallus, which much resembles the original description.



Text-fig. 10.—*Chondrites patulus*, Fischer-Ooster. Showing the rectangular form of the branches. Nat. size. After Fischer-Ooster.

Genus **ALGITES**, Seward (emend.).

[Catal. Mesoz. Plants Brit. Mus., Wealden, pt. i, 1894, p. 4.]

Seward's diagnosis, if emended, may be taken:—A generic term to cover a presumably composite genus for those *flattened* impressions "which in all probability belong to the class Algæ, but which by reason of the absence of reproductive organs, internal structure, or characters of a trustworthy nature in the determination of affinity, cannot be referred with any degree of certainty to a particular genus or family."

The separation of the cylindrical forms into another group may at times seem arbitrary, but it is convenient at present. Where all the "genera" are based on very unsatisfactory data, it is well to use a clearly marked and stable character.

Algites furcatus (Brongn.), comb. nov.

[Plate I, fig. 1 a.]

1824. *Fucoides* (*Sphaerococcus*?) *furcatus*, Brongniart, Observations sur *Fucoides*, p. 309, pl. xix, fig. 3.
 1828. *Fucoides furcatus*, Brongniart, Hist. Végét. Foss., p. 62, pl. v, fig. 1.
 1828. *Fucoides furcatus*, Brongniart, Prodrôme, p. 20.
 1838. *Chondrites furcatus*, Sternberg, Versuch Flora Vorwelt, p. 27.
 1858. *Chondrites furcatus* (some varieties), Fischer-Ooster, Fossilen *Fucoiden*, p. 51, pls. ix-x.
 1863. *Chondrites affinis* & *C. genuinus*, Ettingshausen, Foss. Algen Wiener Karpath.-Sandst., pp. 460, 465.
 1869. *Chondrites furcatus*, Schimper, Traité Paléont. Végét., vol. 1, p. 169, pl. iii, fig. 8.
 1896. *Phycopsis affinis*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 48, p. 885, pl. xxii, figs. 1, 2.

Algites with thallus branching alternately, the side branches averaging 3 mm. in width and with rounded ends, springing from a larger branch 4-7 mm. in diameter. The main branches are characterised by dichotomous division, and the ultimate branches are sympodial.

HORIZON.—Cretaceous—Flysch.

LOCALITY.—Switzerland and other mid-European countries.

Rothpletz states that it is the fucoid with the widest thallus among those of the Flysch deposits. It appears to me to be one of the best established of the doubtful group.

53020. A branching thallus showing some of the ultimate ramifications with their rounded ends, illustrated in Pl. I, fig. 1 a, with *Chondrites intricatus*. Cretaceous Flysch; Sievering. Ettingshausen Coll., 1879.

Algites sp.

41410. Very obscure markings on a block of white chalk, which may possibly be fucoidal impressions. Upper Chalk; Charlton, Kent. Purchased, 1863.

Algites ?

- V. 3947. Unidentifiable fragments of plant tissue in a mass together. On sandstone. They may, however, be macerated fragments of higher plants. Senonian; Baumberg, Westphalia. Purchased, 1899.

NOTE.—Several more or less doubtful species which could technically be placed in the genus *Algites* have been described from time to time. Their bibliographic details, however, are not worth bringing together. Many of the "species" will be found in the list of Cretaceous species given at the beginning of this volume.

Order RHODOPHYCEÆ—CORALLINACEÆ.

Algæ, principally marine, with branched and sometimes very complex thallus. Multicellular, with some cell-differentiation and a more highly organised reproductive system than in the other groups of Algæ. The calcareous encrusting of the thallus and the form of growth give a number of the species the appearance of corals. The chromatophores (as in all Rhodophyceæ) contain a red colouring-matter.

Genus **LITHOTHAMNIUM**, Philippi.


[Living genus.]

A coralline Alga with an encrusting, spreading thallus growing on to the sub-stratum, with irregular, coralline, or branching upward outgrowths. Thallus entirely calcified and as hard as stone. Basal part of thallus with a regular arrangement of the cells in rows. Conceptacles developed in the upward growing part of the thallus, generally in series of growth-zones.

As the living genus was long unrecognised as belonging to the Algæ, it is not surprising that fossil representatives of the group remained even longer without detection. It was in 1874 that the first systematic description of the fossil forms of *Lithothamnium* appeared by Gumbel under the title "Die sogenannten Nulliporen." In this work seven Cretaceous species are described in addition to those of Tertiary and Jurassic age. Rothpletz (1891 B), under the title "Fossile Kalkalgen aus den Familien der Codiaceen und der Corallineen," has added the only other important contribution to the subject, in which he describes four new Cretaceous species of *Lithothamnium*. Solms-Laubach in his 'Text Book of Fossil Botany' says that "we shall do well . . . to put them all together as

Lithothamnium ramosissimum," but he wrote before the more detailed measurements and study of the fossils had been undertaken. Seward (1898) gives some of Rothpletz's figures and describes *L. mamillosum*, Gümbel, in his more recent Text-book.

Lithothamnium has not hitherto been recognised in British Cretaceous rocks, although it is so widely spread in European deposits of the same age. Though no specimens have been identified in the British Museum Collection, I have had the privilege of examining Prof. Rothpletz's original specimens in Munich. In several of these slides the internal structure is remarkably clear, and there appear to be several true species among those described from the Cretaceous, even if all the eleven species are not really distinct.



Text-fig. 11.—*Lithothamnium mamillosum*, Gümbel, to show external features. Nat. size. After Gümbel.

***Lithothamnium mamillosum*, Gümbel.**

1871. *Lithothamnium mamillosum*, Gümbel, Abhandl. k. Akad. Wiss. München, vol. 11, p. 41, pl. ii, figs. 7 a, 7 b.

1891. *Lithothamnium mamillosum*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 43, p. 315, pl. xvii, fig. 7.

1898. *Lithothamnium mamillosum*, Seward, Fossil Plants, p. 188, text-fig. 32 A, p. 155.

Algal thallus irregularly branched, nodular and crust-like, the branches about 5 mm. long and 4.5 mm. thick. In microscopic section the individual cells of the perithallium are 5.5–8 μ in diameter and of square outline, and of the hypothallium about 18 μ in length.

Apparently sterile.

As text-fig. 11 shows, this irregular coralline mass may well have been mistaken for an animal nullipore. Previous to

Gümbel's work, it had been described as *Cellepora bipunctata*, Goldf., *Membranipora bipunctata*, Blainv., *Discopora bipunctata*, Edw., and *Marginaria bipunctata*, Roem.

The internal structure was studied in more detail by



× 80

Text-fig. 12.—*Lithothamnium mamillosum*, Gümbel, to show microscopic details of the tissue of the perithallium. × 80. After Rothpletz.

Rothpletz (1891 B), in the hope of detecting reproductive organs, but the available specimens seem to be sterile.

HORIZON.—Uppermost Cretaceous (Danian).

LOCALITY.—Petersberg, near Maestricht.

TYPE.—Goldfuss' specimen of *Cellepora bipunctata* in the Palæontological Museum, Old Academy, Munich.



Text-fig. 13.—*Lithothamnium perulatum*, Gümbel. External appearance. Natural size. After Gümbel.

***Lithothamnium perulatum*, Gümbel.**

1871. *Lithothamnium perulatum*, Gümbel, Abhandl. k. Akad. Wiss. München, vol. 11, p. 44, pl. ii, figs. 11 a, 11 c.

Thallus somewhat flat, spreading and irregularly encrusting. In section the cells are rectangular and $10\mu \times 8\mu$ in diameter.

HORIZON.—Uppermost Cretaceous (Danian).

LOCALITY.—Maestricht.

TYPE.—Bavarian State Collection, Old Academy, Munich.


***Lithothamnium procoenum*, Gumbel.**

1871. *Lithothamnium procoenum*, Gumbel, Abhandl. k. Akad. Wiss. München, vol. 11, p. 43, pl. ii, fig. 10.

The thallus with irregularly divided branches about 3–4 mm. in diam. springing from a thicker stock. In section the cells are rectangular and about $12\mu \times 8\mu$.

The specimen was originally known as the Nullipore *Ceriopora dichotoma*, Goldf.

It is probable that the species *Lithothamnium parisiense*, Gumbel, described from a small fragment, and separated from



Text-fig. 14.—*Lithothamnium procoenum*, Gumbel. External form.
Natural size. After Gumbel.

this because its cells are $9\mu \times 6\mu$ in diameter, really belongs to this species.

HORIZON.—Uppermost Cretaceous (Danian).

LOCALITY.—Petersberg, near Maestricht.

TYPE.—Palæontological Museum, Old Academy, Munich.

***Lithothamnium palmatum* (Goldf.), Gumbel.**

1826. *Nullipora palmata*, Goldfuss, Petrefacta Germaniæ, p. 20, pl. viii, fig. 9.

1871. *Lithothamnium palmatum*, Gumbel, Abhandl. k. Akad. Wiss. München, vol. 11, p. 284, pl. v, figs. L 1 a & L 1 b.


Irregularly branched, the branches much grown together, and the ends of the branches somewhat swollen. In section the cells are rectangular and $7\mu \times 8\mu$.

HORIZON.—Upper Cretaceous.

LOCALITY.—Gosau, and in France.

TYPE.—Goldfuss' *Nullipora palmata*, in the University Collections at Bonn.

It is probable that *Lithothamnium racemosum* (Goldf.), Gumbel, p. 284, pl. d iv, figs. L 2 a & L 2 b, is really a small



Text-fig. 15.—*Lithothamnium palmatum* (Goldf.), Gumbel. External form. Natural size. After Gumbel.

fragment of this species. The size of the cells, viz. $9\mu \times 10\mu$ instead of $7\mu \times 8\mu$, is the only apparent difference and hardly seems sufficient basis for a specific distinction where the internal details are so slightly described.

***Lithothamnium Goldfussi*, Gumbel.**

1871. *Lithothamnium Goldfussi*, Gumbel, Abhandl. k. Akad. Wiss. München, vol. 11, p. 285, pl. d iv, figs. L 3 a & L 3 b.

A large, spreading form with wing-like expansions and short branches with rounded ends. The cells are exceptionally large, and measure $70\mu \times 24\mu$.


LOCALITY.—In the Cretaceous deposits, apparently several localities.

TYPE.—Goldfuss' specimens in the University Collections of Bonn.


***Lithothamnium cenomanicum*, Rothpletz.**

1891. *Lithothamnium cenomanicum*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 43, p. 313, pl. xv, figs. 1, 2, 16.

The thallus consists of rounded, mamillöse masses which are outwardly quite distinct from the other species of *Lithothamnium*.



Text-fig. 16.—*Lithothamnium Goldfussi*, Gümbel. External form.
Natural size.



Text-fig. 17.—*Lithothamnium cenomanicum*, Rothpletz. A. External appearance, natural size. B. Cross section showing a zone of tetraspore development, $\times 2$ After Rothpletz.

The growth is very regular and there is no growing over of the perithallium through the hypothallium. In section the cells are $12-14\ \mu \times 20-25\ \mu$. The tetraspores lie in a number of concentric zones in the tissues and are $50-60\ \mu$ by $70-80\ \mu$.

HORIZON.—Cenomanian.

LOCALITY.—St. Patern, Sarthe, France.

TYPE.—Palæontological Museum, Old Academy, Munich.

Rothpletz considers that this species should be placed near *L. amphiroæformis*, *L. nummuliticum*, *L. ramosissimum*, and *L. racemus*, because of the size of the cells, though in its outward form it differs from these.

***Lithothamnium turonicum*, Rothpletz.**

1891. *Lithothamnium turonicum*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 43, p. 313, pl. xv, figs. 9, 13.

Small, dichotomously branching form, the branches 3 mm. thick by 12 mm. long. When broken across the concentric banding is visible to the naked eye, and these concentric rings are the tetraspore zones. Cells $9-10\ \mu$ by $12-15\ \mu$, and in the hypothallium up to $30\ \mu$ long. The tetraspores are $30-35\ \mu$ wide and $75\ \mu$ high.

HORIZON.—Turonian.

LOCALITY.—Beausset, Var, France.

TYPE.—Palæontological Museum, Old Academy, Munich.


Both the external shape and the internal cells separate this immediately from *L. cenomanicum*, while the other forms to which it has an external likeness have not been studied by means of sections, so that this form stands rather isolated.

***Lithothamnium amphiroæformis*, Rothpletz.**


1891. *Lithothamnium amphiroæformis*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 43, p. 314, pl. xv, figs. 10, 14.

Very delicate cylindrical branches, dichotomously divided, and only 0.5 mm. in diameter. The cells are $12-15\ \mu$ wide and $20-30\ \mu$ long, and in the hypothallium up to $100\ \mu$ in length. No tetraspores or conceptacles have been recognised.

The fact that the cells are so much larger in this form than in *L. turonicum*, although it is externally of smaller size, precludes the likelihood of this species being a juvenile stage



Text-fig. 18.—*Lithothamnium turonicum*, Rothpletz. A. External form, natural size. B. Part of the thallus to show the cells and tetraspore zone, $\times 25$. After Rothpletz.



Text-fig. 19.—*Lithothamnium amphiroëformis*, Rothpletz. A. External form, natural size. B. Transverse section, and C. Longitudinal section, showing the zones of cells. B $\times 45$ and C $\times 50$. After Rothpletz.

of *L. turonicum*, as might at first be imagined from the external similarity.

HORIZON.—Turonian.


LOCALITY.—Beausset, Var, France.

TYPE.—Palaeontological Museum, Old Academy, Munich.

***Lithothamnium gosaviense*, Rothpletz.**

1891. *Lithothamnium gosaviense*, Rothpletz, Zeitschr. deutsch. geol. Ges., vol. 43, p. 315, pl. xvii, fig. 3.

An encrusting form, taking on the shape of the object it covers; the upper surface branching or warty. The cells are



Text-fig. 20.—*Lithothamnium gosaviense*, Rothpletz. Section showing several tetraspore zones in the tissue. $\times 30$. After Rothpletz.

9–12 μ wide and 9–12 μ high, but in the hypothallium up to 25 μ long. The tetraspores are 30–40 μ wide and 50–70 μ high. Tetraspores are developed once the perithallium reaches a thickness of 300 μ .

HORIZON.—Senonian.

LOCALITY.—Gosau and Martignes, fairly common.

TYPE.—Palaeontological Museum, Old Academy, Munich.

ALGÆ?

Indeterminate remains.

- V. 11065. Supposed fucoidal remains in a coarse sandstone from North Kop Khan, Armenia (see Oswald, 1906, p. 339). The fragments are carbonised and permeate the matrix, and are entirely unrecognisable. They resemble fragments of higher plants rather than Algæ.

Armenia.

Presented by Dr. F. Oswald
and H. F. B. Lynch, Esq., 1904.

DESCRIBED CRETACEOUS "ALGÆ" OF EXCEEDINGLY DOUBTFUL
NATURE.

The following *may* represent traces of Algæ:—

Caulerpites bryodes, Debey & Ettings.

Caulerpites Eseri, Unger.

Chondrites cretaceus, Presl.

Chondrites dichotomus, Feistmantel.

Chondrites divaricatus, Debey & Ettings.

Chondrites filiciformis, Lesquereux.

Chondrites neocomensis, Heer.

Chondrites polymorphus, Hosius & v. d. Marck.

Chondrites Riemsdyki, Miquel.

Chondrites serpentinus, Heer.

Conservites Woodwardi, Mantell.

Fucoides æqualis, Brongniart.

Fucoides strictus, Brongniart.

Halyserites gracilis, Debey & Ettings.

Sphærococcites Meyrati, Fischer-Ooster.

The following appear to be poor impressions of Dicotyledonous or Monocotyledonous plants:—

Delesserites Thierensi, Miquel, in Hosius & v. d. Marck, and in Debey & Ettings. (dicot. leaf?).

Fucoides Brardii, Brongniart (leafy twig of gymnosperm).

Fucoides tuberculatus, Brongniart.

Halyserites Reichii, Sternberg.

Laminarites polystigma, Debey & Ettings.

Lochmophycus caulerpoides, Debey & Ettings.

Neosporangium foliaceum, Debey & Ettings.

Neosporangium undulatum, Debey & Ettings.

The following may be of vegetable origin, but are indeterminate:—

- | | |
|---|---|
| <i>Aulacophycus pedatus</i> , Heer. | <i>Confervites æquensis</i> , Debey & Ettings. |
| <i>Caulerpa Lehmanni</i> , Heer. | <i>Confervites dubius</i> , Berry. |
| <i>Caulerpites tenuis</i> , Fischer-Ooster (probably a small conifer twig). | <i>Delessertites Hampeanus</i> , Stiehler. |
| <i>Chondrites Bosqueti</i> , Miquel. | <i>Fucoides Brongniarti</i> , Mantell. |
| <i>Chondrites elegans</i> , Debey & Ettings. | <i>Fucoides cauliformis</i> , Fritsch (pith-cast?). |
| <i>Chondrites flexuosus</i> , Newberry (rootlets?). | <i>Fucoides lygnbianus</i> , Brongniart. |
| <i>Chondrites jugiformis</i> , Debey & Ettings. | <i>Fucoides orbignianus</i> , Brongniart. |
| <i>Chondrites rigidus</i> , Debey & Ettings. | <i>Gelidium trajectomosanum</i> , Debey & Ettings. |
| <i>Chondrites subcurvatus</i> , Hosius & v. d. Marek (rootlets?). | <i>Gyrochorte porrecta</i> , de Stefani. |
| <i>Chondrites subintricatus</i> , Debey & Ettings. (rootlets?). | <i>Haliserites contortuplicatus</i> , v. d. Marek. |
| <i>Chondrites subverticillatus</i> , Presl in Sternberg. | <i>Keckia ambigua</i> , Eichwald (fern petioles?). |
| <i>Chondrites vagus</i> , Debey & Ettings. (rootlets). | <i>Phycodes sericeus</i> , Debey & Ettings. |
| | <i>Sphærococcites Laubei</i> , Engelhardt. |
| | <i>Sphærococcites pinnatifidus</i> , Unger. |
- The following appear to be indeterminate as Algæ, and are probably the tracks of animals, trickling water, etc.:—
- | | |
|--|--|
| <i>Caulerpites pyramidalis</i> , Sternberg. | <i>Fucoides strangulatus</i> , Fritsch. |
| <i>Caulerpites Diesingi</i> , Unger. | <i>Keckia annulata</i> , Glocker. |
| <i>Chondrites fusiformis</i> , Fischer-Oost. | <i>Keckia cylindrica</i> , Otto. |
| <i>Codites neocomiensis</i> , Saporta & Marion. | <i>Keckia nodulosa</i> , Otto. |
| <i>Confervites cæspitosus</i> , Debey & Ettings. | <i>Keckia vesiculosa</i> , Otto. |
| <i>Confervites ramosus</i> , Debey & Ettings. | <i>Münsteria Schneideriana</i> , Goeppert. |
| <i>Cylindrites arteriaeformis</i> , Goeppert. | <i>Neomeris annulus</i> , Böhm. |
| <i>Cylindrites conicus</i> , Hosius & v. d. Marek. | <i>Nulliporites granulosus</i> , Heer. |
| <i>Cylindrites dedaleus</i> , Goeppert. | <i>Tenidium alysioides</i> , Hosius & v. d. Marek. |
| <i>Cylindrites spongioides</i> , Goeppert. | <i>Taonurus Saportai</i> , Dewalque. |
| <i>Fucoides? columnaris</i> , Fritsch. | <i>Taonurus tenuistriatus</i> , Heer. |
| <i>Fucoides funiformis</i> , Fritsch. | <i>Zonarites allicornis</i> , Fischer-Ooster. |
| <i>Fucoides fritburgensis</i> , Heer. | <i>Zoophycos Brianteus</i> , Massalongo. |
| <i>Fucoides latifrons</i> , Heer. | <i>Zoophycos Villæ</i> , Massalongo. |
| | <i>Hexagonaria senonica</i> , Deecke (a sponge). |

These lists do not comprise all the supposed Algæ from Cretaceous rocks, but include the majority of them and indicate the extent of the described material. Some synonyms and further

"species," with references to the original descriptions, will be found in the list of Cretaceous plants at the beginning of the volume.

Class FUNGI.

A group of plants parallel to the Algæ, ranging from unicellular to large multicellular forms. They are all without chlorophyll, unable to manufacture their own food, and are therefore parasitic or saprophytic.

In comparison with the numerous fossils, vegetable and otherwise, which have been described as Algæ, there are very few Cretaceous specimens which have been placed in the group of Fungi.

A few "species" of parasitic fungi have been described as having attacked several of the dicotyledonous leaves which are so numerous in Cretaceous times; and while it is possible that a few of these are in reality the remains of fungal infection-spots on the leaves, most of them suggest that they owe their origin to the accidents of petrification. In 1896, Krasser (p. 116, pls. xi-xii) described and figured some traces of fungi on the supposed Monocotyledon *Typhaloipum cretaceum*, but he refrained from naming them, and said "aus der Kreideformation ist bisher überhaupt kein einziger Pilz bekannt über dessen systematische Stellung man sich mit Sicherheit äussern könnte."

There are now known, however, undoubted fungal petrifications. Most of these are found in the woods which have their anatomical structure preserved, and in which many of the cells are permeated by fungi. Among the petrified remains of the Japanese Cretaceous forests recently described by Stopes & Fujii (1910), parasitic fungi showing their reproductive organs have been discovered inhabiting the leaves and stems of the higher plants. These are among the best petrifications of fungi known from any geological formation. Septate and aseptate forms are recorded, and Suzuki (1910) even figures the microscopic section of a petrification showing the details of an ascomycetous perithecium.


Sub-Class **Ascomycetes.**

Order PYRENOMYCETES.

Genus **PLEOSPORITES**, Suzuki.

[Bot. Mag. Tokyo, vol. 24, 1910, p. 194.]

A monotypic genus with the following species.

Pleosporites Shirainus, Suzuki.1910. *Pleosporites Shirainus*, Suzuki, Bot. Mag. Tokyo, vol. 24, pp. 191-194, text-figs. 2, 3, pl. vii, phot. 6.Microscopic fungus in which the "hyphæ are well developed, septate, 2-5 μ in diameter. Perithecium formed under the

Text-fig. 21.—*Pleosporites Shirainus*, Suzuki. Perithecium cut longitudinally through the neck and orifice, showing asci (*as*); paraphyses (*pr*); hyphæ (*hy*); and epidermis of host (*ep*). $\times 370$. After Suzuki.

hypoderma of the leaves of the host, well defined, spherical or somewhat flask-shaped, with an orifice, short-necked, thick-walled, the wall consisting of 5-7 or more irregular layers of thin-walled plectenchyma cells. Asci formed at the bottom

of perithecium, without development of stroma; paraphyses present."

HOST.—Shoot of *Cryptomeriopsis*.

HORIZON.—Upper Cretaceous.

LOCALITY.—Hokkaido, Japan.

TYPE.—In Prof. Fujii's collection in the Botanical Department, Science College, Imperial University, Tokio, Japan.

This fungus is found excellently preserved, infecting most of the shoots and leaves of the Cretaceous Gymnosperm *Cryptomeriopsis*. The fructifications form under the hypoderm of the leaves, as is illustrated in the text-fig. 21.

The prevalence of this fungus in the remains of *Cryptomeriopsis* is commented on by Fujii (1910), who compares it with those infesting the living genus *Cryptomeria*, and considers that it may have been one of the factors in the extinction of the host.

- V. 11988. A microscopic section of three leaves of *Cryptomeriopsis antiqua*, Stopes & Fujii, all affected by a fungus which is in all probability the same as that described by Suzuki as *Pleosporites Shirainus* infecting *Cryptomeriopsis mesozoica*, Suzuki. One of the fungal perithecia is cut nearly medianly in this section, and two other perithecia in the other leaves are cut very tangentially. Throughout the mesophyll of all the leaves there are numerous ramifying hyphae which show their septations in many places. In their description of the host-plant, *Cryptomeriopsis antiqua*, Stopes & Fujii (1910, p. 55) noted the presence of this fungus but did not describe it further. It is of course possible that it is not of the same species as that more recently described by Suzuki (1910) on his species of *Cryptomeriopsis*; indeed, judging from the "biological species" of fungi which result from their adaptations to different hosts, it is more than likely that the two fungi would be classed as different species were they both alive. In the present imperfect state of our knowledge, however, it seems better to place this fungus in Suzuki's species, *Pleosporites Shirainus*, with a note to the effect that

it inhabits a different host from that attacked by his original species, than to multiply specific names.

Upper Cretaceous; Hokkaido, Japan.

Presented by Dr. M. C. Stopes, 1910.

Genus **PETROSPHÆRIA**, Stopes & Fujii.

[Phil. Trans. Roy. Soc. Lond., 1910 B, p. 6.]

Monotypic genus with the following species.

Petrosphæria japonica, Stopes & Fujii.

[Plate II.]

1909-10. *Petrosphæria japonica*, Stopes & Fujii, Proc. Roy. Soc. Lond., vol. 81, p. 599 (abstract), and Phil. Trans. Roy. Soc. Lond., 1910 B, pp. 4-6, pl. i, figs. 1-6.

Microscopic fungus in which the "hyphæ are septate, 2-4 μ in diameter, the septa at right angles to the long walls. Many cells of the hyphæ irregularly swollen to a large size. These swollen cells usually thickened, and forming round or irregular nests of plectenchyma which has often extremely thick walls. These clusters, entirely within the tissue of the host, are principally in the periderm."

True spore-formation at present unknown.

HOST.—*Saururus*, on the apparently underground stems.

HORIZON.—Upper Cretaceous.

LOCALITY.—Hokkaido, Japan.


TYPE.—British Museum (Natural History).

The stem of the Angiospermic genus *Saururus* was found to be thickly infested with septate hyphæ and also masses of reproductive or resting cells (see text-fig. 22).

No other fossil form is known with which this can be compared, but resting masses of thickened cells are not uncommon among living members of the Sphæriaceæ, with which it is probable the fossil has some affinity. Whether the fossil was a parasitic or saprophytic form is not determinable, but it appears to have lived underground, for the part of the host it was infesting was a rhizome or underground stem.

V. 11968. Holotype. Stopes & Fujii, 1910, Phil. Trans. Roy. Soc. Lond., pl. i, figs. 1, 2, 3, 5, & 6. There is little

to add to the original description of this specimen. The stem affected by the fungus is one of several lying in the mineral matrix of the slide, but is the only one of the genus *Saururopsis* hitherto discovered. Throughout the cortex numerous blackened masses are to be seen which are possibly the effete clusters of the fungal tissues, but may be due to some secretion



Text-fig. 22.—*Petrosphæria japonica*, Stopes & Fujii. A & B. Ends of hyphae cut off by transverse septum, and much swollen and thickened. C & D. Nests of thickened fungal "cells" in the periderm of the host showing the form in which they commonly occur. $\times 750$. After Stopes & Fujii.

natural to the host-stem. All round the stem just below and within the periderm the hyphae and reproductive masses have been petrified in the midst of their activity. Hyphae with swollen ends similar to those in the text-figure are seen in considerable variety. Micro. section in slide 1 B2 in Stopes Coll.

Upper Cretaceous; Hokkaido, Japan.

Presented by Dr. M. C. Stopes, 1910.

V. 11967. Paratype. Section through the same stem with fungus. Slide 1 B1 in Stopes Coll.

Upper Cretaceous; Hokkaido, Japan.

Presented by Dr. M. C. Stopes, 1910.

V. 11969. Paratype; section through the same stem showing the fungus very widely spread through the tissues. The distribution of the fungus in the periderm layers of the host is illustrated well in this section, a small portion of which is drawn in Plate II to show both the hyphæ and a few of the irregular thickened fungal cells. Slide 1 B4 in Stopes Coll.

Upper Cretaceous; Hokkaido, Japan.

Presented by Dr. M. C. Stopes, 1910.

V. 11970. Paratype; next section to V. 11969. This shows similar hyphal developments to those in V. 11969, and also has in the periderm layers a number of particularly well preserved nests of thickened cells (*cf.* text-fig. 22). Several of these are seen quite at the edge of the host-tissues, though the majority of them are more deeply seated. Slide 1 B5 in Stopes Coll.

Upper Cretaceous; Hokkaido, Japan.

Presented by Dr. M. C. Stopes, 1910.

V. 11971. Paratype. Slide 1 B10 in Stopes Coll.

Upper Cretaceous; Hokkaido, Japan.

Presented by Dr. M. C. Stopes, 1910.

Doubtful members of the Pyrenomycetes:—

Genus **SPHÆRITES**, Meschinelli.

[In Saccardo, *Sylloge Fungorum*, x, p. 753.]

Presumably with the characters of the living *Sphæria*, but incompletely known.

Sphærites cretaceus (Heer), Meschinelli.

1883. *Sphæria cretacea*, Heer, *Flora foss. Grönlands*, p. 1, pl. lx, figs. 2, 2 b.

1892. *Sphærites cretaceus*, Meschinelli, *Syll. Fung. foss.*, p. 753.

1902. *Sphaerites cretaceus*, Meschinelli, Fungor. foss. omnium Iconog., p. 21, pl. x, figs. 6, 6 a.

"Sph. peritheciis numerosis, orbiculatis, 1-1½ mm. latis, ostiolo rotundato pertusis."

Host.—Leaf of *Viburnum zizyphoides*.

HORIZON.—Upper Cretaceous (Patoot Beds).

LOCALITY.—Greenland.

Heer's description (1883 A, p. 1) is followed exactly by Meschinelli, who adds nothing new to the original short statements concerning this doubtful fossil. It is, however, as Heer's original figure shows, rather more likely to be a fungus than most of the circular markings on leaf-impressions which are described as such.

***Sphaerites problematicus* (Knowlton), Meschinelli.**

1892. *Sphaeria problematica*, Knowlton in Lesquereux, Flora Dakota Group, p. 23, pl. xxxi, figs. 2, 2 a.

1895. *Sphaerites problematicus*, Meschinelli, Fungi fossiles, p. 657.

1902. *Sphaerites problematicus*, Meschinelli, Fungor. foss. omnium Iconog., p. 24, pl. x, figs. 26, 26 a.

"Peritheciis sparsis, punctiformibus, rotundatis vel ovalibus, raro triangulatis, 0.5-1 mm. diam., ostiolo centrali duobus annulis prominentibus circumdato."

Host.—Leaf of *Sterculia Snowii*.

HORIZON.—Dakota Group.

LOCALITY.—Kansas, U.S.A.

While it is possible that this species may represent a true pyrenomycetous fungus, there is no real evidence for the view, and the markings on the leaf may be due to galls, and not fungal in their nature.

Order DISCOMYCETES.

Genus **PHACIDITES**, Meschinelli.

[In Saccardo, Sylloge Fungorum, x, p. 776.]


Presumably with the characters of the living genus *Phacidium*, but incompletely known.

Phacidites circumscriptus (Bayer), nom. nov.

1899. *Phacidium circumscriptum*, Bayer, Sitzungsber. k. böhm. Ges. Wiss. Prag, p. 7, text-fig. 3.

1901. *Phacidium circumscriptum*, Bayer in Fritsch & Bayer, Studien böhm. Kreidepfl., p. 67, text-fig. 3.

Fungus on leaves with apothecia scattered or in groups of two or three, mostly round or very slightly angled, 1.5–2 mm.



Text-fig. 23.—*Phacidites circumscriptus* (Bayer), nom. nov. A. Leaf of *Aralia daphnophyllum* with the fungus, natural size. B. Small portion about 3 times enlarged to show the character of the apothecia. After Bayer.

in diameter, with a definite border. In some cases grooves run from the border to a central depression.

HOST.—Leaf of *Aralia daphnophyllum*.

HORIZON.—Upper Cretaceous (Perucr Beds).

LOCALITY.—Vyserovic, Bohemia.

In Bayer's original description (Bayer, 1899) he illustrates this fungus with two text-figures which show its very characteristic appearance. He states that it is rare, but readily distinguished from any of the other Cretaceous fungi.

Supposed fungus:—

Genus **RHYTISMITES**, Meschinelli.

[In Saccardo, *Sylloge Fungorum*, x, p. 781.]

Presumably with the characters of the living genus *Rhytisma*, but incompletely known.

Rhytismites hederæ (Heer), Meschinelli.

1882. *Rhytisma hederæ*, Heer, *Flora foss. Grönlands*, p. 20, pl. xxiv, fig. 6 a.

1892. *Rhytismites hederæ*, Meschinelli, *Syll. Fung. foss.*, p. 781.

1902. *Rhytismites hederæ*, Meschinelli, *Fungor. foss. omnium* *Iconog.*, p. 57, pl. xvii, fig. 5.

HORIZON.—Upper Cretaceous (Atane Beds).

LOCALITY.—Greenland.

TYPE.—Heer's specimen in University Geological Museum, Copenhagen, on leaf of *Hedera primordialis*.

Very many further specimens of the Greenland leaves, not only of *Hedera* but of other genera, show circular markings similar to those described by Heer as his fungus *Rhytisma hederæ*. A number of excellent specimens are in Prof. Nathorst's collection in Stockholm, and a comparison of these with Heer's original, and with other specimens, suggests strongly that the markings are truly organic but are those of galls, and not fungi.

The same conclusion is suggested for *Phacidium myrtophylli* described by Engelhardt (1892 A, p. 80, pl. ii, figs. 10 a, b).

Further "genera" which are probably due to GALLS are:—

Puccinites cretaceus, Velenovsky, 1889, p. 26, pl. iii, fig. 14, which is also the same as *Uredinites cretaceus*, Velenovsky, 1889, p. 29, fig. 1, a-b.

Phacidites communis (Feistmantel), Meschinelli, 1902, p. 54. = *Phacidium commune*, Feistmantel, 1874, p. 266.

Sclerotites sp. (Knowlton), Meschinelli, 1902, p. 99. = *Sclerotium* sp., Knowlton, 1892, p. 23, pl. lix, fig. 4.

OTHER PHENOMENA DESCRIBED AS DISCOMYCETES :—

Genus **HYSTERITES**, Meschinelli.

[In Saccardo, Sylloge Fungorum, x, p. 774.]

Hysterites protogæus (Heer), Meschinelli.1882. *Hysterium protogæum*, Heer, Flora foss. Grönlands, p. 20, pl. xxiv, figs. 9, 9 a.1892. *Hysterites protogæus*, Meschinelli, Syll. Fung. foss., p. 774.1902. *Hysterites protogæus*, Meschinelli, Fungor. foss. omnium Iconog., p. 24, pl. xv, figs. 6, 6 a.

HORIZON.—Upper Cretaceous (Atane Beds).

LOCALITY.—Greenland.

TYPE.—Heer's specimen in University Geological Museum, Copenhagen.

The twig on which the so-called fungus was detected by Heer (1882) appears to be that of a dicotyledon, which though it is not described by Heer is very common in the Greenland beds. The one specimen with the "fungus" which Heer mentions, he supposes to be a gymnosperm; but even superficially it appears much more like a dicotyledonous twig. In Stockholm, in Prof. Nathorst's rich collections of Arctic plant-fossils there are a number of similar stems which all show the small markings Heer took to be a fungus, but which bear the strongest likeness to lenticels.

Hysterites protogæus, in short, seems to be nothing but normal lenticels on woody twigs.

Order HYPHOMYCETES.

Genus **CERCOSPORITES**, nom. nov.


Presumably with the characters of the living genus *Cercospora*, but incompletely known.

Cercosporites coriococcus (Bayer), nom. nov.

1899. *Cercospora coriococcum*, Bayer, Sitzungsber. k. böhm. Ges. Wiss. Prag. p. 4, text-figs. 1, 1 a, 2.

1901. *Cercospora corioccum*, Bayer, in Fritsch & Bayer, Studien böhm. Kreidepfl., p. 66, text-fig. 2.

The fungus forms roundish, sharply delineated patches, generally distinct from each other but sometimes merging.



Text-fig. 24.—*Cercosporites corioccus* (Bayer). *a.* Leaflet of *Devalquea coriacea* covered with the fungus, natural size. *b.* A small portion enlarged twice to show the surface character of the fungal patches. *c.* Portion of leaf of *Hedera primordialis* showing the same fungus. After Bayer.

These are 2-5 mm. in diameter, with a surface marked by series of concentric striations; in some cases the centre is slightly depressed. These patches lie both on the leaf-lamina and over the central nerve.

Host.—Leaves of *Devalquea coriacea*, *Aralia daphnophyllum*, *Hedera primordialis*, and many others.

HORIZON.—Upper Cretaceous (Perucer Beds).

LOCALITY.—Vyserovic and Kounic, Bohemia. Very plentiful.

These markings on the dicotyledonous leaves of the Bohemian Perucer Beds are undoubtedly very characteristic and easy to recognise. I have seen them in other collections of Bohemian plants, and think it very probable that they do truly represent fungal remains, though the nature of the impressions does not allow of microscopic examination or of exact determination.

Order BASIDIOMYCETES?

Genus **TRAMETITES**, Meschinelli.

[In Saccardo, Sylloge Fungorum, x, p. 747.]

Presumably with the characters of the living genus *Trametes* but incompletely known.

Trametes Pini (Conwentz), Meschinelli.

1892. *Trametes Pini*, Conwentz, Unters. foss. Hölz. Schwed., pp. 13, 21, pl. vii, fig. 1.

1892. *Trametes Pini*, Meschinelli, Syll. Fung. foss., p. 747.

1902. *Trametes Pini*, Meschinelli, Fungor. foss. omnium Iconog., p. 6, pl. iv, fig. 5.

Meschinelli defines the species as follows:—Cl. doct. Conwentz innumeras mycelii reliquias, lignum illud dum viveret a parassitis vexatum ostendentes, in speciminibus succicis invenit. Hyphæ tenues, jalinae, cinerae, brunneae et quandoque atro-cinerae ad nigritudinem, ligni in directionem verticalem discurrunt."

In describing some Gymnospermic petrified woods from Sweden, Conwentz (1892) mentioned the frequent occurrence in them of fungal hyphæ. He gives a figure and short description of them and associates the name *Trametes Pini* with the fungus, though he does not absolutely identify it as such.

Host.—Wood of *Pinus Nathorsti*.

HORIZON.—Senonian.

LOCALITY.—Sweden.

TYPE.—Conwentz' slide in Palæobotanical Museum, Stockholm.

Meschinelli's figure is a reproduction of that of Conwentz, which shows the fungus in a low degree of magnification. The species may be taken to include the numerous hyphæ that are noticed in petrified woods of Cretaceous pines, though it is probably not a true biological species.

Genus **TRICHOSPORITES**, Felix.

[Zeitschr. deutsch. geol. Ges., vol. 46, 1894, p. 273.]

Presumably with the characters of the living genus *Trichosporium*, but incompletely known.

Trichosporites Conwentzi, Felix.

1892. "Cf. *Trichosporium fuscum*," Conwentz, Unters. foss. Hölz. Schwedens, p. 27, pl. vii, fig. 9.

1894. *Trichosporites Conwentzi*, Felix, Zeitschr. deutsch. geol. Ges., vol. 46, p. 273.

Branched, septate hyphæ, conidia egg-shaped or oval, circular when seen from one end. Composed of one cell, apparently undivided, and of dark, red-brown colour.

HOST.—Wood of *Cedroxylon Ryedalense*.

HORIZON.—Senonian.


LOCALITY.—Ryedal, Sweden.

TYPE.—Palæobotanical Museum, Stockholm.

The specific name and some description of this fungus appear in Felix's paper on Fossil Fungi (Felix, 1894), but it was figured by Conwentz in his paper on the petrified woods of Sweden (Conwentz, 1892), who described the effects of the fungus on the wood, *Cedroxylon Ryedalense*, and noted that the hyphæ are thick-walled, branched, and septate.

Conwentz' illustrations of the hyphæ and conidia are reproduced in the text-figure 25 (p. 280).

While the name appears to be unfortunate in being too suggestive of affinities with a living genus where such affinity cannot be established from the data at present available, it seems better to leave it than to multiply poorly established specific names.



Text-fig. 25.—*Trichosporites Conwentzi*, Felix ; hyphae and conidia.
After Conwentz.

PHENOMENA DESCRIBED AS FUNGI, OF WHICH THE
NATURE IS VERY DOUBTFUL:—

Xylomites ellipticus, Ettingshausen, 1867 A, p. 243, pl. i, fig. 7.
“Peritheciis ellipticis planis, disco centrali vix distinguendo.”

This “fungus” takes the form of small oval markings on the impression of the leaf of *Ficus Geinitzii*. Ettingshausen considered that there could scarcely be any doubt that the appearance is due to a fungus.

Phacidium Palaeocassiae, Ettingshausen, 1867 A, p. 242, pl. i, figs. 8, 8 b. "Peritheciis irregularibus polygonis depressis, disco subrotundato, pallido."

This is also a mere marking on a leaf-impression which may or may not be due to a fungus, but about which there is no clear evidence in the specimens themselves.

Sphaerites myricae (Lesquereux), Meschinelli, 1892, and 1902, p. 29 (= *Sphaeria myricae*, Lesquereux); and also *Sphaerites Lesquereuxi*, Meschinelli, 1892, & 1902, p. 38 (= *Sphaeria rhytismoides*, Lesquereux), are two other markings on leaves for which there is little evidence to support their claims to being fungi.

Stichus mermisoides, Etheridge, 1904, pp. 255-257, pls. xxx-xxxi. Described as an endophytic fungus in the shells of a Cretaceous bivalve. The plates, however, are far from convincing.

SO-CALLED "FUNGI" WHICH ARE PROBABLY INORGANIC
PSEUDOMORPHS :—

Sclerotites sp., Geinitz, 1842, pp. 99 & xxii, pl. xxiv, figs. 1-3.

Xylomites aggregatus, Heer, 1882, p. 21, pl. xxix, fig. 11.

Rosellinites lepidus (Lesquereux), Knowlton, 1898, p. 204.
= *Sphaeria lapidea*, Lesquereux, 1873, p. 373, & 1878 B, p. 34, pl. i, fig. 3.

INDEX

TO

DESCRIPTIVE CATALOGUE.

- Acetabularia, 247.
 Algæ, 233.
 — furcatus, 253.
 Amphitetras antediluviana, 235.
 Animal tracks etc. described as
 algæ, 266.
 Aralia daphnophyllum, 274.
 Ascomycetes, 268.
 Aulacophycus pedatus, 266.

 Basidiomycetes, 278.
 Boueina, 236.
 — Hochstetteri, 237.

 Caulerpa Lehmanni, 266.
 Caulerpites bryodes, 265.
 — Diesingi, 266.
 — Eseri, 265.
 — pyramidalis, 266.
 — tenuis, 266.
 Cedroxylon Ryedalense, 279.
 Cellepora bipunctata, 258.
 Cercospora coriococeum, 276.
 Cercosporites, 276.
 — coriococeus, 276.
 Ceriopera dichotoma, 259.
 Chondrites, 247.
 — equalis, 252.
 — affinis, 255.
 — Bosqueti, 266.
 — cretaceus, 265.
 — dichotomus, 265.
 — divaricatus, 255.
 — elegans, 266.
 — filiciformis, 265.
 — flexuosus, 266.
 — furcatus, 255.
 — fusiformis, 266.
 — genuinus, 255.
 — intricatus, 252.

 Chondrites jugiformis, 266.
 — neocomensis, 265.
 — patulus, 253.
 — polymorphus, 265.
 — Riemsdyki, 265.
 — rigidus, 266.
 — serpentinus, 265.
 — subcurvatus, 266.
 — subintricatus, 266.
 — subverticillatus, 266.
 — Targionii, 249.
 — vagus, 266.
 — Vindobonensis, 250, 252.
 Codiaceæ, 236.
 Codites neocomiensis, 266.
 Confervites æquensis, 266.
 — cespitosus, 266.
 — dubius, 266.
 — ramosus, 266.
 — Woodwardi, 265.
 Corallinaceæ, 256.
 Coscinodiscus Argus, 235.
 — eccentricus, 235.
 — Patina, 235.
 Cryptomeriopsis antiqua, 269.
 — mesozoica, 269.
 Cylindrites arteriæformis, 266.
 — conicus, 266.
 — dædaleus, 266.
 — spongioides, 266.

 Dasycladaceæ, 239.
 Delessertites Thierensi, 265.
 Delessertites Hampeanus, 266.
 Dewalquea coriacea, 277.
 Diatomaceæ, 235.
 Diplopore, 243.
 — Mühlbergii, 243.
 Discomycetes, 273.
 Discophorites, 247.


- Discophorites angustilobus, 247.
 — Fischeri, 247.
 Discopora bipunctata, 258.
 Ficus Geinitzii, 280.
 Fragilaria rhabdosoma, 235.
 — striolata, 236.
 Fucoides, 248.
 — aequalis, 265.
 — Brardii, 265.
 — Brongniarti, 266.
 — cauliformis, 266.
 — columnaris, 266.
 — friburgensis, 266.
 — funiformis, 266.
 — furcatus, 255.
 — intricatus, 252.
 — latifrons, 266.
 — lygnbianus, 266.
 — orbignianus, 266.
 — strangulatus, 266.
 — strictus, 265.
 — Targionii, 249.
 — tuberculosus, 265.
 Fungus, 267.
 Gallionella aurichalcea, 236.
 — sulcata, 236.
 Galls, 275.
 Gelidium trajectomosanum, 266.
 Gigartinites, 248.
 Goniolina, 246.
 Gyrochorte porrecta, 266.
 Gyrophyllites, 247.
 — obtusifolius, 247.
 — Oosteri, 247.
 — pentamerus, 247.
 Gyroporella, 242, 247.
 Halimeda, 237, 239.
 — Fuggeri, 239.
 Halimedes, 239.
 Haliserites contortuplicatus, 266.
 — Reichii, 265.
 Halyserites gracilis, 265.
 Hedera primordialis, 275, 277.
 Hexagonaria senonica, 266.
 Hyphomycetes, 276.
 Hysterites, 276.
 — protogæus, 276.
 Hysterium protogæum, 276.
 Keckia ambigua, 266.
 — annulata, 266.
 — cylindrica, 266.
 — nodulosa, 266.
 — vesiculosa, 266.
 Laminarites polystigma, 265.
 Lenticels, described as fungi, 276.
 Lithothamnium, 256.
 — amphiroæformis, 262.
 — cenomanicum, 260.
 — Goldfussi, 260.
 — gosaviense, 264.
 — mamillosum, 257.
 — nummuliticum, 262.
 — palmatum, 259.
 — parisiense, 259.
 — perulatum, 258.
 — procoenum, 259.
 — racemosum, 260.
 — racemus, 262.
 — ramosissimum, 257.
 — turonicum, 262.
 Lochmophycus caulerpoides, 265.
 Marginaria bipunctata, 258.
 Membranipora bipunctata, 258.
 Munieria, 240.
 — baconica, 240.
 Münsteria Schneideriana, 266.
 Navicula ventricosa, 236.
 Nematophycus, 234.
 Neomeris, 239.
 — annulus, 266.
 — cretacea, 240.
 Neosporangium foliaceum, 265.
 — undulatum, 265.
 Nullipora palmata, 259.
 Nulliporites granulosus, 266.
 Petrosphæria, 270.
 — japonica, 270.
 Phacidites, 273.
 — circumscriptus, 274.
 — communis, 275.
 Phacidium, 273.
 — circumscriptum, 274.
 — commune, 275.
 — myrtophylli, 275.
 — Palæocassia, 281.
 Phæophyceæ, 247.
 Phycodes sericeus, 266.
 Phycopsis, 249.
 — affinis, 255.
 — intricata, 252.
 — Targioni, 250.
 Pinus Nathorsti, 278.
 Pleosporites, 268.
 — Shiraius, 268.
 Puccinities cretaceus, 275.
 Pyrenomycetes, 268.

- Rhodophyceæ, 256.
 Rhytisma, 275.
 — hederæ, 275.
 Rhytismites, 275.
 — hederæ, 275.
 Rosellinites lepideus, 281.
 Saururopsis, 270.
 Sclerotites, 275, 281.
 Sclerotium, 275.
 Siphonaceæ, 236.
 Sphæria, 272.
 — cretacea, 272.
 — lapidea, 281.
 — myricæ, 281.
 — problematica, 273.
 — rhytismoides, 281.
 Sphærites, 272.
 — cretaceus, 272.
 — Lesquereuxi, 281.
 — myricæ, 281.
 — problematicus, 273.
 Sphærococcites Laubei, 266.
 — Meyrati, 265.
 — pinnatifidus, 266.
 Sphærococcus, 255.
 Sterculia Snowii, 273.
 Stichus mermisoides, 281.
 Tænidium alysioides, 266.
 Taonurus Saportai, 266.
 — tenuistriatus, 266.
 Thallophyta, 233.
 Trametes, 278.
 — Pini, 278.
 Trametites, 278.
 — Pini, 278.
 Trichosporites, 279.
 — Conwentzi, 279.
 Trichosporium fuscum, 279.
 Triploporella, 244.
 — Fraasi, 245.
 Typhæloipum cretaceum, 267.
 Uredinites cretaceus, 275.
 Viburnum zizyphoides, 273.
 Xylomites aggregatus, 281.
 — ellipticus, 280.
 Zonarites alcicornis, 266.
 Zoophycos Brianteus, 266.
 — Villæ, 266.


PLATE I.

Fig. 1. A slab of rock from the Cretaceous Flysch, with *a. Alcyites
furcatus* (p. 255), and *b. Chondrites intricatus* (p. 252).
Nat. size.

Fig. 2. *Chondrites patulus*. Nat. size. (p. 253)



2.



1.

1a. ALGITES FURCATUS. 1b. CHONDRITES INTRICATUS.
2. CHONDRITES PATULUS.

J. Green Photo imp.


PLATE II.

Part of the outer cortex and periderm of the Angiospermic stem *Saururopsis* which is infested by the fungus *Petro-sphaeria japonica*, Stopes & Fujii. V. 11969.

- M. The outer limit of the stem, slightly crushed and destroyed adjacent to the granular mineral matrix of the section.
- K. Cells of periderm layers in which hyphæ and other cells of the fungus are principally developed.
- C. Inner cells of cortex in which are hyphæ of fungus, but no other fungal cells which appear to have been vital at the time of petrification.
- H. Fungal hyphæ, at H' a part showing the septate condition.
- T. Groups of thickened cells of the fungus.
- E. Cells in the cortex containing blackened "frothy" substance which may be due to the fungal cells, or may be due to some natural secretion of the plant.

Magnification $\times 300$.

(p. 270)



M. C. Stopes del.

J. Green lith.

LIST OF THE CURRENT
NATURAL HISTORY PUBLICATIONS
OF THE TRUSTEES OF THE
BRITISH MUSEUM.

The following publications can be purchased through the Agency of *Messrs. LONGMANS, GREEN & Co., 39, Paternoster Row, London, E.C.; Mr. QUARITCH, 11, Grafton Street, New Bond Street, W.; and Messrs. DULAU & Co., Ltd., 37, Soho Square, W.; or at the NATURAL HISTORY MUSEUM, Cromwell Road, London, S.W.*

The History of the Collections contained in the Natural History Departments of the British Museum :—

Vol. I. Libraries; Botany; Geology; Minerals. Pp. xvii., 442. 1904, 8vo. 15s.

Vol. II. Separate historical accounts of the several collections included in the Department of Zoology. Pp. 782. 1906, 8vo. 17. 10s.

Vol. II.—Appendix. General History of the Department of Zoology from 1856 to 1895. By Dr. Albert Günther, F.R.S., formerly Keeper of the Department of Zoology. Pp. ix., 109. [With Index.] 1912, 8vo. 5s.

Catalogue of the Books, Manuscripts, Maps, and Drawings in the British Museum (Natural History). Compiled by B. B. Woodward, Assistant in charge of the General Library, with some clerical assistance :—

Vol. I. A—D. Pp. viii., 500. 1903, 4to. 17.

Vol. II. E—K. Pp. 501–1038. 1904, 4to. 17.

Vol. III. L—O. Pp. 1039–1494. 1910, 4to. 17.

Vol. IV. P—SN. Pp. 1495–1956. 1913, 4to. 17.

Catalogue of the Specimens and Drawings of Mammals, Birds, Reptiles, and Fishes of Nepal and Tibet. Presented by B. H. Hodgson, Esq., to the British Museum. 2nd edition. By John Edward Gray. Pp. xii., 90. [With an account of the Collection by Mr. Hodgson.] 1863, 12mo. 2s. 3d.

Report on the Zoological Collections made in the Indo-Pacific Ocean during the voyage of H.M.S. "Alert," 1881-2. Pp. xxv., 684. 54 Plates. 1884, 8vo. 1*l.* 10s.

Summary of the Voyage...	By Dr. R. W. Coppinger.
Mammalia	" O. Thomas.
Aves...	" R. B. Sharpe.
Reptilia, Batrachia, Pisces	" A. Günther.
Mollusca	" E. A. Smith.
Echinodermata	" F. J. Bell.
Crustacea	" E. J. Miers.
Coleoptera	" C. O. Waterhouse.
Lepidoptera	" A. G. Butler.
Alcyonaria and Spongiida	" S. O. Ridley.

Report on the Collections of Natural History made in the Antarctic Regions during the Voyage of the "Southern Cross." Pp. ix., 344. 53 Plates. 1902, Royal 8vo. 2*l.*

Mammalia	By Capt. G. E. H. Barrett-Hamilton.
Notes on Antarctic Seals ...	"	E. A. Wilson, M.B.
Extracts from the "Diary" of the late Nicolai Hanson.	}	
Aves	" Dr. R. Bowdler Sharpe.
Pisces...	...	" G. A. Boulenger, F.R.S.
Tunicata	" Prof. W. A. Herdman, F.R.S.
Mollusca	" E. A. Smith.
Echinoderma	" Prof. F. J. Bell.
Insecta	" G. H. Carpenter and Hon. N. C. Rothschild.
Arachnida	" Dr. E. L. Trouessart.
Crustacea	" T. V. Hodgson.
Polychaeta	" Dr. A. Willey.
Gephyrea	" A. E. Shipley.
Nematoda	" Dr. von Linstow.
Cestoda	
Polyzoa	" R. Kirkpatrick.
Porifera	
Anthozoa	" Dr. L. Roule and S. J. Hickson, F.R.S.
Actiniae	" J. A. Clubb.
Hydrozoa	" E. T. Browne.
Cryptogamia	" A. Gepp, V. H. Blackman, and Miss E. S. Barton.
Rock Specimens	" Dr. G. T. Prior.

National Antarctic Expedition, 1901-1904. [*Discovery Report.*] Natural History:—

Vol. I. Geology (Field-Geology: Petrography). Pp. xii., 160: 10 plates, 72 text-figures, 2 maps. [With Index.] 1907, 4to. 1*l.* 10s.

Field-Geology	By H. T. Ferrar, M.A., F.G.S.
Rock-Specimens	" G. T. Prior, M.A., D.Sc., F.G.S.

Vol. II. Zoology (Vertebrata: Mollusca: Crustacea). Pp. xiv., 355: 33 plates (17 coloured), 146 text-figures, 1 map. 1907, 4to. 3*l.*

Mammalia (pp. 69: 5 pls. [3 col.], 36 figs.)	}	By E. A. Wilson, M.B.
Aves (pp. 121: 13 pls. col., 46 figs.)		

National Antarctic Expedition, 1901-1904—*continued*.Vol. II—*continued*.

- Anatomy of Penguins (pp. 28: 1 pl., 8 figs.) By W. P. Pyecraft.
- Fishes (pp. 5; 2 pls.) " G. A. Boulenger, F.R.S.
- Cephalodiscus (pp. 67: 7 pls. [1 col.], 17 figs.) " W. G. Ridewood, D.Sc.
- Mollusca. I.—Cephalopoda (pp. 2: 1 fig.) " W. E. Hoyle, D.Sc.
- " II.—Gastropoda { (pp. 12, 1: 1 fig.)
- " III.—Amphineura { 2 pls.) " E. A. Smith, I.S.O.
- " IV.—Nudibranchiata (pp. 28: 1 pl., 30 figs.) " Sir Charles Eliot, K.C.M.G., LL.D.
- " V.—Lamellibranchiata (pp. 7: 1 pl.) " E. A. Smith, I.S.O.
- Brachiopoda (pp. 2: 4 figs.)
- Crustacea I.—Decapoda (pp. 7)
- " II.—Cumacea (pp. 6: 1 pl., 4 figs.) " W. T. Calman, D.Sc.

Vol. III. Zoology and Botany (Invertebrata: Marine Algae, Musci). Pp. vi., 273: 51 plates, 8 text-figures, and 1 chart. 1907, 4to. 2l. 10s.

- On Collecting in Antarctic Seas (pp. 10) By T. V. Hodgson, F.L.S.
- Mollusca. VI.—Pteropoda (pp. 15: 2 pls.) " Sir Charles Eliot, K.C.M.G., LL.D.
- Crustacea. III.—Amphipoda (pp. 39: 13 pls.) " A. O. Walker, F.L.S.
- " IV.—Leptostraca (pp. 2: 2 figs.) " Dr. J. Thiele.
- " V.—Ostracoda (pp. 9: 3 pls.) " Prof. G. S. Brady, F.R.S.
- " VI.—Cirripedia (pp. 4: 1 pl.) " Prof. A. Gruvel.
- Pycnogonida (pp. 72: 10 pls.) " T. V. Hodgson, F.L.S.
- Acari (pp. 6: 1 pl.) " Dr. E. L. Trouessart.
- Chaetognatha (pp. 6: 1 chart) " Dr. G. H. Fowler, F.L.S.
- Nematoda (pp. 4: 1 pl.) " Dr. O. von Linstow.
- Cestoda (pp. 6: 1 pl.) " A. E. Shipley, F.R.S.
- Coelentera. I.—Alcyonaria (pp. 15: 2 pls.) " Prof. S. J. Hickson, F.R.S.
- " II.—Hydroid Zoophytes (pp. 34: 4 pls.) " Prof. S. J. Hickson, F.R.S., and F. H. Gravely.
- " III.—Tentacles of a Siphonophore (pp. 3: 5 figs.) " Dr. J. Rennie.
- Porifera. I.—Hexactinellida (pp. 25: 7 pls.) " R. Kirkpatrick.
- Marine Algae. I.—Phaeophyceae and Florideae (pp. 15: 4 pls.) " A. Gepp and Mrs. E. S. Gepp.
- " II.—Corallinaceae (pp. 2: 1 fig.) " M. Foslie.
- Musci (pp. 6: 2 pls.) " J. Cardot.

Vol. IV. Zoology (various Invertebrata). Pp. iv., 281: 65 plates, 1 text-figure. 1908, 4to. 1l. 15s.

- Mollusca. VII.—Solenogastres (pp. 13: 2 pls.) By Dr. H. F. Nierstrasz.

National Antarctic Expedition, 1901-1904—*continued.*Vol. IV—*continued.*

- Insecta.—Aptera (pp. 5 : 1 pl.) ... By G. H. Carpenter, B.Sc.,
M.R.I.A.
Crustacea. VII.—Schizopoda (pp. 42 :
8 pls.) .. W. M. Tattersall, M.Sc.
" VIII.—Copepoda (pp. 44 :
7 pls.) .. R. Norris Wolfenden,
M.D.
Echinoderma (pp. 16 : 5 pls.) ... By F. Jeffrey Bell, M.A.
Echinoderm Larvæ (pp. 9 : 1 pl.) ... " E. W. McBride, M.A.,
F.R.S., and J. C.
Simpson, B.Sc.
Myzostomidae (pp. 26 : 1 pl. and 1 figure) .. Dr. Rudolf Ritter von
Stummer-Traunfels.
Sipunculoidea (pp. 6) " W. F. Lanchester, M.A.
Coelentera. IV.—Actiniae (pp. 12 : 3 pls.) .. J. A. Clubb, M.Sc.
Porifera. II.—Tetraxonida (pp. 56 :
19 pls.) .. R. Kirkpatrick.
" III.—Calcarea (pp. 52 : 12 pls.) .. C. F. Jenkin, B.A.

Vol. V. Zoology and Botany. Pp. ix., 212 : 28 plates, and
19 text-figures. [With a Summary of the Contents of
Vols. II.-V.] 1910, 4to. 17. 10s.

- Vertebrata. IA.—Mammalia (Seal-Em- By Dr. H. W. Marett
bryos) (pp. 21 : 2 pls., 1 text-figure) Tims.
Tunicata (pp. 26 : 7 pls., 2 text-figures) .. Prof. W. A. Herdman,
D.Sc., F.R.S.
Crustacea. IX.—Isopoda (pp. 77 : 10 pls) .. T. V. Hodgson, F.L.S.
Nemertinea (pp. 15 : 1 pl., 16 text- Prof. L. Joubin.
figures)
Coelentera. V.—Medusæ (pp. 62 : 7 pls.) .. Edward T. Browne.
Lichenes (pp. 11 : 1 pl.)... .. Dr. O. V. Darbishire.

Vol. VI. Zoology and Botany. Pp. xvi., 101 : 8 plates,
1 key-plate, and 1 text-figure. [With Synopsis of
Contents of the several volumes of Memoirs, Classified
Summary of the Contents of Vols. I.-VI., Subject
Index, and Index of Authors.] 1912, 4to. 16s.

- Echinoderma. III.—On a collection of By Prof. E. W. MacBride,
young Holothurioids (pp. 9 : 2 pls., D.Sc., LL.D., F.R.S.
and key to pl. 1.)
Polychæta (pp. 32 : 3 pls.) " Prof. E. Ehlers.
Freshwater Algæ (pp. 60 : 3 pls., 1 text-figure) .. Dr. F. E. Fritsch, F.L.S.

A Monograph of Christmas Island (Indian Ocean) : Physical
Features and Geology by C. W. Andrews, B.A., B.Sc.,
F.G.S., with descriptions of the Fauna and Flora by
numerous contributors. Pp. xv., 337 : 22 plates (7 coloured),
a map, and 27 illustrations in text. [With Index.] 1900,
8vo. 17.First Report on Economic Zoology. By Fred. V. Theobald
M.A., &c. [With an Introduction, containing a Classifica-
tion of Animals from the point of view of Economic
Zoology, by Prof. E. Ray Lankester, LL.D., F.R.S.]
Pp. xxxiv., 192. 18 Woodcuts. 1903, Roy. 8vo. 6s.

Second Report on Economic Zoology. By Fred. V. Theobald, M.A., &c. Pp. x., 197. 29 Illustrations. 1904, Roy. 8vo. 6s.

MAMMALS.

Catalogue of the Bones of Mammalia in the Collection of the British Museum. By Edward Gerrard. Pp. iv., 296, 1862, 8vo. 5s.

Catalogue of Monkeys, Lemurs, and Fruit-eating Bats in the Collection of the British Museum. By Dr. J. E. Gray, F.R.S., &c. Pp. viii., 137. 21 Woodcuts. 1870, 8vo. 4s.

Catalogue of the Chiroptera in the Collection of the British Museum. Second Edition. By Knud Andersen. Vol. I. Megachiroptera. Pp. ci., 854: 85 text-figures. [With Table of Contents, Introduction, and Index.] 1912, 8vo. 2l. 10s.

Catalogue of Carnivorous Pachydermatous, and Edentate Mammalia in the British Museum. By John Edward Gray, F.R.S., &c. Pp. vii., 398. 47 Woodcuts. 1869, 8vo. 6s. 6d.

Catalogue of Seals and Whales in the British Museum. By John Edward Gray, F.R.S., &c. 2nd Edition. Pp. vii., 402. 101 Woodcuts. 1866, 8vo. 8s.

—Supplement. By John Edward Gray, F.R.S., &c. Pp. vi., 103. 11 Woodcuts. 1871, 8vo. 2s. 6d.

Catalogue of Ruminant Mammalia (*Pecora*, Linnæus) in the British Museum. By John Edward Gray, F.R.S., &c. Pp. viii., 102. 4 Plates. 1872, 8vo. 3s. 6d.

Catalogue of the Ungulate Mammals in the British Museum (Natural History). Vol. I. Artiodactyla, Family Bovidae, Subfamilies Bovinae to Ovibovinae (Cattle, Sheep, Goats, Chamois, Serows, Takin, Musk-Oxen, etc.). By R. Lydekker, F.R.S. Pp. xvii., 249: 55 text-figures. [With List of Illustrations and Index of Genera and Species.] 1913, 8vo. 7s. 6d.

Monograph of the Okapi. By Sir E. Ray Lankester, K.C.B., M.A., D.Sc., F.R.S., &c. Atlas. Compiled with the assistance of W. G. Ridewood, D.Sc. 48 Plates, with xxii. pp. of Explanations, &c. 1910, 4to. 1l. 5s.

Catalogue of the Marsupialia and Monotremata in the Collection of the British Museum. By Oldfield Thomas. Pp. xiii., 401. 4 Coloured and 24 plain Plates. [With Systematic and Alphabetical Indexes.] 1888, 8vo. 1l. 8s.

Catalogue of the Mammals of Western Europe (Europe exclusive of Russia) in the Collection of the British Museum. By Gerrit S. Miller. Pp. xv., 1019. 213 Text-figures. [With Systematic and Alphabetical Indexes.] 1912, 8vo. 1l. 6s.

Catalogue of the Heads and Horns of Indian Big Game bequeathed by A. O. Hume, C.B., to the British Museum (Natural History). By R. Lydekker, F.R.S. Pp. xvi., 45. 16 Text-figures, and a Portrait of Mr. Hume. [With an Alphabetical Index.] 1913, 8vo. 2s.

BIRDS.

Catalogue of the Birds in the British Museum :—

Vol. VII. Catalogue of the Passeriformes, or Perching Birds. *Cichlomorphæ* : Part IV., containing the concluding portion of the family Timeliidæ (Babbling Thrushes). By R. Bowdler Sharpe. Pp. xvi., 698. Woodcuts and 15 coloured Plates. [With Systematic and Alphabetical Indexes.] 1883, 8vo. 17. 6s.

Vol. VIII. Catalogue of the Passeriformes, or Perching Birds. *Cichlomorphæ* : Part V., containing the families Paridæ and Laniidæ (Titmice and Shrikes) ; and *Certhiomorphæ* (Creepers and Nuthatches). By Hans Gadow, M.A., Ph.D. Pp. xiii., 386. Woodcuts and 9 coloured Plates. [With Systematic and Alphabetical Indexes.] 1883, 8vo. 17s.

Vol. X. Catalogue of the Passeriformes, or Perching Birds. *Fringilliformes* : Part I., containing the families Dicaeidae, Hirundinidae, Ampelidae, Mniotiltidae, and Motacillidae. By R. Bowdler Sharpe. Pp. xiii., 682. Woodcuts and 12 coloured Plates. [With Systematic and Alphabetical Indexes.] 1885, 8vo. 17. 2s.

Vol. XI. Catalogue of the Passeriformes, or Perching Birds. *Fringilliformes* : Part II., containing the families Cœrebidæ, Tanagridæ, and Icteridæ. By Philip Lutley Sclater, M.A., F.R.S. Pp. xvii., 431. Woodcuts and 18 coloured Plates. [With Systematic and Alphabetical Indexes.] 1886, 8vo. 17.

Vol. XII. Catalogue of the Passeriformes, or Perching Birds. *Fringilliformes* : Part III., containing the family Fringillidae. By R. Bowdler Sharpe. Pp. xv., 871. Woodcuts and 16 coloured Plates. [With Systematic and Alphabetical Indexes.] 1888, 8vo. 17. 8s.

Vol. XIII. Catalogue of the Passeriformes, or Perching Birds. *Sturniformes*, containing the families Artamidæ, Sturnidæ, Ploceidæ, and Alaudidæ. Also the

Catalogue of the Birds in the British Museum—*continued*.

families Atrichiidae and Menuridae. By R. Bowdler Sharpe. Pp. xvi., 701. Woodcuts and 15 coloured Plates. [With Systematic and Alphabetical Indexes.] 1890, 8vo. 1*l.* 8*s.*

Vol. XIV. Catalogue of the Passeriformes, or Perching Birds. *Oligomyzodæ*, or the families Tyrannidae, Oxyrhamphidae, Pipridæ, Cotingidae, Phytotomidae, Philepittidae, Pittidae, Xenicidae, and Eurylamidae. By Philip Lutley Sclater, M.A., F.R.S. Pp. xix., 494. Woodcuts and 26 coloured Plates. [With Systematic and Alphabetical Indexes.] 1888, 8vo. 1*l.* 4*s.*

Vol. XV. Catalogue of the Passeriformes, or Perching Birds. *Tracheophonæ*, or the families Dendrocolaptidae, Formicariidae, Conopophagidae, and Pteroptochidae. By Philip Lutley Sclater, M.A., F.R.S. Pp. xvii., 371. Woodcuts and 20 coloured Plates. [With Systematic and Alphabetical Indexes.] 1890, 8vo. 1*l.*

Vol. XVI. Catalogue of the Picariæ. *Upupæ* and *Trochili*, by Osbert Salvin. *Coraciæ*, of the families Cypselidae, Caprimulgidae, Podargidae, and Steatorniithidae, by Ernst Hartert. Pp. xvi., 703. Woodcuts and 14 coloured Plates. [With Systematic and Alphabetical Indexes.] 1892, 8vo. 1*l.* 16*s.*

Vol. XVII. Catalogue of the Picariæ. *Coraciæ* (contin.) and *Halcyones*, with the families Leptosomatidae, Coraciidae, Meropidae, Alcedinidae, Momotidae, Totidae and Coliidae, by R. Bowdler Sharpe. *Bucerotes* and *Trogonæ*, by W. R. Ogilvie Grant. Pp. xi., 522. Woodcuts and 17 coloured Plates. [With Systematic and Alphabetical Indexes.] 1892, 8vo. 1*l.* 10*s.*

Vol. XVIII. Catalogue of the Picariæ. *Scansores*, containing the family Picidae. By Edward Hargitt. Pp. xv., 597. Woodcuts and 15 coloured Plates. [With Systematic and Alphabetical Indexes.] 1890, 8vo. 1*l.* 6*s.*

Vol. XIX. Catalogue of the Picariæ. *Scansores* and *Coccyges*: containing the families Rhamphastidae, Galbulidae, and Bucconidae, by P. L. Sclater; and the families Indicatoridae, Capitonidae, Cuculidae, and Musophagidae, by G. E. Shelley. Pp. xii., 484: 13 coloured Plates. [With Systematic and Alphabetical Indexes.] 1891, 8vo. 1*l.* 5*s.*

Catalogue of the Birds in the British Museum—*continued*.

Vol. XX. Catalogue of the Psittaci, or Parrots. By T. Salvadori. Pp. xvii., 658. Woodcuts and 18 coloured Plates. [With Systematic and Alphabetical Indexes.] 1891, 8vo. 1*l.* 10*s.*

Vol. XXI. Catalogue of the Columbæ, or Pigeons. By T. Salvadori. Pp. xvii., 676. 15 coloured Plates. [With Systematic and Alphabetical Indexes.] 1893, 8vo. 1*l.* 10*s.*

Vol. XXII. Catalogue of the Game Birds (*Pterocletes*, *Gallinæ*, *Opisthocomi*, *Hemipodii*). By W. R. Ogilvie Grant. Pp. xvi., 585. 8 coloured Plates. [With Systematic and Alphabetical Indexes.] 1893, 8vo. 1*l.* 6*s.*

Vol. XXIII. Catalogue of the Fulicariæ (Rallidæ and Heliornithidæ) and Alektorides (Aramidæ, Eurypygidæ, Mesitidæ, Rhinocetidæ, Gruidæ, Psophiidæ, and Otididæ). By R. Bowdler Sharpe. Pp. xiii., 353. 9 coloured Plates. [With Systematic and Alphabetical Indexes.] 1894, 8vo. 1*l.*

Vol. XXIV. Catalogue of the Limicolæ. By R. Bowdler Sharpe. Pp. xii., 794. Woodcuts and 7 coloured Plates. [With Systematic and Alphabetical Indexes.] 1896, 8vo. 1*l.* 5*s.*

Vol. XXV. Catalogue of the Gaviæ and Tubinares. Gaviæ. (Terns, Gulls, and Skuas), by Howard Saunders. Tubinares (Petrels and Albatrosses), by Osbert Salvin. Pp. xv., 475. Woodcuts and 8 coloured Plates. [With Systematic and Alphabetical Indexes.] 1896, 8vo. 1*l.* 1*s.*

Vol. XXVI. Catalogue of the Plataleæ, Herodiones, Steganopodes, Pygopodes, Alcæ, and Impennes. Plataleæ (Ibises and Spoonbills) and Herodiones (Hérons and Storks), by R. Bowdler Sharpe. Steganopodes (Cormorants, Gannets, Frigate-birds, Tropic-birds, and Pelicans), Pygopodes (Divers and Grebes), Alcæ (Auks), and Impennes (Penguins), by W. R. Ogilvie-Grant. Pp. xvii., 687. Woodcuts and 14 coloured Plates. [With Systematic and Alphabetical Indexes.] 1898, 8vo. 1*l.* 5*s.*

Catalogue of the Birds in the British Museum—*continued*.

Vol. XXVII. Catalogue of the Chenomorphæ (Palamedæ, Phœnicopteri, Anseres), Crypturi, and Ratitæ. By T. Salvadori. Pp. xv., 636. 19 coloured Plates. [With Systematic and Alphabetical Indexes.] 1895, 8vo. 1*l.* 12*s.*

A Hand-list of the Genera and Species of Birds. [Nomenclator Avium tum Fossilium tum Viventium.] By R. Bowdler Sharpe, LL.D. :—

Vol. V. Pp. xx., 678. [With Systematic and Alphabetical Indexes.] 1909, 8vo. 1*l.*

General Index to . . . Volumes I.–V. Edited by W. R. Ogilvie-Grant. Pp. v., 199. 1912, 8vo. 10*s.*

———— Edition on large paper, printed on one side only. Pp. v., 388. 1912, fol. 1*l.*

List of the Specimens of Birds in the Collection of the British Museum. By George Robert Gray :—

Part III., Sections III. and IV. Capitonidæ and Pcidæ. Pp. 137. [With Index.] 1868, 12mo. 1*s.* 6*d.*

Part IV. Columbæ. Pp. 73. [With Index.] 1856, 12mo. 1*s.* 9*d.*

Part V. Gallinæ. Pp. iv., 120. [With an Alphabetical Index.] 1867, 12mo. 1*s.* 6*d.*

Catalogue of the Birds of the Tropical Islands of the Pacific Ocean in the Collection of the British Museum. By George Robert Gray, F.L.S., &c. Pp. 72. [With an Alphabetical Index.] 1859, 8vo. 1*s.* 6*d.*

Catalogue of the Collection of Birds' Eggs in the British Museum (Natural History) :—

Vol. I. Ratitæ. Carinatæ (Tinamiformes—Lariformes). By Eugene W. Oates. Pp. xxiii., 252. 18 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1901, 8vo. 1*l.* 10*s.*

Vol. II. Carinatæ (Charadriiformes—Strigiformes). By Eugene W. Oates. Pp. xx., 400. 15 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1902 8vo. 1*l.* 10*s.*

Catalogue of the Collection of Birds' Eggs in the British Museum (Natural History)—*continued*.

- Vol. III. Carinatae (Psittaciformes — Passeriformes)
By Eugene W. Oates and Capt. Savile G. Reid.
Pp. xxiii., 349. 10 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1903, 8vo. 1l. 5s.
- Vol. IV. Carinatae (Passeriformes continued). By Eugene W. Oates, assisted by Capt. Savile G. Reid.
Pp. xviii., 352. 14 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1905, 8vo. 1l. 10s.
- Vol. V. Carinatae (Passeriformes completed). By W. R. Ogilvie-Grant. Pp. xxiii., 547. 22 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1912, 8vo. 2l. 7s. 6d.

REPTILES.

Catalogue of the Tortoises, Crocodiles, and Amphisbæniæ in the Collection of the British Museum. By Dr. J. E. Gray, F.R.S., &c. Pp. viii., 80. [With an Alphabetical Index.] 1844, 12mo. 1s.

Catalogue of Shield Reptiles in the Collection of the British Museum. By John Edward Gray, F.R.S., &c.:—

Appendix. Pp. 28. 1872, 4to. 2s. 6d.

Part II. Emydosaurians, Rhynchocephalia, and Amphisbæniæ. Pp. vi., 41. 25 Woodcuts. 1872, 4to. 3s. 6d.

Hand-List of the Specimens of Shield Reptiles in the British Museum. By Dr. J. E. Gray, F.R.S., F.L.S., &c. Pp. iv., 124. [With an Alphabetical Index.] 1873, 8vo. 4s.

Catalogue of the Chelonians, Rhynchocephalians, and Crocodiles in the British Museum (Natural History). New Edition. By George Albert Boulenger. Pp. x., 311. 73 Woodcuts and 6 Plates. [With Systematic and Alphabetical Indexes.] 1889, 8vo. 15s.

Catalogue of the Lizards in the British Museum (Natural History). Second Edition. By George Albert Boulenger:—

Vol. II. Iguanidæ, Xenosauridæ, Zonuridæ, Anguidæ, Anniellidæ, Helodermatidæ, Varanidæ, Xantusiidæ,

Catalogue of the Lizards in the British Museum—*continued*.

Teiidae, Amphisbænidae. Pp. xiii., 497. 24 Plates.
[With Systematic and Alphabetical Indexes.] 1885,
8vo. 1*l*.

Vol. III. Lacertidae, Gerrhosauridae, Scincidae, Anelytro-
pidæ, Dibamidæ, Chamæleontidae. Pp. xii., 575. 40
Plates. [With a Systematic Index and an Alphabetical
Index to the three volumes.] 1887, 8vo. 1*l*. 6*s*.

Catalogue of the Snakes in the British Museum (Natural History). By George Albert Boulenger, F.R.S., &c. :—

Vol. I., containing the families Typhlopidae, Glauconiidae,
Boidæ, Ilysiidae, Uropeltidae, Xenopeltidae, and Colu-
bridæ aglyphæ (part). Pp. xiii., 448 : 26 Woodcuts
and 28 Plates. [With Systematic and Alphabetical
Indexes.] 1893, 8vo. 1*l*. 1*s*.

Vol. II., containing the conclusion of the Colubridæ
aglyphæ. Pp. xi., 382 : 25 Woodcuts and 20 Plates.
[With Systematic and Alphabetical Indexes.] 1894,
8vo. 17*s*. 6*d*.

Vol. III., containing the Colubridæ (Opisthoglyphæ and
Proteroglyphæ), Amblycephalidae, and Viperidae.
Pp. xiv., 727 : 37 Woodcuts and 25 Plates. [With
Systematic Index, and Alphabetical Index to the 3
volumes.] 1896, 8vo. 1*l*. 6*s*.

Catalogue of Colubrine Snakes in the Collection of the British Museum. By Dr. Albert Günther. Pp. xvi., 281.
[With Geographic, Systematic, and Alphabetical Indexes.] 1858, 12mo. 4*s*.

BATRACHIANS.

Catalogue of the Batrachia Salientia in the Collection of the British Museum. By Dr. Albert Günther. Pp. xvi., 160.
12 Plates. [With Systematic, Geographic, and Alphabetical Indexes.] 1858, 8vo. 6*s*.

FISHES.

Catalogue of the Fishes in the British Museum. Second Edition. Vol. I. Catalogue of the Perciform Fishes in the British Museum. Vol. I. Containing the Centrarchidae Percidae, and Serranidae (part). By George Albert Boulenger, F.R.S. Pp. xix., 394. Woodcuts and 15 Plates. [With Systematic and Alphabetical Indexes.] 1895, 8vo. 15*s*.

Catalogue of Lophobranchiate Fish in the Collection of the British Museum. By J. J. Kaup, Ph.D., &c. Pp. iv., 80. 4 Plates. [With an Alphabetical Index.] 1856, 12mo. 2s.

Catalogue of the Fresh-water Fishes of Africa in the British Museum (Natural History). By G. A. Boulenger, F.R.S. :—

Vol. I. Pp. xi., 373 : 270 text-figures. [With Systematic and Alphabetical Indexes.] 1909, imp. 8vo. 17. 12s. 6d.

Vol. II. Pp. xii., 529 : 382 text-figures. [With Systematic and Alphabetical Indexes.] 1911, imp. 8vo. 27. 5s.

MOLLUSCA.

Guide to the Systematic Distribution of Mollusca in the British Museum. Part I. By John Edward Gray, Ph.D., F.R.S., &c. Pp. xii., 230. 121 Woodcuts. 1857, 8vo. 5s.

Catalogue of Pulmonata, or Air Breathing Mollusca, in the Collection of the British Museum. Part I. By Dr. Louis Pfeiffer. Pp. iv., 192. Woodcuts. 1855, 12mo. 2s. 6d.

Catalogue of the Auriculidæ, Proserpinidæ, and Truncatellidæ in the Collection of the British Museum. By Dr. Louis Pfeiffer. Pp. iv., 150. Woodcuts. 1857, 12mo. 1s. 9d.

List of the Mollusca in the Collection of the British Museum. By John Edward Gray, Ph.D., F.R.S., &c. :—

Part II. Olividæ. Pp. 41. 1865, 12mo. 1s.

Catalogue of the Conchifera, or Bivalve Shells, in the Collection of the British Museum. By M. Deshayes :—

Part I. Veneridæ, Cyprinidæ, Glauconomidæ, and Petricoladæ. Pp. iv., 216. 1853, 12mo. 3s.

Part II. Petricoladæ (concluded); Corbiculadæ Pp. 217-292. [With an Alphabetical Index to the two parts.] 1854, 12mo. 6d.

BRACHIOPODA.

Catalogue of Brachiopoda Ancylopoda or Lamp Shells in the Collection of the British Museum. [Issued as "Catalogue of the Mollusca, Part IV."] Pp. iv., 128. 25 Woodcuts. [With an Alphabetical Index.] 1853, 12mo. 3s.

POLYZOA.

Catalogue of Marine Polyzoa in the Collection of the British Museum. Part III. Cyclostomata. By George Busk, F.R.S. Pp. viii., 39. 38 plates. [With a Systematic Index.] 1875, 8vo. 5s.

CRUSTACEA.

Catalogue of the Specimens of Amphipodous Crustacea in the Collection of the British Museum. By C. Spence Bate, F.R.S., &c. Pp. iv., 399. 58 Plates. [With an Alphabetical Index.] 1862, 8vo. 1*l.* 5*s.*

ARACHNIDA.

Descriptive Catalogue of the Spiders of Burma, based upon the Collection made by Eugene W. Oates and preserved in the British Museum. By T. Thorell. Pp. xxxvi., 406. [With Systematic List and Alphabetical Index.] 1895, 8vo. 10*s.* 6*d.*

INSECTS.

Coleopterous Insects.

Nomenclature of Coleopterous Insects in the Collection of the British Museum :—

Part VII. Longicornia, I. By Adam White. Pp. iv., 174. 4 Plates. 1853, 12mo. 2*s.* 6*d.*

Part VIII. Longicornia, II. By Adam White. Pp. 237. 6 Plates. 1855, 12mo. 3*s.* 6*d.*

Illustrations of Typical Specimens of Coleoptera in the Collection of the British Museum. Part I. Lycidæ. By Charles Owen Waterhouse. Pp. x., 83. 18 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1879, 8vo. 16*s.*

Catalogue of the Coleopterous Insects of Madeira in the Collection of the British Museum. By T. Vernon Wollaston, M.A., F.L.S. Pp. xvi., 234 : 1 Plate. [With a Topographical Catalogue and an Alphabetical Index.] 1857, 8vo. 3*s.*

Catalogue of the Coleopterous Insects of the Canaries in the Collection of the British Museum. By T. Vernon Wollaston, M.A., F.L.S. Pp. xiii., 648. [With Topographical and Alphabetical Indexes.] 1864, 8vo. 10*s.* 6*d.*

Catalogue of Halticidæ in the Collection of the British Museum. By the Rev. Hamlet Clark, M.A., F.L.S. Physapodes and Œdipodes. Part I. Pp. xii., 301. Frontispiece and 9 Plates. 1860, 8vo. 7*s.*

Catalogue of Hispidæ in the Collection of the British Museum. By Joseph S. Baly, M.E.S., &c. Part I. Pp. x., 172. 9 Plates. [With an Alphabetical Index.] 1858, 8vo. 6s.

Hymenopterous Insects.

Catalogue of Hymenopterous Insects in the Collection of the British Museum. By Frederick Smith. 12mo. :—

Part II. Apidæ. Pp. 199–465. 6 Plates. [With an Alphabetical Index.] 1854. 6s.

Part III. Mutillidæ and Pompilidæ. Pp. 206. 6 Plates. 1855. 6s.

Part IV. Sphegidæ, Larridæ, and Crabronidæ. Pp. 207–497. 6 Plates. [With an Alphabetical Index.] 1856. 6s.

Part V. Vespidæ. Pp. 147. 6 Plates. [With an Alphabetical Index.] 1857. 6s.

Part VII. Dorylidæ and Thynnidæ. Pp. 76. 3 Plates. [With an Alphabetical Index.] 1859. 2s.

List of Hymenoptera, with descriptions and figures of the Typical Specimens in the British Museum. Vol. I., Tenthredinidæ and Siricidæ. By W. F. Kirby. Pp. xxviii., 450. 16 Coloured Plates. [With Systematic and Alphabetical Indexes.] 1882, 8vo. 17. 18s.

A Revision of the Ichneumonidæ, based on the Collection in the British Museum (Natural History). With descriptions of new Genera and Species. By Claude Morley, F.Z.S., F.E.S. :—

Part I. Tribes Ophionides and Metopiides. Pp. xi., 88 : 1 Coloured Plate. [With Systematic and Alphabetical Indexes.] 1912, 8vo. 4s.

Part II. Tribes Rhyssides, Echthromorphides, Anomalides, and Paniscides. Pp. viii., 140. 1 Coloured Plate. [With Systematic and Alphabetical Indexes, &c.] 1913, 8vo. 5s. 6d.

Dipterous Insects.

A Monograph of the Culicidæ, or Mosquitoes. Mainly compiled from the Collections received at the British Museum from various parts of the world in connection with the Investigation into the cause of Malaria conducted by the

Colonial Office and the Royal Society. By Fred. V. Theobald, M.A., &c. :—

Vol. III. Pp. xvii., 359 : 17 plates, 1 diagram, and 193 illustrations in text. 1903, 8vo. 1*l.* 1*s.*

Vol. IV. Pp. xix., 639 : 16 plates and 297 text-figures. [With Index.] 1907, 8vo. 1*l.* 12*s.* 6*d.*

Vol. V. Pp. xv., 646 : 6 plates and 261 text-figures. [With Index.] 1910, 8vo. 1*l.* 5*s.*

Handbook of the Tsetse-Flies [Genus *Glossina*]. By Ernest Edward Austen. With 10 coloured plates and 24 text-figures, by A. J. Engel Terzi, and 1 map. Pp. x., 110. [With Index.] 1911, roy. 8vo. 5*s.* 6*d.*

Illustrations of African Blood-sucking Flies other than Mosquitoes and Tsetse-Flies. By Ernest Edward Austen, with coloured figures by Grace Edwards. Pp. xv., 221 : 13 coloured plates, 3 text-figures. 1909, roy. 8vo. 1*l.* 7*s.* 6*d.*

Lepidopterous Insects.

Catalogue of the Lepidoptera Phalænæ in the British Museum. By Sir George F. Hampson, Bart. :—

Vol. I. Catalogue of the Syntomidæ. Pp. xxi., 559 : 285 woodcuts. [With Systematic and Alphabetical Indexes.] 1898, 8vo. 15*s.*

—Atlas of 17 Coloured Plates, 8vo. 15*s.*

Vol. II. Catalogue of the Arctiadæ (Nolinæ, Lithosianæ). Pp. xx., 589 : 411 woodcuts. [With Systematic and Alphabetical Indexes.] 1900, 8vo. 18*s.*

—Atlas of 18 Coloured Plates (xviii.-xxxv.), 8vo. 15*s.*

Vol. III. Catalogue of the Arctiadæ (Arctianæ) and Agaristidæ. Pp. xix., 690 : 294 woodcuts. [With Systematic and Alphabetical Indexes.] 1901, 8vo. 15*s.*

—Atlas of 19 Coloured Plates (xxxvi.-liv.), 8vo. 16*s.*

Vol. IV. Catalogue of the Noctuidæ [Agrotinæ]. Pp. xx., 689 : 125 woodcuts. [With Systematic and Alphabetical Indexes.] 1903, 8vo. 15*s.*

—Atlas of 23 Coloured Plates (lv.-lxxvii.), 8vo. 16*s.*

Vol. V. Catalogue of the Noctuidæ [Hadeninæ]. Pp. xvi., 634 : 172 woodcuts. [With Systematic and Alphabetical Indexes.] 1905, 8vo. 15*s.*

—Atlas of 18 Coloured Plates (lxxviii.-xcv.), 8vo. 15*s.*

Catalogue of Lepidoptera Phalaenæ—*continued*.

Vol. VI. Catalogue of the Noctuidæ [Cucullianæ]. Pp. xiv., 532 : 172 woodcuts. [With Systematic and Alphabetical Indexes.] 1906, 8vo. 15s.

—Atlas of 12 Coloured Plates (xcvi.-cvii.), 8vo. 10s.

Vol. VII. Catalogue of the Noctuidæ [Acronyctinæ]. Pp. xv., 709 : 184 woodcuts. [With Systematic and Alphabetical Indexes.] 1908, 8vo. 17s.

—Atlas of 15 Coloured Plates (cviii.-cxxii.), 8vo. 13s.

Vol. VIII. Catalogue of the Noctuidæ [Acronyctinæ, II.]. Pp. xiv., 583 : 162 woodcuts. [With Table of the Phylogeny of the Acronyctinæ, and Systematic and Alphabetical Indexes.] 1909, 8vo. 15s.

—Atlas of 14 Coloured Plates (cxxiii.-cxxxvi.), 8vo. 12s.

Vol. IX. Catalogue of the Noctuidæ [Acronyctinæ, III.]. Pp. xv., 552 : 247 woodcuts. [With Table of the Phylogeny of the Acronyctinæ, and Systematic and Alphabetical Indexes.] 1910, 8vo. 15s.

—Atlas of 11 Coloured Plates (cxxxvii.-cxlvii.), 8vo. 12s.

Vol. X. Catalogue of the Noctuidæ [Eratrionæ]. Pp. xix., 829 : 214 woodcuts. [With Table of the Phylogeny of the Eratrionæ, and Systematic and Alphabetical Indexes.] 1910, 8vo. 17s.

—Atlas of 26 Coloured Plates (cxlviii.-clxxiii.). 1911, 8vo. 17s.

Vol. XI. Catalogue of the Noctuidæ [Sub-families Eutelianæ, Stictopterinae, Sarrothripinae, and Acontianæ.]. Pp. xvii., 689 : 275 woodcuts. [With Systematic and Alphabetical Indexes.] 1912, 8vo. 17s.

—Atlas of 16 Coloured Plates (clxxiv.-cxc.), 8vo. 17s. 6d.

Vol. XII. Catalogue of the Noctuidæ [Sub-family Catocalinae, part]. Pp. xiii., 626 : 134 woodcuts. [With Table of the Phylogeny of the Catocalinae, and Systematic and Alphabetical Indexes.] 1913, 8vo. 17s. 6d.

—Atlas of 30 Coloured Plates (cxcii.-ccxxi.), 8vo. 17s. 5s.

Illustrations of Typical Specimens of Lepidoptera Heterocera
in the Collection of the British Museum:—

Part V. By Arthur Gardiner Butler. Pp. xii., 74.
78–100 Coloured Plates. [With a Systematic Index.]
1881, 4to. 2l. 10s.

Part VI. By Arthur Gardiner Butler. Pp. xv., 89.
101–120 Coloured Plates. [With a Systematic Index.]
1886, 4to. 2l. 4s.

Part VII. By Arthur Gardiner Butler. Pp. iv., 124.
121–138 Coloured Plates. [With a Systematic List.]
1889, 4to. 2l.

Part VIII. The Lepidoptera Heterocera of the Nilgiri
District. By George Francis Hampson. Pp. iv., 144.
139–156 Coloured Plates. [With a Systematic List.]
1891, 4to. 2l.

Part IX. The Macrolepidoptera Heterocera of Ceylon.
By George Francis Hampson. Pp. v., 182. 157–176
Coloured Plates. [With a General Systematic List of
Species collected in, or recorded from, Ceylon.] 1893.
4to. 2l. 2s.

Catalogue of the Collection of Palæarctic Butterflies formed
by the late John Henry Leech, and presented to the
Trustees of the British Museum by his Mother, Mrs. Eliza
Leech. By Richard South, F.E.S. Pp. vi., 228. 2 Coloured
Plates. With a Portrait and Biographical Memoir of Mr.
Leech. 1902, 4to. 1l.

Catalogue of Diurnal Lepidoptera described by Fabricius in
the Collection of the British Museum. By Arthur Gardiner
Butler, F.L.S., &c. Pp. iv., 303. 3 Plates. 1869, 8vo. 7s. 6d.

List of the Specimens of Lepidopterous Insects in the
Collection of the British Museum. By Francis Walker.
12mo. :—

Part XXVIII. Tortricites and Tineites. Pp. 287–561.
1863. 4s.

Part XXXI. Supplement. Pp. 1–321. 1864. 5s.

Part XXXIII. ————— Part 3. Pp. 707–1120.
1865. 6s.

Neuropterous Insects.

Catalogue of the Specimens of Neuropterous Insects in the
Collection of the British Museum. By Dr. H. Hagen.
Part I. Termitina. Pp. 34. 1858, 12mo. 6d.

Orthopterous Insects.

Catalogue of Orthopterous Insects in the Collection of the British Museum. Part I. Phasmidæ. By John Obadiah Westwood, F.L.S., &c. Pp. 195. 48 Plates. [With an Alphabetical Index.] 1859, 4to. 3l.

Catalogue of the Specimens of Blattariæ in the Collection of the British Museum. By Francis Walker, F.L.S., &c. Pp. 239. [With an Alphabetical Index.] 1868, 8vo. 5s. 6d.

Catalogue of the Specimens of Dermaptera Saltatoria in the Collection of the British Museum. By Francis Walker, F.L.S., &c. :—

Part II. Locustidæ (continued). Pp. 225–423. [With an Alphabetical Index.] 1869, 8vo. 4s. 6d.

Part III. Locustidæ (continued).—Acrididæ. Pp. 425–604. [With an Alphabetical Index.] 1870, 8vo. 4s.

Part IV. Acrididæ (continued). Pp. 605–809. [With an Alphabetical Index.] 1870, 8vo. 6s.

Part V. Tettigidæ.—Supplement to the Catalogue of Blattariæ.—Supplement to the Catalogue of Dermaptera Saltatoria (with remarks on the Geographical Distribution of Dermaptera). Pp. 811–850; 43; 116. [With Alphabetical Indexes.] 1870, 8vo. 6s.

Synonymic Catalogue of Orthoptera. By W. F. Kirby :—

Vol. I. Orthoptera Euplexoptera, Cursoria, et Gressoria. (Forficulidæ, Hemimeridæ, Blattidæ, Mantidæ, Phasmidæ.) Pp. x., 501. [With Index.] 1904, 8vo. 10s.

Vol. II. Orthoptera Saltatoria, Part I. (Achetidæ et Phasgonuridæ.) Pp. viii., 562. [With Index.] 1906, 8vo. 15s.

Vol. III. Orthoptera Saltatoria, Part II. (Locustidæ vel Acridiidæ.) Pp. vii., 674. [With Index.] 1910, 8vo. 1l.

Homopterous Insects.

A Synonymic Catalogue of Homoptera. Part I. Cicadidæ By W. L. Distant. Pp. 207. [Index.] 1906, 8vo. 5s.

VERMES.

Catalogue of the Species of Entozoa, or Intestinal Worms, contained in the Collection of the British Museum. By Dr. Baird. Pp. iv., 132. 2 Plates. [With an Index of the Animals in which the Entozoa mentioned in the Catalogue are found, and an Index of Genera and Species.] 1853, 12mo. 2s.

Catalogue of the Chætopoda in the British Museum (Natural History). A. Polychæta: Part I.—Arenicolidæ. By J. H. Ashworth, D.Sc. Pp. xii., 175. 15 Plates, 68 Text-figures. [With Systematic and General Indexes, List of Text-figures, Description of Plates, &c.] 1912, roy. 8vo. 1l. 7s. 6d.

ANTHOZOA.

Catalogue of Sea-pens or Pennatulariidæ in the Collection of the British Museum. By J. E. Gray, F.R.S., &c. Pp. iv., 40. 2 Woodcuts. 1870, 8vo. 1s. 6d.

Catalogue of Lithophytes or Stony Corals in the Collection of the British Museum. By J. E. Gray, F.R.S., &c. Pp. iv., 51. 14 Woodcuts. 1870, 8vo. 3s.

Catalogue of the Madreporarian Corals in the British Museum (Natural History):—

Vol. I. The Genus Madrepora. By George Brook. Pp. xi., 212. 35 Collotype Plates. [With Systematic and Alphabetical Indexes, and Explanation of the Plates.] 1893, 4to. 1l. 4s.

Vol. II. The Genus Turbinaria; the Genus Astræopora. By Henry M. Bernard, M.A. Cantab., F.L.S., F.Z.S. Pp. iv., 106. 30 Collotype and 3 Lithographic Plates. [With Index of Generic and Specific Names, and Explanation of the Plates.] 1896, 4to. 18s.

Vol. III. The Genus Montipora; the Genus Anacropora. By Henry M. Bernard, M.A., &c. Pp. vii., 192. 30 Collotype and 4 Lithographic Plates. [With Systematic Index, Index of Generic and Specific Names, and Explanation of the Plates.] 1897, 4to. 1l. 4s.

Vol. IV. The Family Poritidæ. I.—The Genus Goniopora. By Henry M. Bernard, M.A. Pp. viii., 206. 12 Collotype and 4 Lithographic Plates. [With Index of Generic and Specific Names, and Explanation of the Plates.] 1903, 4to. 1l.

Vol. V. The Family Poritidæ. II.—The Genus Porites. Part I.—Porites of the Indo-Pacific Region. By Henry M. Bernard, M.A. Pp. vi., 303. 35 Plates. [With Index of Generic and Specific Names and Explanation of the Plates.] 1905, 4to. 1l. 15s.

Catalogue of the Madreporarian Corals in the British Museum (Natural History)—*continued*.

Vol. VI. The Family Poritidæ. II.—The Genus *Porites*. Part II.—*Porites* of the Atlantic and West Indies, with the European Fossil Forms. The Genus *Goniopora*, a supplement to Vol. IV. By Henry M. Bernard, M.A. Pp. vi., 173. 16 Collotype and 1 Lithographic Plates. [With Index of Generic and Specific Names, and Explanation of the Plates.] 1906, 4to. 1*l*.

BRITISH ANIMALS.

Catalogue of British Birds in the Collection of the British Museum. By George Robert Gray, F.L.S., F.Z.S., &c. Pp. xii., 248. [With a List of Species.] 1863, 8vo. 3*s*. 6*d*.

Catalogue of the British Species of *Pisidium* (Recent and Fossil) in the Collections of the British Museum (Natural History), with Notes on those of Western Europe. By B. B. Woodward, F.L.S., &c. Pp. ix., 144. 30 Plates. [With Bibliography and Index.] 1913, 8vo. 10*s*. 6*d*.

Catalogue of British Hymenoptera in the Collection of the British Museum. Second edition. Part I. *Andrenide* and *Apidæ*. By Frederick Smith, M.E.S. New issue Pp. xi. 236. 11 Plates. [With Systematic and Alphabetical Index.] 1891, 8vo. 6*s*.

Catalogue of British Fossorial Hymenoptera, *Formicidæ*, and *Vespidæ* in the Collection of the British Museum. By Frederick Smith, V.P.E.S. Pp. 236. 6 Plates. [With an Alphabetical Index.] 1858, 12mo. 6*s*.

Catalogue of British Hymenoptera of the Family *Chalcididæ*. By Claude Morley, F.Z.S., F.E.S. Pp. 74. [Index.] 1910, 8vo. 3*s*. 6*d*.

Illustrations of British Blood-sucking Flies, with notes by Ernest Edward Austen, Assistant, Department of Zoology, British Museum (N.H.). Pp. 74. 34 Coloured Plates. 1906, roy. 8vo. 1*l*. 5*s*.

A Catalogue of the British Non-parasitical Worms in the Collection of the British Museum. By George Johnston, M.D., Edin., F.R.C.L., Ed., LL.D., Marischal Coll., Aberdeen, &c. Pp. 365. Woodcuts and 24 Plates. [With an Alphabetical Index.] 1865, 8vo. 7*s*.

Catalogue of the British Echinoderms in the British Museum (Natural History). By F. Jeffrey Bell, M.A. Pp. xvii., 202. Woodcuts and 16 Plates (2 Coloured). [With Table of Contents, Tables of Distribution, Alphabetical Index, Description of the Plates, &c.] 1892, 8vo. 12s. 6d.

List of the Specimens of British Animals in the Collection of the British Museum; with Synonyma and References to figures. 12mo.:—

Part V. Lepidoptera. By J. F. Stephens. 2nd Edition. Revised by H. T. Stainton and E. Shepherd. Pp. iv., 224. 1856. 1s. 9d.

Part VI. Hymenoptera. By F. Smith. Pp. 134. 1851. 2s.

Part VII. Mollusca, Acephala and Brachiopoda. By Dr. J. E. Gray. Pp. iv., 167. 1851. 3s. 6d.

Part VIII. Fish. By Adam White. Pp. xxiii., 164. (With Index and List of Donors.) 1851. 3s. 6d.

Part XI. Anoplura, or Parasitic Insects. By H. Denny. Pp. iv., 51. 1852. 1s.

Part XII. Lepidoptera (continued). By James F. Stephens. Pp. iv., 54. 1852. 9d.

Part XIII. Nomenclature of Hymenoptera. By Frederick Smith. Pp. iv., 74. 1853. 1s. 4d.

Part XIV. Nomenclature of Neuroptera. By Adam White. Pp. iv., 16. 1853. 6d.

Part XV. Nomenclature of Diptera, I. By Adam White. Pp. iv., 42. 1853. 1s.

Part XVI. Lepidoptera (completed). By H. T. Stainton. Pp. 199. [With an Index.] 1854. 3s.

PLANTS.

Illustrations of Australian Plants collected in 1770 during Captain Cook's Voyage round the World in H.M.S. "Endeavour." By the Right Hon. Sir Joseph Banks, Bart., K.B., P.R.S., and Dr. Daniel Solander, F.R.S. [Being a series of lithographic reproductions of copper-plates engraved after paintings by F. P. Nodder, James Miller, J. F. Miller, and John Cleveley.] With Introduction and Determinations by James Britten, F.L.S., Senior Assistant, Department of Botany, British Museum:—

Part I.—101 Plates, with 31 pages of descriptive text. 1900, fol. 17. 5s.

Part II.—142 Plates (pls. 101-243), with 41 pages of descriptive text (pp. 35-75). 1901, fol. 17. 15s.

Illustrations of Australian Plants &c.—*continued*.

Part III.—77 Plates (pls. 244–318, 45A, and 122), with 26 pages of descriptive text, including Index to the whole work (pp. 77–102), and 3 maps. 1905, fol. 17. 5s.

Catalogue of the African Plants collected by Dr. Friedrich Welwitsch in 1853–61 :—

Vol. I. Dicotyledons. By William Philip Hiern, M.A. F.L.S., &c. :—

Part I. [Ranunculaceæ to Rhizophoraceæ.] Pp. xxvi., 336. [With Portrait of Dr. Welwitsch. Introduction, Bibliography, and Index of Genera.] 1896. 8vo. 7s. 6d.

Part II. Combretaceæ to Rubiaceæ. Pp. 337–510. [With Index of Genera.] 1898, 8vo. 4s.

Part III. Dipsacæ to Scrophulariaceæ. Pp. 511–784. [With Index of Genera.] 1898, 8vo. 5s.

Part IV. Lentibulariaceæ to Ceratophylleæ. Pp. 785–1035. [With Index.] 1900, 8vo. 5s.

Vol. II. Monocotyledons, Gymnosperms, and Cryptogams :—

Part I. Monocotyledons and Gymnosperms. By Alfred Barton Rendle, M.A., D.Sc., F.L.S., Assistant, Department of Botany. Pp. 260. [With Index of Genera.] 1899, 8vo. 6s.

Part II. Cryptogamia. Pp. 261–566. [With Table of Errata, and General Index to the whole work.] 1901, 8vo. 6s.

Vascular Cryptogams	...	By William Carruthers, F.R.S.
Mosses	...	Antony Gepp, M.A., F.L.S.
Hepatics	...	F. Stephani.
Marine Algæ	...	Ethel S. Barton.
Freshwater Algæ	...	W. West, F.L.S., and G. S. West, B.A.
Diatomaceæ	...	Thomas Comber, F.L.S.
Lichenes	...	E. A. Wainio.
Fungi	...	Annie Lorrain Smith.
Mycetozoa	...	Arthur Lister, F.R.S.

Catalogue of the Plants collected by Mr. and Mrs. P. A. Talbot in the Oban District, South Nigeria. By A. B. Rendle, M.A., D.Sc., F.R.S., E. G. Baker, F.L.S., H. F. Wernham, B.Sc., S. Moore, F.L.S., and others. Pp. x., 157 : 17 Plates. [With Index and List of Plates.] 1913, 8vo. 9s.

Flora of Jamaica, containing descriptions of the Flowering Plants known from the Island. By William Fawcett, B.Sc., F.L.S., etc., and A. B. Rendle, M.A., D.Sc., F.R.S., F.L.S., etc. Vol. I. Orchidaceæ. Pp. xx., 150 : 32 Plates. [With Index of Genera and Species.] 1910, 8vo. 10s. 6d.

Synopsis of the British Basidiomycetes: a Descriptive Catalogue of the Drawings and Specimens in the Department of Botany, British Museum. By Worthington George Smith, F.L.S. Pp. 531. 5 Plates and 145 Figures in Text. [With Index.] 1908, 8vo. 10s.

A Monograph of the British Lichens: a Descriptive Catalogue of the Species in the Department of Botany, British Museum. Part II. By Annie Lorrain Smith, F.L.S. Pp. [viii.,] 409: 59 Plates. [With List of Plates, Glossary, and Index.] 1911, 8vo. 1l.

A Monograph of the Mycetozoa: a Descriptive Catalogue of the Species in the Herbarium of the British Museum. By Arthur Lister, F.R.S., F.L.S. Second Edition, revised by Gulielma Lister, F.L.S. Pp. 302. 201 Plates (120 coloured). 56 Woodcuts. [With Indexes, Bibliography, Glossary, etc.] 1911, 8vo. 1l. 10s.

List of British Diatomaceæ in the Collection of the British Museum. By the Rev. W. Smith, F.L.S., &c. Pp. iv., 55. 1859, 12mo. 1s.

FOSSILS.

Catalogue of the Fossil Mammalia in the British Museum (Natural History). By Richard Lydekker, B.A., F.G.S.:—

Part I. Containing the Orders Primates, Chiroptera, Insectivora, Carnivora, and Rodentia. Pp. xxx., 268. 33 Woodcuts. [With Systematic and Alphabetical Indexes.] 1885, 8vo. 5s.

Part II. Containing the Order Ungulata, Suborder Artiodactyla. Pp. xxii., 324. 39 Woodcuts. [With Systematic and Alphabetical Indexes.] 1885, 8vo. 6s.

Part III. Containing the Order Ungulata, Suborders Perissodactyla, Toxodontia, Condylarthra, and Amblypoda. Pp. xvi., 186. 30 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1886, 8vo. 4s.

Part IV. Containing the Order Ungulata, Suborder Proboscidea. Pp. xxiv., 235. 32 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1886, 8vo. 5s.

Part V. Containing the Group Tillodontia, the Orders Sirenia, Cetacea, Edentata, Marsupialia, Monotremata, and Supplement. Pp. xxxv., 345. 55 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1887, 8vo. 6s.

Catalogue of the Fossil Birds in the British Museum (Natural History). By Richard Lydekker, B.A. Pp. xxvii., 368. 75 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1891, 8vo. 10s. 6d.

Catalogue of the Fossil Reptilia and Amphibia in the British Museum (Natural History). By Richard Lydekker, B.A., F.G.S. :—

Part I. Containing the Orders Ornithosauria, Crocodilia, Dinosauria, Squamata, Rhynchocephalia, and Proterosauria. Pp. xxviii., 309. 69 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1888, 8vo. 7s. 6d.

Part II. Containing the Orders Ichthyopterygia and Sauropterygia. Pp. xxi., 307. 85 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1889, 8vo. 7s. 6d.

Part III. Containing the Order Chelonina. Pp. xviii., 239. 53 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species including Synonyms.] 1889, 8vo. 7s. 6d.

Part IV. Containing the Orders Anomodontia, Ecaudata, Caudata, and Labyrinthodontia; and Supplement. Pp. xxiii., 295. 66 Woodcuts. [With Systematic Index, Alphabetical Index of Genera and Species, including Synonyms, and Alphabetical Index of Genera and Species to the entire work.] 1890, 8vo. 7s. 6d.

A descriptive Catalogue of the Marine Reptiles of the Oxford Clay. Based on the Leeds Collection in the British Museum (Natural History), London. By C. W. Andrews, D.Sc., F.R.S. :—

Part I. Families Ophthalmosauridae, Elasmosauridae. Pp. xxiii., 205: 9½ Text-figures, 11 Plates. [With Systematic and Alphabetical Indexes, List of Illustrations, Explanations of Plates, &c.] 1910, 4to. 1l. 5s.

Part II. Families Pliosauridae, Teleosauridae, and Geosauridae. Pp. xxiv., 206: 73 Text-figures, 14 Plates. [With Systematic and Alphabetical Indexes, List of Illustrations in Text, Explanations of Plates, &c.] 1913, 4to. 1l. 5s.

Catalogue of the Fossil Fishes in the British Museum (Natural History). By Arthur Smith Woodward, LL.D., F.R.S., F.G.S., &c. :—

Part I. Containing the Elasmobranchii. Pp. xlvii., 474. 13 Woodcuts and 17 Plates. [With Alphabetical Index, and Systematic Index of Genera and Species.] 1889, 8vo. 1l. 1s.

Part II. Containing the Elasmobranchii (Acanthodii), Holocephali, Ichthyodermulites, Ostracodermi, Dipnoi, and Teleostomi (Crossopterygii and Chondrosteian Actinopterygii). Pp. xlv., 567. 58 Woodcuts and 16 Plates. [With Alphabetical Index, and Systematic Index of Genera and Species.] 1891, 8vo. 1l. 1s.

Catalogue of the Fossil Fishes in the British Museum (Natural History)—*continued*.

Part III. Containing the Actinopterygian Teleostomi of the Orders Chondrostei (concluded), Protospondyli, Aethospondyli, and Isospondyli (in part). Pp. xlii., 544. 45 Woodcuts and 18 Plates. [With Alphabetical Index, and Systematic Index of Genera and Species.] 1895, 8vo. 1*l.* 1*s.*

Part IV. Containing the Actinopterygian Teleostomi of the Suborders Isospondyli (in part), Ostariophysii, Apodes, Percesoces, Hemibranchii, Acanthopterygii, and Anacanthini. Pp. xxxix., 636. 22 Woodcuts and 19 Plates. [With Alphabetical Index, and Systematic Index of Genera and Species.] 1901, 8vo. 1*l.* 1*s.*

A descriptive Catalogue of the Tertiary Vertebrata of the Fayûm, Egypt. Based on the Collection of the Egyptian Government in the Geological Museum, Cairo, and on the Collection in the British Museum (Natural History), London. By C. W. Andrews, D.Sc. Pp. xxxvii., 324 : 98 Text Figures and 26 Plates. [With Systematic and Alphabetical Indexes.] 1906, 4to. 1*l.* 15*s.*

Systematic List of the Edwards Collection of British Oligocene and Eocene Mollusca in the British Museum (Natural History), with references to the type-specimens from similar horizons contained in other collections belonging to the Geological Department of the Museum. By Richard Bullen Newton, F.G.S. Pp. xxviii., 365. [With table of Families and Genera, Bibliography, Correlation-table, Appendix, and Alphabetical Index.] 1891, 8vo. 6*s.*

Catalogue of Tertiary Mollusca in the Department of Geology, British Museum (Natural History). Part I. The Australasian Tertiary Mollusca. By George F. Harris, F.G.S., &c. Pp. xxvi., 407. 8 Plates. [With Table of Families, Genera, and Sub-Genera, and Index.] 1897, 8vo. 10*s.*

Catalogue of the Fossil Cephalopoda in the British Museum (Natural History) :—

Part I. Containing part of the Suborder Nautiloidea, consisting of the families Orthoceratidæ, Endoceratidæ, Actinoceratidæ, Gomphoceratidæ, Ascoceratidæ, Poterioceratidæ, Cyrtoceratidæ, and Supplement. By Arthur H. Foord, F.G.S. Pp. xxxi., 344. 51 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1888, 8vo. 10*s.* 6*d.*

Catalogue of the Fossil Cephalopoda in the British Museum
(Natural History)—*continued*.

Part II. Containing the remainder of the Suborder Nautiloidea, consisting of the families Lituitidæ, Trochoceratidæ, Nautilidæ, and Supplement. By Arthur H. Foord, F.G.S. Pp. xxviii., 407. 86 Woodcuts. [With Systematic Index, and Alphabetical Index of Genera and Species, including Synonyms.] 1891, 8vo. 15s.

Part III. Containing the Bactritidæ, and part of the Suborder Ammonoidea. By Arthur H. Foord, Ph.D., F.G.S., and George Charles Crick, A.R.S.M., F.G.S. Pp. xxxiii., 303. 146 Woodcuts. [With Systematic Index of Genera and Species, and Alphabetical Index.] 1897, 8vo. 12s. 6d.

List of the Types and Figured Specimens of Fossil Cephalopoda in the British Museum (Natural History). By G. C. Crick, F.G.S. Pp. 103. [With Index.] 1898, 8vo. 2s. 6d.

A Catalogue of British Fossil Crustacea, with their Synonyms and the Range in Time of each Genus and Order. By Henry Woodward, F.R.S. Pp. xii., 155. [With an Alphabetical Index.] 1877, 8vo. 5s.

Catalogue of the Fossil Bryozoa in the Department of
Geology, British Museum (Natural History):—

The Jurassic Bryozoa. By J. W. Gregory, D.Sc., F.G.S., F.Z.S. Pp. [viii.,] 239 : 22 Woodcuts and 11 Plates. [With List of Species and Distribution, Bibliography, Index, and Explanation of Plates.] 1896, 8vo. 10s.

The Cretaceous Bryozoa. By J. W. Gregory, D.Sc., F.R.S., &c. :—

Vol. I. Pp. xiv., 457 : 64 Woodcuts and 17 Plates. [With Index and Explanation of Plates.] 1899, 8vo. 16s.

Vol. II. Pp. xlviii., 346. 75 Woodcuts and 9 Plates. [With List of Localities, Bibliography, Subject and Systematic Indexes, and Explanation of Plates.] 1909, 8vo. 13s.

Catalogue of the Blastoida in the Geological Department of the British Museum (Natural History), with an account of the morphology and systematic position of the group, and a revision of the genera and species. By Robert Etheridge, jun., of the Department of Geology, British Museum (Natural History), and P. Herbert Carpenter, D.Sc., F.R.S.,

F.L.S. (of Eton College). Pp. xv., 322. 20 Plates. [With Preface by Dr. H. Woodward, Table of Contents, General Index, Explanations of the Plates, &c.] 1886, 4to. 1*l.* 5*s.*

The Genera and Species of Blastoidea, with a List of the Specimens in the British Museum (Natural History). By F. A. Bather, M.A., F.G.S., of the Geological Department. Pp. x., 70. 1 Woodcut. 1899, 8vo. 3*s.*

Catalogue of the Palæozoic Plants in the Department of Geology and Palæontology, British Museum (Natural History). By Robert Kidston, F.G.S. Pp. viii., 288. [With a list of works quoted, and an Index.] 1886, 8vo. 5*s.*

Catalogue of the Mesozoic Plants in the Department of Geology, British Museum (Natural History). By A. C. Seward, M.A., F.R.S., F.G.S., University Lecturer in Botany and Fellow of Emanuel College, Cambridge:—

Part I. The Wealden Flora. Part I. Thallophyta—Pteridophyta. Pp. xxxviii., 179. 17 Woodcuts and 11 Plates. [With Alphabetical Index, Explanations of the Plates, &c.] 1894, 8vo. 10*s.*

Part II. The Wealden Flora. Part II. Gymnospermæ. Pp. viii., 259. 9 Woodcuts and 20 Plates. [With Alphabetical Index, Explanations of the Plates, &c.] 1895, 8vo. 15*s.*

Part III. The Jurassic Flora. Part I. The Yorkshire Coast. Pp. xii., 341. 53 Woodcuts and 21 Plates. [With Alphabetical Index, Explanations of the Plates, &c.] 1900, 8vo. 1*l.*

Part IV. The Jurassic Flora. II.—Liassic and Oolitic Floras of England (excluding the Inferior Oolite Plants of the Yorkshire Coast). Pp. xv., 192. 20 Woodcuts and 13 Plates. [With Alphabetical Index, Explanations of the Plates, &c.] 1904, 8vo. 10*s.*

Catalogue of the Fossil Plants of the Glossopteris Flora in the Department of Geology, British Museum (Natural History). Being a Monograph of the Permo-carboniferous Flora of India and the Southern Hemisphere. By E. A. Newell Arber, M.A., F.L.S., F.G.S. Pp. lxxiv., 255: 51 Text-Figures and 8 Plates. [With Bibliography and Alphabetical Index.] 1905, 8vo. 12*s.* 6*d.*

GUIDE-BOOKS, ETC.

- General Guide to the British Museum (Natural History), Cromwell Road, London, S.W. Thirteenth Edition. With 58 text-figures, 2 plans, 2 views of the building, and an illustrated cover. Pp. x., 121. 1913, 8vo. 3*d*.
- Guide to the Specimens illustrating the Races of Mankind (Anthropology), exhibited in the Department of Zoology, British Museum (Natural History). Second Edition. [By R. Lydekker, F.R.S.] Illustrated by 16 Figures. Pp. 35. 1912, 8vo. 4*d*.
- Guide to the Galleries of Mammals (other than Ungulates) in the Department of Zoology of the British Museum (Natural History). Eighth Edition. Pp. 101. 52 Woodcuts and 4 plans. Index. 1906, 8vo. 6*d*.
- Guide to the Specimens of Great Game Animals (Ungulata) exhibited in the Department of Zoology, British Museum (Natural History). Second Edition. [By R. Lydekker, F.R.S.] Pp. 95. 53 Text and other figures. With list of Horns, Antlers and Tusks, and Index. 1913, 8vo. 1*s*.
- Guide to the Elephants (Recent and Fossil) exhibited in the Department of Geology and Palæontology in the British Museum (Natural History). [By Dr. C. W. Andrews, F.R.S.] Illustrated by 31 text-figures. Pp. 46. 1908, 8vo. 6*d*.
- Guide to the Specimens of the Horse Family (Equidæ) exhibited in the Department of Zoology, British Museum (Natural History). [By R. Lydekker, F.R.S.] Pp. 42. 26 Figures. 1907, 8vo. 1*s*.
- Guide to the Domesticated Animals (other than Horses) exhibited in the Central and North Halls of the British Museum (Natural History). Second Edition. [By R. Lydekker, F.R.S.] Illustrated by 25 Figures. Pp. 56. [With table of Contents, List of Illustrations, and Index.] 1912, 8vo. 6*d*.
- Guide to the Whales, Porpoises, and Dolphins (order Cetacea) exhibited in the Department of Zoology, British Museum (Natural History). [By R. Lydekker, F.R.S.] Illustrated by 33 Figures. Pp. 47. [With Index.] 1909, 8vo. 4*d*.
- Guide to the Gallery of Birds in the Department of Zoology, British Museum (Natural History). [By W. R. Ogilvie Grant.] Second Edition. Pp. iv., 228. 25 Plates, and 7 Illustrations in text. [With Index.] 1910, 4to. 2*s*. 6*d*

Guide to the Gallery of Birds in the Department of Zoology British Museum (Natural History). [By W. R. Ogilvie Grant.] :—

Part I. General Series. Pp. 149. [With Index.] 1905, 4to. 6*d*.

Part II. Nesting Series of British Birds. Second Edition. Pp. 62. 4 Plates. [Index.] 1909, 4to. 4*d*.

Guide to the Gallery of Reptilia and Amphibia in the Department of Zoology of the British Museum (Natural History). [By R. Lydekker, F.R.S.] Illustrated by 76 text and other Figures. Second Edition. Pp. iv., 85. [With Table of Contents and Index.] 1913, 8vo. 1*s*.

Guide to the Gallery of Fishes in the Department of Zoology of the British Museum (Natural History). [By Dr. W. G. Ridewood.] Illustrated by 96 Figures. Pp. v., 209. [With Preface by Sir E. Ray Lankester, Table of Classification, and Index.] 1908, 8vo. 1*s*.

Guide to the British Vertebrates Exhibited in the Department of Zoology, British Museum (Natural History). [By W. P. Pycraft.] Pp. iv., 122. 26 Text-Figures, 1 Plan. [With Index.] 1910, 8vo. 1*s*.

Guide to the exhibited series of Insects in the Department of Zoology, British Museum (Natural History). By C. O. Waterhouse. Second Edition. Pp. 65 : 62 text- and full-page Illustrations. [With Table of Contents and Index.] 1909, 8vo. 1*s*.

Guide to the Crustacea, Arachnida, Onychophora and Myriopoda exhibited in the Department of Zoology, British Museum (Natural History). (By W. T. Calman, D.Sc., A. S. Hirst, and F. J. Bell.) Pp. 133 : 90 Text-Figures. [With Table of Contents and Index.] 1910, 8vo. 1*s*.

Guide to the Shell and Starfish Galleries (Mollusca, Polyzoa, Brachiopoda, Tunicata, Echinoderma, and Worms), Department of Zoology, British Museum (Natural History). Fifth Edition. Pp. iv., 133. 125 Woodcuts, Plan, and Indexes. 1908, 8vo. 6*d*.

Guide to the Coral Gallery (Protozoa, Porifera or Sponges, Hydrozoa, and Anthozoa) in the Department of Zoology, British Museum (Natural History). Second Edition. Pp. [iv., 8] 73. 90 Illustrations, Plan, and Index. 1907, 8vo. 1*s*.

A Guide to the Fossil Mammals and Birds in the Department of Geology and Palaeontology in the British Museum (Natural History). Ninth Edition. [By A. S. Woodward, LL.D., F.R.S.] Pp. xvi., 100. 6 Plates, 88 Text-Figures. [With List of Illustrations, Table of Stratified Rocks, and Index.] 1909, 8vo. 6*d*.

- A Guide to the Fossil Reptiles, Amphibians, and Fishes in the Department of Geology and Palæontology in the British Museum (Natural History). Ninth Edition. [By A. S. Woodward, LL.D., F.R.S.] Pp. xviii., 110. 8 Plates and 116 Text-Figures. [With Table of Contents, Lists of Illustrations, Geological Time-Scale, and Index.] 1910, 8vo. 9*d*.
- A Guide to the Fossil Invertebrate Animals in the Department of Geology and Palæontology in the British Museum (Natural History). [By F. A. Bather, D.Sc., F.R.S.] Second Edition. Pp. x., 183. 7 Plates and 96 Text-Figures. [With List of Illustrations, Geological Time-scale, and Index.] 1911, 8vo. 1*s*.
- A Guide to the Mineral Gallery of the British Museum (Natural History). Eleventh Edition. Pp. 32. Plan. 1911, 8vo. 1*d*.
- The Student's Index to the Collection of Minerals, British Museum (Natural History). Twenty-fourth Edition. Pp. 36. [With a Plan of the Mineral Gallery.] 1911, 8vo. 2*d*.
- An Introduction to the Study of Minerals, with a Guide to the Mineral Gallery of the British Museum (Natural History). By L. Fletcher, M.A., F.R.S. Thirteenth Edition. Pp. 123. 41 Woodcuts. [With Plan of the Mineral Gallery and Index.] 1910, 8vo. 6*d*.
- An Introduction to the Study of Rocks and Guide to the Museum Collection. Fourth Edition. By L. Fletcher, M.A., F.R.S. Pp. 155. [With Plan of the Mineral Gallery, Table of Contents, and Index.] 1909, 8vo. 1*s*.
- An Introduction to the Study of Meteorites, with a List of the Meteorites represented in the Collection. By L. Fletcher, M.A., F.R.S., &c. Tenth Edition. Pp. 120. [With a Plan of the Mineral Gallery, and an Index to the Meteorites represented in the Collection.] 1908, 8vo. 6*d*.
- List of British Seed-plants and Ferns exhibited in the Department of Botany, British Museum (Natural History). [By A. B. Rendle, D.Sc., F.R.S., and J. Britten, F.L.S.] With table of Sequence of Orders, and Index of Genera. Pp. 44. 1907, 8vo. 4*d*.
- Guide to Sowerby's Models of British Fungi in the Department of Botany, British Museum (Natural History). Second Edition, revised. By Worthington G. Smith, F.L.S. Pp. 85. 91 Woodcuts. [With Table of Diagnostic Characters, Glossary, and Index.] 1908, 8vo. 4*d*.

*Guide to Mr. Worthington Smith's Drawings of Field and Cultivated Mushrooms, and Poisonous or Worthless Fungi, often mistaken for Mushrooms, exhibited in the Department of Botany. British Museum (Natural History). Pp. 24: 2 Plates, containing 28 coloured figures; 4 text-Figures. 1910, 8vo., 1s.

Guide to the British Mycetoza exhibited in the Department of Botany, British Museum (Natural History). [By Arthur Lister, F.R.S.] Third Edition, revised. Pp. 49. 46 Woodcuts. Index. 1909, 8vo. 3d.

SPECIAL GUIDES.

No. 2.—Books and Portraits illustrating the History of Plant Classification exhibited in the Department of Botany. Second Edition. [By A. B. Rendle, M.A., D.Sc., F.R.S.] Pp. 19. 4 Plates. 1909, 8vo. 4d.

No. 4.—Memorials of Charles Darwin: a Collection of Manuscripts, Portraits, Medals, Books, and Natural History Specimens to commemorate the Centenary of his Birth and the Fiftieth Anniversary of the Publication of "The Origin of Species." (Second Edition.) [By W. G. Ride-wood, D.Sc.] Pp. vi., 50. 2 Plates. 1910, 8vo. 6d.

No. 5.—Guide to the Exhibition of Animals, Plants, and Minerals mentioned in the Bible. Second Edition. Pp. vii., 78. 7 Text-figures. [With Index.] 1911, 8vo. 6d.

No. 6.—Guide to the Exhibition of Specimens illustrating the modification of the Structure of Animals in relation to Flight. Pp. viii., 80. 1 Plate, 44 Text-figures. [With List of Illustrations and Index.] 1913, 8vo. 6d.

INSTRUCTIONS FOR COLLECTORS.

Handbook of Instructions for Collectors, issued by the British Museum (Natural History). With Illustrations. Third Edition. Pp. 144. Index. 1906, 8vo. 1s. 6d.

Instructions for Collectors:—

No. 1.—Mammals. Fourth Edition. Pp. 8. Text illust. 1912, 8vo. 3d.

No. 2.—Birds and their Eggs. Fifth Edition. Pp. 13. 6 Text-figures. 1912, 8vo. 3d.

* The plates may be had separately in one sheet mounted on linen and varnished. Price 1s., or 1s. 2d. post free.

Instructions for Collectors—*continued*.

- No. 3.—Reptiles, Batrachians, and Fishes. [Third Edition.] Pp. 12. 1903, 8vo. 4*d*.
No. 4.—Insects. Fifth Edition. Pp. 11. Text illust. 1911, 8vo. 3*d*.
No. 5.—Diptera (Two-winged Flies). Third Edition. Pp. 16. Text illust. 1908, 8vo. 3*d*.
No. 6.—Mosquitoes (Culicidæ). [Third Edition.] Pp. 8. 1 Plate, 1 figure in text. 1904, 8vo. 3*d*.
No. 7.—Blood-sucking Flies, Ticks, &c. By E. E. Austen. Third Edition. Pp. 24 : 13 figures in text. 1907, 8vo. 3*d*.
No. 8.—Spiders, Centipedes, &c. Second Edition. Pp. 4. 1906, 8vo. 3*d*.
No. 9.—Soft-bodied and other Invertebrate Animals; Shells of Molluscs. Third Edition. Pp. 18. 1909, 8vo. 3*d*.
No. 10.—Plants. Fourth Edition. Pp. 10 : 3 figures in text. 1909, 8vo. 3*d*.
No. 11.—Fossils and Minerals. Fourth Edition. Pp. 8. 1913, 8vo. 3*d*.

ECONOMIC SERIES.

- No. 1.—The House-Fly as a Danger to Health. Its Life-history, and how to deal with it. By Ernest E. Austen. Second Edition. Pp. 12 : 2 plates (containing 4 figures), and 3 figures in text. 1913, 8vo. 1*d*.

British Museum (Natural History),
Cromwell Road,
London, S.W.
November, 1913.